

Tatsunori Hanai

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1340027/publications.pdf>

Version: 2024-02-01

38
papers

1,516
citations

430442

18
h-index

344852

36
g-index

39
all docs

39
docs citations

39
times ranked

1982
citing authors

#	ARTICLE	IF	CITATIONS
1	Handgrip strength stratifies the risk of covert and overt hepatic encephalopathy in patients with cirrhosis. <i>Journal of Parenteral and Enteral Nutrition</i> , 2022, 46, 858-866.	1.3	20
2	Proposal for new sleep disorder criteria in patients with chronic liver disease: Influence of liver-related complications. <i>Hepatology Research</i> , 2022, 52, 364-370.	1.8	7
3	Survival benefit of L-carnitine supplementation in patients with cirrhosis. <i>Journal of Parenteral and Enteral Nutrition</i> , 2022, , .	1.3	2
4	Zinc deficiency predicts overt hepatic encephalopathy and mortality in liver cirrhosis patients with minimal hepatic encephalopathy. <i>Hepatology Research</i> , 2021, 51, 662-673.	1.8	21
5	Increased Visceral Adipose Tissue and Hyperinsulinemia Raise the Risk for Recurrence of Non-B Non-C Hepatocellular Carcinoma after Curative Treatment. <i>Cancers</i> , 2021, 13, 1542.	1.7	9
6	Proposal of Stroop test cutoff values as screening for neuropsychological impairments in cirrhosis: A Japanese multicenter study. <i>Hepatology Research</i> , 2021, 51, 674-681.	1.8	11
7	Usefulness of the Stroop Test in Diagnosing Minimal Hepatic Encephalopathy and Predicting Overt Hepatic Encephalopathy. <i>Hepatology Communications</i> , 2021, 5, 1518-1526.	2.0	10
8	Nutritional assessment tool for predicting sarcopenia in chronic liver disease. <i>JCSM Rapid Communications</i> , 2021, 4, 150-158.	0.6	2
9	Utility of the SARC-F Questionnaire for Sarcopenia Screening in Patients with Chronic Liver Disease: A Multicenter Cross-Sectional Study in Japan. <i>Journal of Clinical Medicine</i> , 2021, 10, 3448.	1.0	10
10	Usefulness of nutritional therapy recommended in the Japanese Society of Gastroenterology/Japan Society of Hepatology evidence-based clinical practice guidelines for liver cirrhosis 2020. <i>Journal of Gastroenterology</i> , 2021, 56, 928-937.	2.3	7
11	Higher Accumulation of Visceral Adipose Tissue Is an Independent Risk Factor for Hepatocellular Carcinoma among Viral Hepatitis Patients with Non-Cirrhotic Livers. <i>Cancers</i> , 2021, 13, 5980.	1.7	1
12	Effect of l-carnitine supplementation on muscle cramps induced by stroke: A case report. <i>Nutrition</i> , 2020, 71, 110638.	1.1	5
13	Response to letter to the editor regarding: "Effect of l-carnitine supplementation on muscle cramps induced by stroke: A case report". <i>Nutrition</i> , 2020, 75-76, 110800.	1.1	0
14	Usefulness of Carnitine Supplementation for the Complications of Liver Cirrhosis. <i>Nutrients</i> , 2020, 12, 1915.	1.7	33
15	Rapid Depletion of Subcutaneous Adipose Tissue during Sorafenib Treatment Predicts Poor Survival in Patients with Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 1795.	1.7	15
16	Sustained virological response by direct-acting antivirals reduces the recurrence risk of hepatitis C-related hepatocellular carcinoma after curative treatment. <i>Molecular and Clinical Oncology</i> , 2020, 12, 111-116.	0.4	16
17	Development of diffuse large B-cell lymphoma after sofosbuvir/ledipasvir treatment for chronic hepatitis C: A case report and literature review. <i>Molecular and Clinical Oncology</i> , 2020, 13, 1.	0.4	8
18	Reduced handgrip strength is predictive of poor survival among patients with liver cirrhosis: A sex-stratified analysis. <i>Hepatology Research</i> , 2019, 49, 1414-1426.	1.8	51

#	ARTICLE	IF	CITATIONS
19	Rapid Depletions of Subcutaneous Fat Mass and Skeletal Muscle Mass Predict Worse Survival in Patients with Hepatocellular Carcinoma Treated with Sorafenib. <i>Cancers</i> , 2019, 11, 1206.	1.7	38
20	Homeostatic Model Assessment of Insulin Resistance for Predicting the Recurrence of Hepatocellular Carcinoma after Curative Treatment. <i>International Journal of Molecular Sciences</i> , 2019, 20, 605.	1.8	18
21	Prognostic significance of minimal hepatic encephalopathy in patients with liver cirrhosis in Japan: A propensity score-matching analysis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2019, 34, 1809-1816.	1.4	17
22	Effect of loop diuretics on skeletal muscle depletion in patients with liver cirrhosis. <i>Hepatology Research</i> , 2019, 49, 82-95.	1.8	31
23	Prevalence of Sarcopenia and Its Relationship with Nutritional State and Quality of Life in Patients with Digestive Diseases. <i>Journal of Nutritional Science and Vitaminology</i> , 2018, 64, 445-453.	0.2	22
24	Increased visceral fat volume raises the risk for recurrence of hepatocellular carcinoma after curative treatment. <i>Oncotarget</i> , 2018, 9, 14058-14067.	0.8	15
25	Sarcopenia predicts minimal hepatic encephalopathy in patients with liver cirrhosis. <i>Hepatology Research</i> , 2017, 47, 1359-1367.	1.8	78
26	Sarcopenia Impairs Prognosis of Patients with Hepatocellular Carcinoma: The Role of Liver Functional Reserve and Tumor-Related Factors in Loss of Skeletal Muscle Volume. <i>Nutrients</i> , 2017, 9, 1054.	1.7	44
27	Rapid skeletal muscle wasting predicts worse survival in patients with liver cirrhosis. <i>Hepatology Research</i> , 2016, 46, 743-751.	1.8	138
28	Impact of serum glycosylated <i>Wisteria floribunda</i> agglutinin positive Mac-2 binding protein levels on liver functional reserves and mortality in patients with liver cirrhosis. <i>Hepatology Research</i> , 2015, 45, 1083-1090.	1.8	35
29	Skeletal Muscle Depletion Predicts the Prognosis of Patients with Hepatocellular Carcinoma Treated with Sorafenib. <i>International Journal of Molecular Sciences</i> , 2015, 16, 9612-9624.	1.8	56
30	Efficacy and safety of cisplatin versus miriplatin in transcatheter arterial chemoembolization and transarterial infusion chemotherapy for hepatocellular carcinoma: A randomized controlled trial. <i>Hepatology Research</i> , 2015, 45, 514-522.	1.8	18
31	Sarcopenia impairs prognosis of patients with liver cirrhosis. <i>Nutrition</i> , 2015, 31, 193-199.	1.1	321
32	Skeletal muscle depletion is an independent prognostic factor for hepatocellular carcinoma. <i>Journal of Gastroenterology</i> , 2015, 50, 323-332.	2.3	205
33	Impact of Serum Chemerin Levels on Liver Functional Reserves and Platelet Counts in Patients with Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2014, 15, 11294-11306.	1.8	26
34	Pharmaceutical and nutraceutical approaches for preventing liver carcinogenesis: Chemoprevention of hepatocellular carcinoma using acyclic retinoid and branched-chain amino acids. <i>Molecular Nutrition and Food Research</i> , 2014, 58, 124-135.	1.5	8
35	Free fatty acid as a marker of energy malnutrition in liver cirrhosis. <i>Hepatology Research</i> , 2014, 44, 218-228.	1.8	28
36	Branched-Chain Amino Acids Prevent Hepatocarcinogenesis and Prolong Survival of Patients With Cirrhosis. <i>Clinical Gastroenterology and Hepatology</i> , 2014, 12, 1012-1018.e1.	2.4	84

#	ARTICLE	IF	CITATIONS
37	Hepatocellular carcinoma patients with increased oxidative stress levels are prone to recurrence after curative treatment: a prospective case series study using the d-ROM test. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 845-852.	1.2	43
38	Nutritional status and quality of life in current patients with liver cirrhosis as assessed in 2007â€“2011. <i>Hepatology Research</i> , 2013, 43, 106-112.	1.8	63