Yoshihisa Matsukawa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1352166/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The presence of intraductal carcinoma of the prostate in needle biopsy is a significant prognostic factor for prostate cancer patients with distant metastasis at initial presentation. Modern Pathology, 2016, 29, 166-173.	5.5	59
2	Efficacy of Silodosin for Relieving Benign Prostatic Obstruction: Prospective Pressure Flow Study. Journal of Urology, 2009, 182, 2831-2835.	0.4	38
3	De novo detrusor underactivity after laparoscopic radical prostatectomy. International Journal of Urology, 2010, 17, 643-648.	1.0	37
4	Urodynamic Evaluation of the Efficacy of Mirabegron on Storage and Voiding Functions in Women With Overactive Bladder. Urology, 2015, 85, 786-790.	1.0	30
5	De novo overactive bladder after robotâ€assisted laparoscopic radical prostatectomy. Neurourology and Urodynamics, 2018, 37, 2008-2014.	1.5	27
6	Effects of tadalafil on storage and voiding function in patients with male lower urinary tract symptoms suggestive of benign prostatic hyperplasia: A urodynamicâ€based study. International Journal of Urology, 2018, 25, 246-250.	1.0	23
7	Clinical characteristics and useful signs to differentiate detrusor underactivity from bladder outlet obstruction in men with nonâ€neurogenic lower urinary tract symptoms. International Journal of Urology, 2020, 27, 47-52.	1.0	23
8	Efficacy of Silodosin for Relieving Benign Prostatic Obstruction: Prospective Pressure Flow Study. Journal of Urology, 2013, 189, S117-21.	0.4	20
9	Longâ€term efficacy of a combination therapy with an anticholinergic agent and an α1â€blocker for patients with benign prostatic enlargement complaining both voiding and overactive bladder symptoms: A randomized, prospective, comparative trial using a urodynamic study. Neurourology and Urodynamics. 2017. 36. 748-754.	1.5	18
10	Effects of dutasteride on storage and voiding symptoms in male patients with lower urinary tract symptoms as a result of benign prostatic obstruction: The 1â€year outcomes from a prospective urodynamic study. International Journal of Urology, 2014, 21, 826-830.	1.0	17
11	Laparoscopic versus open radical prostatectomy: Urodynamic evaluation of vesicourethral function. International Journal of Urology, 2009, 16, 393-396.	1.0	16
12	Effects of Withdrawing α1-Blocker from Combination Therapy with α1-Blocker and 5α-Reductase Inhibitor in Patients with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia: A Prospective and Comparative Trial Using Urodynamics. Journal of Urology, 2017, 198, 905-912.	0.4	15
13	Comparison in the efficacy of fesoterodine or mirabegron addâ€on therapy to silodosin for patients with benign prostatic hyperplasia complicated by overactive bladder: A randomized, prospective trial using urodynamic studies. Neurourology and Urodynamics, 2019, 38, 941-949.	1.5	15
14	Clinical efficacy of a loading dose of naftopidil for patients with benign prostate hyperplasia. World Journal of Urology, 2011, 29, 225-231.	2.2	14
15	Effect of Tamsulosin on Bladder Microcirculation in Rat Model of Bladder Outlet Obstruction Using Pencil Lens Charge-coupled Device Microscopy System. Urology, 2013, 81, 155-159.	1.0	14
16	Effect of Naftopidil on Bladder Microcirculation in a Rat Model of Bladder Outlet Obstruction. LUTS: Lower Urinary Tract Symptoms, 2017, 9, 111-116.	1.3	14
17	Comparison of Silodosin and Naftopidil for Efficacy in the Treatment of Benign Prostatic Enlargement Complicated by Overactive Bladder: A Randomized, Prospective Study (SNIPER Study). Journal of Urology, 2017, 197, 452-458.	0.4	14
18	Role of microglia in the spinal cord in colonâ€ŧoâ€bladder neural crosstalk in a rat model of colitis. Neurourology and Urodynamics, 2018, 37, 1320-1328.	1.5	14

#	Article	IF	CITATIONS
19	Clinical diagnostic criteria for detrusor underactivity: A report from the Japanese Continence Society working group on underactive bladder. LUTS: Lower Urinary Tract Symptoms, 2021, 13, 13-16.	1.3	14
20	Effect of Tamsulosin on Bladder Microcirculation in a Rat Ischemia-reperfusion Model, Evaluated by Pencil Lens Charge-coupled Device Microscopy System. Urology, 2010, 76, 1266.e1-1266.e5.	1.0	13
21	What are the factors contributing to failure in improvement of subjective symptoms following silodosin administration in patients with benign prostatic hyperplasia? investigation using a pressureâ€flow study. Neurourology and Urodynamics, 2013, 32, 266-270.	1.5	13
22	Variability of postâ€void residual urine volume and bladder voiding efficiency in patients with underactive bladder. LUTS: Lower Urinary Tract Symptoms, 2021, 13, 51-55.	1.3	13
23	Objective impacts of tadalafil on storage and voiding function in male patients with benign prostatic hyperplasia: 1-year outcomes from a prospective urodynamic study. World Journal of Urology, 2019, 37, 867-872.	2.2	12
24	Causative factors for de novo inguinal hernia after robot-assisted radical prostatectomy. Journal of Robotic Surgery, 2018, 12, 277-282.	1.8	11
25	Intravesical prostatic protrusion can predict therapeutic response to silodosin in male patients with lower urinary tract symptoms. International Journal of Urology, 2017, 24, 454-459.	1.0	10
26	What are the predicting factors for the therapeutic effects of dutasteride in male patients with lower urinary tract symptoms? Investigation using a urodynamic study. Neurourology and Urodynamics, 2017, 36, 1809-1815.	1.5	7
27	Investigation of the relationship between bladder function and sarcopenia using pressure flow studies in elderly male patients. Neurourology and Urodynamics, 2019, 38, 1417-1422.	1.5	7
28	A small pilot study to evaluate the accuracy and feasibility of a novel automated voiding diary device for recording urine output measurements. Neurourology and Urodynamics, 2021, 40, 272-277.	1.5	7
29	Feasibility of the Lilium α-200 portable ultrasound bladder scanner for accurate bladder volume measurement. Investigative and Clinical Urology, 2020, 61, 613.	2.0	7
30	A Slow Stream Is Pathophysiologically Related to a Poor Response to α1-Adrenoceptor Therapy in the Treatment of Storage Symptoms Associated With Benign Prostatic Hyperplasia. Urology, 2015, 86, 558-564.	1.0	6
31	Comparison of cernitin pollen extract vs tadalafil therapy for refractory chronic prostatitis/chronic pelvic pain syndrome: A randomized, prospective study. Neurourology and Urodynamics, 2020, 39, 1994-2002.	1.5	6
32	Two-year follow up of silodosin on lower urinary tract functions and symptoms in patients with benign prostatic hyperplasia based on prostate size: a prospective investigation using urodynamics. Therapeutic Advances in Urology, 2018, 10, 263-272.	2.0	5
33	Pre-treatment serum testosterone level can be a useful factor to predict the improvement in bladder outlet obstruction by tadalafil for male patients with lower urinary tract symptoms induced by benign prostatic obstruction. Aging Male, 2020, 23, 641-647.	1.9	4
34	Effects of tadalafil versus silodosin on voiding function in male patients with nonâ€neurogenic detrusor underactivity: A comparative study using propensity score matching. International Journal of Urology, 2021, 28, 411-416.	1.0	4
35	Effect of chemotherapy on survival for intraductal carcinoma of the prostate (IDC-P) in patients with castration-resistant prostate cancer Journal of Clinical Oncology, 2017, 35, e601-e601.	1.6	4
36	Associations among Physician–Patient Communication, Patient Satisfaction, and Clinical Effectiveness of Overactive Bladder Medication: A Survey of Patients with Overactive Bladder. Journal of Clinical Medicine, 2022, 11, 4087.	2.4	4

Yoshihisa Matsukawa

#	Article	IF	CITATIONS
37	Urodynamic evaluation before and after to RARP to identify pre and intraoperative factors affecting postoperative continence. Neurourology and Urodynamics, 2021, 40, 1147-1153.	1.5	3
38	Development of an artificial intelligence diagnostic system for lower urinary tract dysfunction in men. International Journal of Urology, 2021, 28, 1143-1148.	1.0	3
39	The Change of Testosterone Secretion During the Treatment of Alpha-1 Blocker in Patients with Benign Prostatic Hyperplasia. Urology, 2016, 88, 149-154.	1.0	2
40	What Are Factors Contributing to Improvement of Overactive Bladder Symptoms after Alpha-1 Blocker Treatment in Patients with Both Storage and Voiding Symptoms?. Urologia Internationalis, 2019, 103, 439-443.	1.3	2
41	Factors contributing to the efficacy of two addâ€on therapies of fesoterodine or mirabegron to silodosin monotherapy for persistent overactive bladder in men with lower urinary tract symptoms. International Journal of Urology, 2020, 27, 85-86.	1.0	2
42	Useful parameters to predict the presence of detrusor overactivity in male patients with lower urinary tract symptoms. Neurourology and Urodynamics, 2020, 39, 1394-1400.	1.5	2
43	Daily urine loss immediately after urethral catheter removal may be an effective predictor of longâ€term urinary incontinence following robotâ€assisted laparoscopic radical prostatectomy. International Journal of Clinical Practice, 2021, 75, e13736.	1.7	2
44	Validation and clinical utility of the diagnostic criteria for detrusor underactivity in men proposed by the Japanese Continence Society. International Journal of Urology, 2022, 29, 595-596.	1.0	2
45	Clinical features and urodynamic findings in elderly men with chronic prostatitis. International Journal of Urology, 2022, , .	1.0	1
46	Reply. Urology, 2015, 86, 563-564.	1.0	0
47	The clinical benefit of sequential therapy with androgen receptor axisâ€ŧargeted agents alone in patients with castrationâ€resistant prostate cancer: A propensity scoreâ€matched comparison study. Prostate, 2020, 80, 1373-1380.	2.3	0
48	Retrospective analysis of post-TKI everolimus therapy for metastatic renal cell carcinoma using the doubling-time of tumor growth Journal of Clinical Oncology, 2014, 32, e15574-e15574.	1.6	0