
## Mari Kobayashi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1360910/publications.pdf Version: 2024-02-01



| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Foraging activity of harbour porpoises around a bottom-gillnet in a coastal fishing ground, under the risk of bycatch. PLoS ONE, 2021, 16, e0246838.   | 2.5 | 3         |
| 2  | Morphological identification in skull between spotted seal and harbor seal using geometric morphometrics. Journal of Morphology, 2021, 282, 1455-1465.   | 1.2 | 3         |
| 3  | Intraspecific differences in the diet of Kuril harbor seals (Phoca vitulina stejnegeri) in Erimo,<br>Hokkaido, using DNA barcoding diet analysis. Mammal Research, 2021, 66, 553-563.  | 1.3 | Ο         |
| 4  | Notocotylus ikutai n. sp. (Digenea: Notocotylidae) from lymnaeid snails and anatid birds in Hokkaido,<br>Japan. Parasitology International, 2021, 83, 102318.  | 1.3 | 3         |
| 5  | Distribution and abundance of dalli-type Dall's porpoises Phocoenoides dalli migrating into waters off southeastern Hokkaido, Japan, during summer: results of 2014–2016 aerial surveys. Fisheries Science, 2020, 86, 287-298.     | 1.6 | Ο         |
| 6  | Current population genetics of Japanese harbor seals: Two distinct populations found within a small area. Marine Mammal Science, 2020, 36, 915-924.  | 1.8 | 5         |
| 7  | One-sided infections by intestinal parasites in two sympatric porpoises bycaught from the Nemuro<br>Strait of Hokkaido, Japan. Parasitology International, 2020, 77, 102118.   | 1.3 | Ο         |
| 8  | Infection status of commercial fish with cystacanth larvae of the genus Corynosoma<br>(Acanthocephala: Polymorphidae) in Hokkaido, Japan. International Journal of Food Microbiology,<br>2019, 305, 108256.                        | 4.7 | 19        |
| 9  | Dependency of Japanese harbor seals ( <i>Phoca vitulina</i> ) on salmon set nets at Cape Erimo,<br>Hokkaido, Japan. Marine Mammal Science, 2019, 35, 58-71.  | 1.8 | 2         |
| 10 | Surveillance of amyloidosis in stranded and bycaught cetaceans off Hokkaido, Japan. Journal of<br>Veterinary Medical Science, 2019, 81, 897-902.   | 0.9 | 8         |
| 11 | Host characteristics and infection level of an intestinal parasite Corynosoma strumosum<br>(Acanthocephala) in the Kuril harbor seal of Erimo Cape, Hokkaido, Japan. Parasitology International,<br>2018, 67, 237-244.             | 1.3 | 8         |
| 12 | Mitochondrial DNA reveals secondary contact in Japanese harbour seals, the southernmost population in the western Pacific. PLoS ONE, 2018, 13, e0191329.   | 2.5 | 13        |
| 13 | Seasonal and Spatial Occurrence of Northern Fur Seals <i>Callorhinus ursinus</i> Around Northern<br>Japan. Mammal Study, 2017, 42, 51-56.  | 0.6 | 3         |
| 14 | Brown adipose tissue expresses uncoupling protein 1 in newborn harbor seals ( <i>Phoca vitulina</i> ).<br>Marine Mammal Science, 2015, 31, 818-827.  | 1.8 | 3         |
| 15 | Stable isotope ratios of carbon, nitrogen and oxygen in killer whales ( Orcinus orca ) stranded on the coast of Hokkaido, Japan. Marine Pollution Bulletin, 2014, 86, 238-243.   | 5.0 | 6         |
| 16 | Genetic Variation in the Harbor Seal ( <i>Phoca vitulina</i> ) and Spotted Seal ( <i>Phoca largha</i> )<br>Around Hokkaido, Japan, Based on Mitochondrial Cytochrome <i>b</i> Sequences. Zoological Science,<br>2010, 27, 263-268. | 0.7 | 17        |
| 17 | Growth variation in skull morphology of Kuril harbor seals (Phoca vitulina stejnegeri) and spotted<br>seals (Phoca largha) in Hokkaido, Japan. Japanese Journal of Veterinary Research, 2009, 57, 147-62.                          | 0.7 | 2         |