

Ireneia Melo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1377660/publications.pdf>

Version: 2024-02-01

20
papers

257
citations

1163117
8
h-index

940533
16
g-index

20
all docs

20
docs citations

20
times ranked

578
citing authors

#	ARTICLE	IF	CITATIONS
1	New record of Hyphoderma ettriae (Basidiomycota) in Portugal and its combination in Lawryomyces. Phytotaxa, 2020, 472, 193-200.	0.3	1
2	Considerations and consequences of allowing DNA sequence data as types of fungal taxa. IMA Fungus, 2018, 9, 167-175.	3.8	45
3	Corticoid fungi (Basidiomycota) from the Biosphere Reserve of Arganeraie, Morocco: a preliminary survey. Nova Hedwigia, 2016, 103, 193-210.	0.4	1
4	New national and regional bryophyte records, 44. Journal of Bryology, 2015, 37, 228-241.	1.2	32
5	Spelling out Jaapia species. Mycological Progress, 2015, 14, 1.	1.4	6
6	Sistotremastrum guttuliferum: a new species from the Macaronesian islands. Mycological Progress, 2013, 12, 687-692.	1.4	8
7	Molecular analyses confirm Brevicellicium in Trechisporales. IMA Fungus, 2013, 4, 21-28.	3.8	23
8	Regional annotated mycobiotas new to the Mycotaxon website. Mycotaxon, 2013, 123, 491-492.	0.3	13
9	<i>Steccherinum straminellum</i>, a new record for Italy. Mycotaxon, 2013, 121, 281-284.	0.3	0
10	<i>Gloeocystidiellum kenyense</i> in Azores and Madeira. Mycotaxon, 2012, 119, 337-343.	0.3	4
11	Amanita poisonings resulting in acute, reversible renal failure: new cases, new toxic Amanita mushrooms. Nephrology Dialysis Transplantation, 2012, 27, 1380-1386.	0.7	45
12	Peniophora aluticolor (Fungi, Basidiomycota), an orphaned species restudied. Nova Hedwigia, 2012, 94, 437-440.	0.4	4
13	Morphological and molecular studies of Hyphodermella in the Western Mediterranean area. Mycological Progress, 2010, 9, 585-596.	1.4	18
14	A re-evaluation of <i>Hypochnicium</i> (Polyporales) based on morphological and molecular characters. Mycologia, 2010, 102, 1426-1436.	1.9	24
15	Diversity and richness of corticioid fungi (Basidiomycota) on Azores Islands: a preliminary survey. Nova Hedwigia, 2009, 88, 285-308.	0.4	7
16	The aphyllophorales (<I>Basidiomycota</I>) of a Mediterranean biodiversity "hotspot" â€" Cazorla, Segura & Las Villas" Natural Park (Spain). Mycotaxon, 2009, 109, 465-468.	0.3	1
17	<i>Lagarobasidium calongei</i> (Aphyllophorales, Basidiomycota), a new species of corticioid fungi from Azores Islands. Anales Del Jardin Botanico De Madrid, 2009, 66, 41-46.	0.4	10
18	<i>Gloedontia xerophila</i> (Aphyllophorales, Basidiomycota), a new species with corticioid basidioma from the Canary Islands. Mycologia, 2008, 100, 673-676.	1.9	8

#	ARTICLE	IF	CITATIONS
19	A new species of <i>Hyphodermella</i> (Basidiomycetes, Aphylophorales) from Portugal. <i>Nova Hedwigia</i> , 2003, 77, 351-355.	0.4	4
20	<i>Australohydnum dregeanum</i> (Basidiomycetes, Stereaceae) in Europe. <i>Nova Hedwigia</i> , 2002, 74, 527-532.	0.4	3