

Fumitaka Koga

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1383414/fumitaka-koga-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

159
papers

3,497
citations

32
h-index

53
g-index

224
ext. papers

3,944
ext. citations

3
avg. IF

4.98
L-index

#	Paper	IF	Citations
159	Molecular chaperone TRAP1 regulates a metabolic switch between mitochondrial respiration and aerobic glycolysis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E1604-12	11.5	175
158	A small molecule cell-impermeant Hsp90 antagonist inhibits tumor cell motility and invasion. <i>Oncogene</i> , 2008 , 27, 2478-87	9.2	140
157	Diagnostic performance of diffusion-weighted magnetic resonance imaging in bladder cancer: potential utility of apparent diffusion coefficient values as a biomarker to predict clinical aggressiveness. <i>European Radiology</i> , 2011 , 21, 2178-86	8	123
156	Impaired p63 expression associates with poor prognosis and uroplakin III expression in invasive urothelial carcinoma of the bladder. <i>Clinical Cancer Research</i> , 2003 , 9, 5501-7	12.9	107
155	Sarcopenia as a prognostic biomarker of advanced urothelial carcinoma. <i>PLoS ONE</i> , 2015 , 10, e0115895	3.7	100
154	Hsp90 inhibition transiently activates Src kinase and promotes Src-dependent Akt and Erk activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 11318-22	11.5	98
153	Impact of C-reactive protein kinetics on survival of patients with metastatic renal cell carcinoma. <i>European Urology</i> , 2009 , 55, 1145-53	10.2	92
152	Initial experience of diffusion-weighted magnetic resonance imaging to assess therapeutic response to induction chemoradiotherapy against muscle-invasive bladder cancer. <i>Urology</i> , 2010 , 75, 387-91	1.6	85
151	Role of diffusion-weighted magnetic resonance imaging in predicting sensitivity to chemoradiotherapy in muscle-invasive bladder cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 83, e21-7	4	82
150	Prognostic Significance of Sarcopenia in Patients with Metastatic Renal Cell Carcinoma. <i>Journal of Urology</i> , 2016 , 195, 26-32	2.5	78
149	Inhibition of cancer invasion and metastasis by targeting the molecular chaperone heat-shock protein 90. <i>Anticancer Research</i> , 2009 , 29, 797-807	2.3	77
148	Long-term outcome of bladder papillary urothelial neoplasms of low malignant potential. <i>BJU International</i> , 2003 , 92, 559-62	5.6	73
147	Apparent diffusion coefficient value reflects invasive and proliferative potential of bladder cancer. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 172-8	5.6	71
146	Development, validation, and head-to-head comparison of logistic regression-based nomograms and artificial neural network models predicting prostate cancer on initial extended biopsy. <i>European Urology</i> , 2008 , 54, 601-11	10.2	69
145	High diagnostic ability of multiparametric magnetic resonance imaging to detect anterior prostate cancer missed by transrectal 12-core biopsy. <i>Journal of Urology</i> , 2013 , 190, 867-73	2.5	64
144	Impaired Delta Np63 expression associates with reduced beta-catenin and aggressive phenotypes of urothelial neoplasms. <i>British Journal of Cancer</i> , 2003 , 88, 740-7	8.7	59
143	Usefulness of pre-biopsy multiparametric magnetic resonance imaging and clinical variables to reduce initial prostate biopsy in men with suspected clinically localized prostate cancer. <i>Journal of Urology</i> , 2013 , 190, 502-8	2.5	58

142	C-reactive protein level predicts prognosis in patients with muscle-invasive bladder cancer treated with chemoradiotherapy. <i>BJU International</i> , 2008 , 101, 978-81	5.6	58
141	New three-dimensional head-mounted display system, TMDU-S-3D system, for minimally invasive surgery application: procedures for gasless single-port radical nephrectomy. <i>International Journal of Urology</i> , 2012 , 19, 886-9	2.3	53
140	Potential role of Hsp90 inhibitors in overcoming cisplatin resistance of bladder cancer-initiating cells. <i>International Journal of Cancer</i> , 2012 , 131, 987-96	7.5	53
139	Low dose geldanamycin inhibits hepatocyte growth factor and hypoxia-stimulated invasion of cancer cells. <i>Cell Cycle</i> , 2007 , 6, 1393-402	4.7	52
138	Risk factors for the development of bladder transitional cell carcinoma following surgery for transitional cell carcinoma of the upper urinary tract. <i>Urologia Internationalis</i> , 2001 , 67, 135-41	1.9	52
137	Diffusion-weighted magnetic resonance imaging in the differentiation of angiomyolipoma with minimal fat from clear cell renal cell carcinoma. <i>International Journal of Urology</i> , 2011 , 18, 727-30	2.3	50
136	Selective bladder preservation with curative intent for muscle-invasive bladder cancer: a contemporary review. <i>International Journal of Urology</i> , 2012 , 19, 388-401	2.3	47
135	Heat shock protein 90 targeting therapy: state of the art and future perspective. <i>EXCLI Journal</i> , 2015 , 14, 48-58	2.4	46
134	Selective bladder-sparing protocol consisting of induction low-dose chemoradiotherapy plus partial cystectomy with pelvic lymph node dissection against muscle-invasive bladder cancer: oncological outcomes of the initial 46 patients. <i>BJU International</i> , 2012 , 109, 860-6	5.6	42
133	Loss of DeltaNp63alpha promotes invasion of urothelial carcinomas via N-cadherin/Src homology and collagen/extracellular signal-regulated kinase pathway. <i>Cancer Research</i> , 2009 , 69, 9263-70	10.1	41
132	RhoA is associated with invasion and lymph node metastasis in upper urinary tract cancer. <i>BJU International</i> , 2003 , 91, 234-8	5.6	41
131	Gasless single-port access endoscopic surgery in urology: minimum incision endoscopic surgery, MIES. <i>International Journal of Urology</i> , 2009 , 16, 791-800	2.3	39
130	Effects of intravenous administration of high dose-diethylstilbestrol diphosphate on serum hormonal levels in patients with hormone-refractory prostate cancer. <i>Endocrine Journal</i> , 1999 , 46, 659-64	2.9	38
129	Apparent diffusion coefficient value as a biomarker reflecting morphological and biological features of prostate cancer. <i>International Urology and Nephrology</i> , 2014 , 46, 555-61	2.3	34
128	Risk stratification for bladder recurrence of upper urinary tract urothelial carcinoma after radical nephroureterectomy. <i>BJU International</i> , 2015 , 115, 705-12	5.6	33
127	External validation of the Mayo Clinic cancer specific survival score in a Japanese series of clear cell renal cell carcinoma. <i>Journal of Urology</i> , 2008 , 180, 1290-5; discussion 1295-6	2.5	32
126	Prognostic significance of sarcopenia in upper tract urothelial carcinoma patients treated with radical nephroureterectomy. <i>Cancer Medicine</i> , 2016 , 5, 2213-20	4.8	31
125	Characteristics and clinical significance of prostate cancers missed by initial transrectal 12-core biopsy. <i>BJU International</i> , 2012 , 109, 665-71	5.6	31

124	Bladder Cancer Stem-Like Cells: Their Origin and Therapeutic Perspectives. <i>International Journal of Molecular Sciences</i> , 2015 , 17,	6.3	31
123	Favourable outcomes of patients with clinical stage T3N0M0 bladder cancer treated with induction low-dose chemo-radiotherapy plus partial or radical cystectomy vs immediate radical cystectomy: a single-institutional retrospective comparative study. <i>BJU International</i> , 2009 , 104, 189-94	5.6	30
122	Apparent diffusion coefficient as a prognostic biomarker of upper urinary tract cancer: a preliminary report. <i>European Radiology</i> , 2013 , 23, 2206-14	8	29
121	ErbB2 and NFB overexpression as predictors of chemoradiation resistance and putative targets to overcome resistance in muscle-invasive bladder cancer. <i>PLoS ONE</i> , 2011 , 6, e27616	3.7	29
120	Phase-II trial of combination treatment of interferon- γ /cimetidine, cyclooxygenase-2 inhibitor and renin-angiotensin-system inhibitor (I-CCA therapy) for advanced renal cell carcinoma. <i>Cancer Science</i> , 2011 , 102, 137-43	6.9	29
119	Serum concentration of type I collagen metabolites as a quantitative marker of bone metastases in patients with prostate carcinoma 1997 , 80, 1760-1767		29
118	Low-dose chemoradiotherapy followed by partial or radical cystectomy against muscle-invasive bladder cancer: an intent-to-treat survival analysis. <i>Urology</i> , 2008 , 72, 384-8	1.6	29
117	Diffusion-weighted magnetic resonance imaging in management of bladder cancer, particularly with multimodal bladder-sparing strategy. <i>World Journal of Radiology</i> , 2014 , 6, 344-54	2.9	28
116	Prognostic significance of endothelial Per-Arnt-sim domain protein 1/hypoxia-inducible factor-2 α expression in a subset of tumor associated macrophages in invasive bladder cancer. <i>Journal of Urology</i> , 2004 , 171, 1080-4	2.5	26
115	Low-dose Hsp90 inhibitors tumor-selectively sensitize bladder cancer cells to chemoradiotherapy. <i>Cell Cycle</i> , 2011 , 10, 4291-9	4.7	25
114	Combination of diffusion-weighted magnetic resonance imaging and extended prostate biopsy predicts lobes without significant cancer: application in patient selection for hemiablativ focal therapy. <i>European Urology</i> , 2014 , 65, 186-92	10.2	24
113	Significance of ERBB2 overexpression in therapeutic resistance and cancer-specific survival in muscle-invasive bladder cancer patients treated with chemoradiation-based selective bladder-sparing approach. <i>International Journal of Radiation Oncology Biology Physics</i> , 2014 , 90, 303-11	4	24
112	Postoperative Changes in Skeletal Muscle Mass Predict Survival of Patients With Metastatic Renal Cell Carcinoma Undergoing Cytoreductive Nephrectomy. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, e229-e238	3.3	24
111	Longitudinal change in renal function after radical nephrectomy in Japanese patients with renal cortical tumors. <i>Journal of Urology</i> , 2011 , 185, 2066-71	2.5	24
110	Perioperative complications of radical cystectomy after induction chemoradiotherapy in bladder-sparing protocol against muscle-invasive bladder cancer: a single institutional retrospective comparative study with primary radical cystectomy. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 1373-9	2.8	24
109	Impact of Sarcopenia as a Prognostic Biomarker of Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	24
108	Selective tetramodal bladder-preservation therapy, incorporating induction chemoradiotherapy and consolidative partial cystectomy with pelvic lymph node dissection for muscle-invasive bladder cancer: oncological and functional outcomes of 107 patients. <i>BJU International</i> , 2019 , 124, 242-250	5.6	23
107	Gasless laparoendoscopic single-port clampless sutureless partial nephrectomy for peripheral renal tumors: perioperative outcomes. <i>International Journal of Urology</i> , 2015 , 22, 349-55	2.3	22

106	Loss of uroplakin III expression is associated with a poor prognosis in patients with urothelial carcinoma of the upper urinary tract. <i>BJU International</i> , 2006 , 97, 1322-6	5.6	22
105	Impact of sarcopenia in the management of urological cancer patients. <i>Expert Review of Anticancer Therapy</i> , 2017 , 17, 455-466	3.5	21
104	Safety of transperineal 14-core systematic prostate biopsy in diabetic men. <i>International Journal of Urology</i> , 2009 , 16, 930-5	2.3	21
103	Young age as favorable prognostic factor for cancer-specific survival in localized renal cell carcinoma. <i>Urology</i> , 2011 , 77, 842-7	1.6	20
102	Antimicrobial prophylaxis is not necessary in clean category minimally invasive surgery for renal and adrenal tumors: a prospective study of 373 consecutive patients. <i>Urology</i> , 2012 , 80, 570-5	1.6	19
101	Equivalent survival and improved preservation of renal function after distal ureterectomy compared with nephroureterectomy in patients with urothelial carcinoma of the distal ureter: a propensity score-matched multicenter study. <i>International Journal of Urology</i> , 2014 , 21, 1098-104	2.3	18
100	Role of diffusion-weighted magnetic resonance imaging as an imaging biomarker of urothelial carcinoma. <i>International Journal of Urology</i> , 2014 , 21, 1190-200	2.3	18
99	Zoledronic acid sensitizes renal cell carcinoma cells to radiation by downregulating STAT1. <i>PLoS ONE</i> , 2013 , 8, e64615	3.7	18
98	High Ki-67 Expression Predicts Favorable Survival in Muscle-Invasive Bladder Cancer Patients Treated With Chemoradiation-Based Bladder-Sparing Protocol. <i>Clinical Genitourinary Cancer</i> , 2015 , 13, e243-e251	3.3	17
97	Discarding antimicrobial prophylaxis for transurethral resection of bladder tumor: a feasibility study. <i>International Journal of Urology</i> , 2009 , 16, 61-3	2.3	17
96	Epithelial-mesenchymal transition promotes SOX2 and NANOG expression in bladder cancer. <i>Laboratory Investigation</i> , 2017 ,	5.9	16
95	Heat shock factor 1 (HSF1)-targeted anticancer therapeutics: overview of current preclinical progress. <i>Expert Opinion on Therapeutic Targets</i> , 2019 , 23, 369-377	6.4	16
94	Renal function after radical nephrectomy: development and validation of predictive models in Japanese patients. <i>International Journal of Urology</i> , 2014 , 21, 238-42	2.3	15
93	Diabetes mellitus with obesity is a predictor of recurrence in patients with non-metastatic renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2013 , 43, 740-6	2.8	15
92	Negative p53/positive p21 immunostaining is a predictor of favorable response to chemotherapy in patients with locally advanced bladder cancer. <i>Japanese Journal of Cancer Research</i> , 2000 , 91, 416-23		15
91	Preoperative chronic kidney disease is predictive of oncological outcome of radical cystectomy for bladder cancer. <i>World Journal of Urology</i> , 2018 , 36, 249-256	4	15
90	Impact of bladder neck involvement on progression in patients with primary non-muscle invasive bladder cancer: a prospective validation study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 38.e29-36	2.8	14
89	Metabolic and Molecular Basis of Sarcopenia: Implications in the Management of Urothelial Carcinoma. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	12

88	Effect of diabetes mellitus on high-grade prostate cancer detection among Japanese obese patients with prostate-specific antigen less than 10 ng/mL. <i>Urology</i> , 2012 , 79, 1329-34	1.6	12
87	Spatial and isoform specific p63 expression in the male human urogenital tract. <i>Journal of Urology</i> , 2006 , 176, 2268-73	2.5	12
86	Impact of Immunohistochemistry-Based Subtypes in Muscle-Invasive Bladder Cancer on Response to Chemoradiation Therapy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2018 , 102, 1408-1416	4.16	12
85	Successful reduction of hospital-acquired methicillin-resistant <i>Staphylococcus aureus</i> in a urology ward: a 10-year study. <i>BMC Urology</i> , 2013 , 13, 35	2.2	11
84	840 NEW THREE-DIMENSIONAL HEAD-MOUNTED DISPLAY SYSTEM (ROBOSURGEON SYSTEM): APPLICATION TO THE INITIAL 80 CASES OF GASLESS SINGLE-PORT ACCESS UROLOGIC SURGERIES. <i>Journal of Urology</i> , 2013 , 189,	2.5	11
83	Impact of advanced age on biochemical recurrence after radical prostatectomy in Japanese men according to pathological stage. <i>Japanese Journal of Clinical Oncology</i> , 2013 , 43, 410-6	2.8	11
82	Higher expression of K-ras is associated with parathyroid hormone-related protein-induced hypercalcaemia in renal cell carcinoma. <i>BJU International</i> , 2001 , 88, 960-6	5.6	11
81	Increased expression of sialyl-Lewis A correlates with poor survival in upper urinary tract urothelial cancer patients. <i>Anticancer Research</i> , 2003 , 23, 3441-6	2.3	11
80	Posttherapeutic skeletal muscle mass recovery predicts favorable prognosis in patients with advanced urothelial carcinoma receiving first-line platinum-based chemotherapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 156.e9-156.e16	2.8	10
79	Effects of cholinesterase inhibition in supraspinal and spinal neural pathways on the micturition reflex in rats. <i>BJU International</i> , 2009 , 104, 1163-9	5.6	10
78	Favorable outcome of preoperative low dose chemoradiotherapy against muscle-invasive bladder cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2003 , 26, 504-7	2.7	10
77	Prognostic significance of the controlling nutritional status (CONUT) score in advanced urothelial carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 76.e11-76.e17	2.8	10
76	Ureteral Involvement Is Associated with Poor Prognosis in Upper Urinary Tract Urothelial Carcinoma Patients Treated by Nephroureterectomy: A Multicenter Database Study. <i>European Urology Focus</i> , 2016 , 2, 296-302	5.1	10
75	Candidate selection for quadrant-based focal ablation through a combination of diffusion-weighted magnetic resonance imaging and prostate biopsy. <i>BJU International</i> , 2016 , 117, 94-101	5.6	10
74	Detection of Muscle-Invasive Bladder Cancer on Biparametric MRI Using Vesical Imaging-Reporting and Data System and Apparent Diffusion Coefficient Values (VI-RADS/ADC). <i>Bladder Cancer</i> , 2020 , 6, 161-169	1	9
73	Significance of positive urine cytology on progression and cancer-specific mortality of non-muscle-invasive bladder cancer. <i>Clinical Genitourinary Cancer</i> , 2014 , 12, e87-93	3.3	9
72	Pathology-based risk stratification of muscle-invasive bladder cancer patients undergoing cystectomy for persistent disease after induction chemoradiotherapy in bladder-sparing approaches. <i>BJU International</i> , 2012 , 110, E203-8	5.6	9
71	Sensitivity to chemoradiation predicts development of metastasis in muscle-invasive bladder cancer patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1270-5	2.8	9

70	Prognostic significance of serum Γ glutamyltransferase in patients with advanced urothelial carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019 , 37, 108-115	2.8	9
69	Biomarkers for Predicting Clinical Outcomes of Chemoradiation-Based Bladder Preservation Therapy for Muscle-Invasive Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	9
68	Extended biopsy based criteria incorporating cumulative cancer length for predicting clinically insignificant prostate cancer. <i>BJU International</i> , 2012 , 110, E564-9	5.6	8
67	Pre-operative risk stratification for cancer-specific survival in patients with renal cell carcinoma with venous involvement who underwent nephrectomy. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 756-61	2.8	8
66	History of malignancy is a predictor of prostate cancer detection: incorporation into a pre-biopsy nomogram. <i>International Journal of Urology</i> , 2008 , 15, 1055-60	2.3	8
65	Endoscopic minilaparotomy partial nephrectomy for solitary renal cell carcinoma smaller than 4 cm. <i>Japanese Journal of Clinical Oncology</i> , 2002 , 32, 417-21	2.8	8
64	Prognostic Significance of the Controlling Nutritional Status (CONUT) Score in Patients with Advanced Renal Cell Carcinoma Treated with Nivolumab after Failure of Prior Tyrosine Kinase Inhibitors. <i>Journal of Urology</i> , 2020 , 204, 1166-1172	2.5	8
63	Prostate-specific antigen response to deferred combined androgen blockade therapy using bicalutamide predicts survival after subsequent oestrogen and docetaxel therapies in patients with castration-resistant prostate cancer. <i>BJU International</i> , 2012 , 110, 1149-55	5.6	7
62	Three-dimensional 26-core biopsy-based patient selection criteria for nerve-sparing radical prostatectomy. <i>International Journal of Urology</i> , 2008 , 15, 1061-6	2.3	7
61	Bimodal pattern of the impact of body mass index on cancer-specific survival of upper urinary tract urothelial carcinoma patients. <i>Anticancer Research</i> , 2014 , 34, 5683-8	2.3	7
60	Impact of Serum Γ Glutamyltransferase on Overall Survival in Patients with Metastatic Renal Cell Carcinoma in the Era of Targeted Therapy. <i>Targeted Oncology</i> , 2020 , 15, 347-356	5	6
59	Diagnostic performance of initial transperineal 14-core prostate biopsy to detect significant cancer. <i>International Urology and Nephrology</i> , 2013 , 45, 645-52	2.3	6
58	Female urethral diverticular abscess clearly depicted by diffusion-weighted magnetic resonance imaging. <i>International Journal of Urology</i> , 2008 , 15, 460-1	2.3	6
57	Favorable response to combination treatment of cimetidine, cyclooxygenase-2 inhibitor and renin-angiotensin system inhibitor in metastatic renal cell carcinoma: Report of three cases. <i>International Journal of Urology</i> , 2008 , 15, 848-50	2.3	6
56	A novel equation and nomogram including body weight for estimating prostate volumes in men with biopsy-proven benign prostatic hyperplasia. <i>Asian Journal of Andrology</i> , 2012 , 14, 703-7	2.8	6
55	Small cell carcinoma of the urinary bladder: a contemporary review with a special focus on bladder-sparing treatments. <i>Expert Review of Anticancer Therapy</i> , 2013 , 13, 1269-79	3.5	5
54	Deep vein thrombosis during chemotherapy in a patient with advanced testicular cancer: successful percutaneous thrombectomy under temporary placement of retrievable inferior vena cava filter. <i>International Journal of Urology</i> , 2001 , 8, 90-3	2.3	5
53	Multifocal Synchronous Upper Urinary Tract Carcinosarcoma (Sarcomatoid Carcinoma) With Rhabdomyoblastic Differentiation. <i>International Journal of Surgical Pathology</i> , 2019 , 27, 547-552	1.2	4

52	Prognostic significance of intensive local therapy to bone lesions in renal cell carcinoma patients with bone metastasis. <i>Clinical and Experimental Metastasis</i> , 2016 , 33, 699-705	4.7	4
51	Serum Γ Glutamyltransferase as a Prognostic Biomarker in Metastatic Castration-resistant Prostate Cancer Treated With Enzalutamide. <i>Anticancer Research</i> , 2019 , 39, 5773-5780	2.3	4
50	Bone abnormal signal incidentally found in pre-biopsy diffusion-weighted MRI for suspected prostate cancer: what does it reflect?. <i>Urologia Internationalis</i> , 2014 , 93, 170-5	1.9	4
49	Impact of lower urinary tract symptoms on prostate cancer risk among Japanese men with prostate-specific antigen . <i>International Journal of Urology</i> , 2013 , 20, 1163-8	2.3	4
48	A novel repeat biopsy nomogram based on three-dimensional extended biopsy. <i>Urology</i> , 2011 , 77, 915-20.6		4
47	Interactions between inducible nitric oxide synthase and cyclooxygenase-2 in response to ischaemia-reperfusion of rabbit bladder. <i>BJU International</i> , 2010 , 106, 716-22	5.6	4
46	Sarcomatoid renal cell carcinoma with scant carcinomatous components. <i>International Journal of Urology</i> , 2000 , 7, 58-60; discussion 61	2.3	4
45	Retroperitoneal Teratoma in an Adult: A Potential Pitfall in the Differential Diagnosis of Adrenal Myelolipoma. <i>Case Reports in Urology</i> , 2016 , 2016, 5141769	0.5	4
44	Who Can Avoid Systematic Biopsy Without Missing Clinically Significant Prostate Cancer in Men Who Undergo Magnetic Resonance Imaging-Targeted Biopsy?. <i>Clinical Genitourinary Cancer</i> , 2019 , 17, e664-e671	3.3	3
43	Diagnostic performance and safety of a three-dimensional 14-core systematic biopsy method. <i>BJU International</i> , 2015 , 115, 412-8	5.6	3
42	Collecting duct carcinoma with acquired cystic disease of the kidney in a long-term hemodialysis patient. <i>International Journal of Urology</i> , 2008 , 15, 93-5	2.3	3
41	Minimum incision endoscopic nephrectomy for giant hydronephrosis. <i>International Journal of Urology</i> , 2007 , 14, 774-6	2.3	3
40	A Systematic Review of Serum Γ Glutamyltransferase as a Prognostic Biomarker in Patients with Genitourinary Cancer. <i>Antioxidants</i> , 2021 , 10,	7.1	3
39	Standardization of the apparent diffusion coefficient value of bladder cancer across different centers: Applicability in predicting aggressive pathologic phenotypes. <i>Clinical Imaging</i> , 2017 , 44, 121-126 ²⁻⁷		2
38	PD13-06 FAVORABLE OUTCOMES OF GASLESS SINGLE-PORT CLAMPLESS PARTIAL NEPHRECTOMY USING NEW THREE DIMENSIONAL HEAD-MOUNTED DISPLAY SYSTEM (ROBOSURGEON SYSTEM). <i>Journal of Urology</i> , 2014 , 191,	2.5	2
37	Performance of prostate-specific antigen mass in estimation of prostate volume in Japanese men with benign prostate hyperplasia. <i>International Journal of Urology</i> , 2012 , 19, 929-35	2.3	2
36	Diagnostic contribution of C-reactive protein kinetics for gastric metastasis from renal cell carcinoma. <i>International Cancer Conference Journal</i> , 2012 , 1, 93-95	0.9	2
35	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Partial Nephrectomy 2015 , 43-64		2

34	Contact with renal sinus is associated with poor prognosis in surgically treated pT1 clear cell renal cell carcinoma. <i>International Journal of Urology</i> , 2020 , 27, 657-662	2.3	1
33	MP55-01 BLADDER-SPARING THERAPY IN MUSCLE-INVASIVE BLADDER CANCER BY INDUCTION CHEMORADIOTHERAPY PLUS PARTIAL CYSTECTOMY WITH PELVIC LYMPH NODE DISSECTION: ASSESSMENTS OF QUALITY OF LIFE AND URINARY FUNCTIONS. <i>Journal of Urology</i> , 2014 , 191,	2.5	1
32	PD17-07 IMPACT OF BLADDER NECK INVOLVEMENT ON PROGRESSION IN PATIENTS WITH PRIMARY NON-MUSCLE-INVASIVE BLADDER CANCER: A MULTICENTER VALIDATION STUDY. <i>Journal of Urology</i> , 2015 , 193,	2.5	1
31	Editorial comment from Dr Koga to local control rate and prognosis after sequential chemoradiation for small cell carcinoma of the bladder. <i>International Journal of Urology</i> , 2013 , 20, 785-6	2.3	1
30	49 THE ABNORMALITY OF GLUCOSE TOLERANCE TEST IN PHEOCHROMOCYTOMA: ONE OF THE SIGNIFICANT FINDINGS OF POSTOPERATIVE HYPOGLYCEMIA. <i>Journal of Urology</i> , 2013 , 189,	2.5	1
29	MP55-09 SELECTIVE BLADDER-SPARING PROTOCOL CONSISTING OF INDUCTION LOW-DOSE CHEMORADIOTHERAPY AND CONSOLIDATIVE PARTIAL CYSTECTOMY IN MUSCLE INVASIVE BLADDER CANCER PATIENTS WITH SYNCHRONOUS OR METACHRONOUS MULTIFOCALITY. <i>Journal of Urology</i> , 2014 , 191,	2.5	1
28	MP8-09 ANTIMICROBIAL PROPHYLAXIS IS AVOIDABLE IN MINIMALLY INVASIVE NEPHROURETERECTOMY WITH NEGATIVE URINE CULTURE. <i>Journal of Urology</i> , 2014 , 191,	2.5	1
27	369 COMBINATION OF 14-CORE BIOPSY AND MAGNETIC RESONANCE IMAGING CAN IDENTIFY APPROPRIATE CANDIDATES FOR HEMIABLATIVE FOCAL THERAPY OF PROSTATE CANCER. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
26	1105 GASLESS SINGLE-PORT CLAMPLESS PARTIAL NEPHRECTOMY FOR PERIPHERAL RENAL TUMOR SURGICAL, SHORT-TERM ONCOLOGICAL, AND FUNCTIONAL OUTCOMES. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
25	Gasless Single Port Surgery for Renal Cell Carcinoma: Minimum Incision Endoscopic Surgery 2012 ,		1
24	506 LOW INCIDENCE OF BENIGN PATHOLOGIC LESIONS AT NEPHRECTOMY FOR RENAL MASSES PRESUMED TO BE STAGE T1N0M0 RENAL CELL CARCINOMA IN JAPANESE PATIENTS: INFLUENCE OF SEX, AGE AND TUMOR SIZE. <i>Journal of Urology</i> , 2010 , 183,	2.5	1
23	1172 ANTIBIOTIC PROPHYLAXIS IS NOT NECESSARY IN MINIMALLY INVASIVE SURGERY FOR RENAL AND ADRENAL TUMORS: A PROSPECTIVE STUDY OF 301 CONSECUTIVE PATIENTS. <i>Journal of Urology</i> , 2011 , 185,	2.5	1
22	False tumor marker surge evoked by peripheral blood stem cell transplantation. <i>Oncologist</i> , 2008 , 13, 526-9	5.7	1
21	Congestion of the corpus spongiosum and necrosis of the glans penis in systemic vasculitis. <i>BJU International</i> , 1996 , 78, 796-7	5.6	1
20	Renal Angiomyolipoma Mimicking a Well-Differentiated Retroperitoneal Liposarcoma. <i>Case Reports in Urology</i> , 2020 , 2020, 8812057	0.5	1
19	Jejunal limb obstruction by a tumor thrombus from pancreatic metastasis of renal cell carcinoma: a case report. <i>Surgical Case Reports</i> , 2021 , 7, 37	0.8	1
18	A Case of Renal Pelvic Cancer Complicated by Horseshoe Kidney Treated with RoboSurgeon Gasless Single-Port Retroperitoneoscopic Nephroureterectomy. <i>Case Reports in Urology</i> , 2018 , 2018, 3231480	0.5	1
17	Who Can Avoid Biopsy of Magnetic Resonance Imaging-Negative Lobes without Compromising Significant Cancer Detection among Men with Unilateral Magnetic Resonance Imaging-Positive Lobes?. <i>Urologia Internationalis</i> , 2021 , 105, 386-393	1.9	0

16	Apparent Diffusion Coefficient Value as a Biomarker for Detecting Muscle-Invasive and High-Grade Bladder Cancer: A Systematic Review. <i>Applied Sciences (Switzerland)</i> , 2022 , 12, 1278	2.6	○
15	Preoperative models incorporating the systemic immune-inflammation index for predicting prognosis and muscle invasion in patients with non-metastatic upper tract urothelial carcinoma. <i>International Journal of Clinical Oncology</i> , 2021 , 1	4.2	○
14	The Controlling Nutritional Status (CONUT) Score is a Prognostic Biomarker in Advanced Urothelial Carcinoma Patients Treated with First-Line Platinum-Based Chemotherapy. <i>Bladder Cancer</i> , 2021 , 7, 13-21	1	○
13	Novel anatomical apical dissection utilizing puboprostatic "open-collar" technique: Impact on apical surgical margin and early continence recovery. <i>PLoS ONE</i> , 2021 , 16, e0249991	3.7	○
12	Impact of fluorodeoxyglucose uptake on positron emission tomography/computed tomography on chemosensitivity and survival in patients with metastatic urothelial carcinoma. <i>International Journal of Urology</i> , 2019 , 26, 820-826	2.3	
11	Clinically node-positive micropapillary bladder cancer in a young female desiring to spare functional bladder and fertility. <i>International Cancer Conference Journal</i> , 2020 , 9, 151-154	0.9	
10	A case of metastatic testicular germ-cell tumor with rhabdomyosarcoma successfully treated with germ-cell tumor-oriented chemotherapy. <i>International Cancer Conference Journal</i> , 2012 , 1, 58-62	0.9	
9	Editorial Comment to Cancer death from non-muscle invasive bladder cancer: Report of the Japanese Urological Association of data from the 1999-2001 registry in Japan. <i>International Journal of Urology</i> , 2010 , 17, 913	2.3	
8	IFN γ enhances the cytotoxicity of 5-fluorouracil against human bladder cancer cells through induction of PD-ECGF / TP. <i>Japanese Journal of Urology</i> , 2002 , 93, 323		
7	Platelet-derived endothelial cell growth factor/thymidine phosphorylase expression in primary bladder cancer. <i>Japanese Journal of Urology</i> , 2002 , 93, 377		
6	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Radical Nephrectomy 2015 , 23-42		
5	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Radical Prostatectomy 2015 , 105-126		
4	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Pelvic Lymph Node Dissection and Inguinal Hernia Prevention 2015 , 159-173		
3	Contribution of C-reactive protein to predicting overall survival in patients with advanced urothelial carcinoma: Development of a nomogram and evaluation by decision curve analysis.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 280-280	2.2	
2	Functional CT and MRI of the Urinary System and Adrenal Glands 2014 , 1173-1182		
1	Prognostic significance of the controlling nutritional status score in patients with metastatic renal cell carcinoma diagnosed before an era of first-line immune-oncology combination therapies. <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 1570-1576	2.8	