
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1384096/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Current Concepts in the Etiology and Treatment of Behçet Disease. Survey of Ophthalmology, 2005, 50, 297-350. | 1.7 | 374 |
| 2 | Serum levels of TNF-α, sIL-2R, IL-6, and IL-8 are increased and associated with elevated lipid peroxidation in patients with Behçet's disease. Mediators of Inflammation, 2002, 11, 87-93. | 1.4 | 201 |
| 3 | Comparative External Versus Endoscopic Dacryocystorhinostomy: Results in 115 Patients (130 Eyes). Otolaryngology - Head and Neck Surgery, 2000, 123, 488-491. | 1.1 | 186 |
| 4 | Comparison of secondary implantation of flexible open-loop anterior chamber and scleral-fixated posterior chamber intraocular lenses. Journal of Cataract and Refractive Surgery, 2003, 29, 301-308. | 0.7 | 137 |
| 5 | The effect of reading and near-work on the development of myopia in emmetropic boys: a prospective, controlled, three-year follow-up study. Vision Research, 2001, 41, 2511-2520. | 0.7 | 107 |
| 6 | Increased nitric oxide production in patients with Behçet's disease: Is it a new activity marker?. Journal of the American Academy of Dermatology, 2002, 46, 50-54. | 0.6 | 98 |
| 7 | Nitric oxide and lipid peroxidation are increased and associated with decreased antioxidant enzyme activities in patients with age-related macular degeneration. Documenta Ophthalmologica, 2003, 106, 129-136. | 1.0 | 94 |
| 8 | Central corneal thickness is lower in osteogenesis imperfecta and negatively correlates with the presence of blue sclera. Ophthalmic and Physiological Optics, 2002, 22, 511-515. | 1.0 | 74 |
| 9 | Caffeic acid phenethyl ester to inhibit posterior capsule opacification in rabbits. Journal of Cataract and Refractive Surgery, 1997, 23, 1572-1576. | 0.7 | 62 |
| 10 | Clinical features of Behcet's disease in children. International Journal of Dermatology, 2006, 45, 713-716. | 0.5 | 62 |
| 11 | Audio-vestibular evaluation in patients with Behçet's syndrome. Journal of Laryngology and Otology, 2001, 115, 704-8. | 0.4 | 56 |
| 12 | Adenosine deaminase enzyme activity is increased and negatively correlates with catalase, superoxide dismutase and glutathione peroxidase in patients with Behçet's disease: original contributions/clinical and laboratory investigations. Mediators of Inflammation, 2003, 12, 107-116. | 1.4 | 56 |
| 13 | Managing the symptoms of Behçet's disease. Expert Opinion on Pharmacotherapy, 2004, 5, 317-328. | 0.9 | 49 |
| 14 | Autonomic nervous system functions in obese children. Brain and Development, 2000, 22, 151-153. | 0.6 | 46 |
| 15 | Increased Plasma Adrenomedullin Levels in Patients with Behçet's Disease. Dermatology, 2000, 201, 312-315. | 0.9 | 45 |
| 16 | Ocular Behçet disease. Current Opinion in Ophthalmology, 2011, 22, 508-516. | 1.3 | 45 |
| 17 | Use of caffeic acid phenethyl ester to prevent sodium-selenite-induced cataract in rat eyes. Journal of Cataract and Refractive Surgery, 2002, 28, 1457-1462. | 0.7 | 44 |
| 18 | Ingestion of IH636 grape seed proanthocyanidin extract to prevent selenite-induced oxidative stress in experimental cataract. Journal of Cataract and Refractive Surgery, 2006, 32, 1041-1045. | 0.7 | 39 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Craniofacial Anthropometry in a Turkish Population. Cleft Palate-Craniofacial Journal, 2002, 39, 208-218. | 0.5 | 38 |
| 20 | Weill-Marchesani syndrome in three generations. Eye, 1999, 13, 773-777. | 1.1 | 35 |
| 21 | Defective visual field tests in chronic heavy smokers. Acta Ophthalmologica, 2001, 79, 53-56. | 0.4 | 30 |
| 22 | Decreased Central Corneal Thickness in Children With Down Syndrome. Journal of Pediatric Ophthalmology and Strabismus, 2002, 39, 274-277. | 0.3 | 30 |
| 23 | Adenosine deaminase enzyme levels, their relation with disease activity, and the effect of colchicine on adenosine deaminase levels in patients with Behçet's disease. Rheumatology International, 2005, 25, 452-456. | 1.5 | 26 |
| 24 | Synovial nitric oxide concentrations are increased and correlated with serum levels in patients with active Behçet's disease: a pilot study. Clinical Rheumatology, 2005, 24, 324-330. | 1.0 | 25 |
| 25 | Lenticular astigmatism in tilted disc syndrome. Journal of Cataract and Refractive Surgery, 2002, 28, 1836-1840. | 0.7 | 22 |
| 26 | Pattern VEP is a useful technique in monitoring the effectiveness of occlusion therapy in amblyopic eyes under occlusion therapy. Documenta Ophthalmologica, 2004, 109, 223-227. | 1.0 | 21 |
| 27 | Effect of gestational nicotine treatment on newborn rat retina: a histopathological and morphometric analysis. Ophthalmic and Physiological Optics, 2003, 23, 527-533. | 1.0 | 19 |
| 28 | Hammer-Chisel Technique in Endoscopic Dacryocystorhinostomy. Annals of Otology, Rhinology and Laryngology, 2003, 112, 444-449. | 0.6 | 19 |
| 29 | Serum Levels of Soluble P-Selectin Are Increased and Associated With Disease Activity in Patients With Behçet's Syndrome. Mediators of Inflammation, 2005, 2005, 237-241. | 1.4 | 19 |
| 30 | The pathophysiological significance of red blood cell nitric oxide concentrations in inflammatory BehA§et's disease. Mediators of Inflammation, 2003, 12, 255-256. | 1.4 | 16 |
| 31 | Determination of Nontoxic Concentrations of Piperacillin/Tazobactam for Intravitreal Application. Ophthalmic Research, 2004, 36, 139-144. | 1.0 | 16 |
| 32 | Aqueous Humor Adrenomedullin Levels Differ in Patients With Different Types of Glaucoma. Japanese Journal of Ophthalmology, 2002, 46, 203-208. | 0.9 | 15 |
| 33 | Urinary adrenomedullin levels are increased and correlated with plasma concentrations in patients with Behcet's syndrome. International Journal of Urology, 2002, 9, 296-303. | 0.5 | 15 |
| 34 | Pattern electroretinogram for monitoring the efficacy of intravitreal triamcinolone injection in diabetic macular edema. Documenta Ophthalmologica, 2004, 109, 139-145. | 1.0 | 14 |
| 35 | Invasive Squamous Cell Carcinoma of the Conjunctiva First Misdiagnosed as a Pterygium: A Clinicopathologic Case Report. Eye and Contact Lens, 2008, 34, 188-190. | 0.8 | 14 |
| 36 | The Efficacy of Piperacillin/Tazobactam in Experimental Pseudomonas aeruginosa Endophthalmitis: A Histopathological and Microbiological Evaluation. Current Eye Research, 2005, 30, 13-19. | 0.7 | 13 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | â€~Kite-tail' fascia lata strips technique: frontalis suspension using a non-endoscopic minimally invasive single-thigh incision approach. British Journal of Ophthalmology, 2012, 96, 570-575. | 2.1 | 11 |
| 38 | Urinary nitric oxide levels are increased and correlated with plasma concentrations in patients with Behcet's disease: Is it a new urinary activity marker?. Nephrology, 2003, 8, 231-238. | 0.7 | 10 |
| 39 | Figure-of-Eight Vertical Mattress Suture Technique for Anterior Flap Suspension to Overlying Tissues in External Dacryocystorhinostomy. American Journal of Ophthalmology, 2007, 143, 328-333.e1. | 1.7 | 10 |
| 40 | Interpupillary index: a new parameter for hypo-hypertelorism. Journal of Cranio-Maxillo-Facial Surgery, 2001, 29, 191-194. | 0.7 | 9 |
| 41 | Do Non-Steroidal Anti-Inflammatory Drugs Delay Posterior Capsule Opacification After Phacoemulsification in Children? A Randomized, Prospective Controlled Trial. Current Eye Research, 2011, 36, 1139-1147. | 0.7 | 9 |
| 42 | mRNA Expression Level of Interleukin Genes in the Determining Phases of Behçet's Disease. Annals of Dermatology, 2015, 27, 291. | 0.3 | 9 |
| 43 | Deep Sclerectomy with Various Implants: An Experimental and Histopathologic Study in a Rabbit Model. Ophthalmologica, 2004, 218, 264-269. | 1.0 | 8 |
| 44 | Normative Values of Craniofacial Measurements in Idiopathic Benign Macrocephalic Children. Cleft Palate-Craniofacial Journal, 2001, 38, 260-263. | 0.5 | 7 |
| 45 | Shifting crystalline pseudohypopyon secondary to lens absorption with spontaneous openings in the anterior lens capsule. Journal of Cataract and Refractive Surgery, 2006, 32, 1971-1973. | 0.7 | 7 |
| 46 | Diagnostic dilemma between intestinal Behçet disease and inflammatory bowel disease with pyoderma gangrenosum. World Journal of Gastroenterology, 2006, 12, 5748. | 1.4 | 7 |
| 47 | The Efficacy of Intravitreal Piperacillin/Tazobactam in Rabbits with Experimental <i>Staphylococcus epidermidis</i> Endophthalmitis: A Comparison with Vancomycin. Ophthalmic Research, 2005, 37, 168-174. | 1.0 | 6 |
| 48 | The migration pattern, patient selection with diagnostic methodological flaw and confusing naming dilemma in Behçet disease. European Journal of Echocardiography, 2007, 8, 167-173. | 2.3 | 6 |
| 49 | Primary Probing with and without Monoka Silastic Stent Intubation for Epiphora in Older Children and Adults. Current Eye Research, 2020, 45, 87-90. | 0.7 | 6 |
| 50 | Bilateral epibulbar and posterior scleral choristomas in linear nevus sebaceous syndrome. Acta Ophthalmologica, 2003, 81, 312-314. | 0.4 | 5 |
| 51 | Excessive maternal caffeine exposure during pregnancy is cataractogenic for neonatal crystalline lenses in rats: a biomicroscopic and histopathologic study. Acta Ophthalmologica, 2004, 82, 552-556. | 0.4 | 5 |
| 52 | High dose of caffeine administered to pregnant rats causes histopathological changes in the cornea of newborn pups. Medical Science Monitor, 2003, 9, BR168-73. | 0.5 | 5 |
| 53 | Tc-99m HMPAO brain SPECT in linear nevus sebaceous syndrome. Annals of Nuclear Medicine, 2003, 17, 703-706. | 1.2 | 4 |
| 54 | Tumour necrosis factor α, lipid peroxidation and NO• are increased and associated with decreased free-radical scavenging enzymes in patients with Weill-Marchesani syndrome. Mediators of Inflammation, 2004, 13, 165-170. | 1.4 | 4 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | A comparison of intravitreal piperacillin/tazobactam with ceftazidime in experimental Pseudomonas aeruginosa endophthalmitis. Experimental Eye Research, 2005, 80, 361-367. | 1.2 | 4 |
| 56 | Author's response:. Survey of Ophthalmology, 2006, 51, 174-177. | 1.7 | 4 |
| 57 | Decreased central corneal thickness in children with Down syndrome. Journal of Pediatric Ophthalmology and Strabismus, 2002, 39, 274-7. | 0.3 | 4 |
| 58 | Frontalis suspension by a minimally invasive "harvesting-stripping technique―for congenital blepharoptosis in children under 3-years-old. European Journal of Ophthalmology, 2023, 33, 161-170. | 0.7 | 3 |
| 59 | Regarding neutrophil and lymphocyte responses to oralStreptococcusin Adamantiades–Behçet's disease. FEMS Immunology and Medical Microbiology, 2006, 47, 311-314. | 2.7 | 2 |
| 60 | Internuclear ophthalmoplegia as a symptom associated with Henoch-Schoenlein purpura. Neuro-Ophthalmology, 2002, 26, 193-196. | 0.4 | 1 |
| 61 | HLA-genotyping and the historical naming process of an old disorder. Jornal Vascular Brasileiro, 0, 21, | 0.1 | 1 |
| 62 | A novel one-step repair of large lower eyelid defects following tumor resection in adults: surgical technique and outcome. European Journal of Plastic Surgery, 0, , . | 0.3 | 1 |
| 63 | Breakage of the Ritleng Tip Within the Interosseous Nasolacrimal Ductus During Monoka Silastic Stent Intubation and its Management. Journal of Maxillofacial and Oral Surgery, 0, , 1. | 0.6 | 0 |