

Cem Evereklioglu

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1384096/cem-evereklioglu-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

1,851
citations

21
h-index

42
g-index

63
ext. papers

2,057
ext. citations

2.9
avg, IF

4.73
L-index

#	Paper	IF	Citations
60	Current concepts in the etiology and treatment of Behçet disease. <i>Survey of Ophthalmology</i> , 2005 , 50, 297-350	6.1	295
59	Serum levels of TNF-alpha, sIL-2R, IL-6, and IL-8 are increased and associated with elevated lipid peroxidation in patients with Behçet's disease. <i>Mediators of Inflammation</i> , 2002 , 11, 87-93	4.3	163
58	Comparative external versus endoscopic dacryocystorhinostomy: results in 115 patients (130 eyes). <i>Otolaryngology - Head and Neck Surgery</i> , 2000 , 123, 488-91	5.5	154
57	Comparison of secondary implantation of flexible open-loop anterior chamber and scleral-fixated posterior chamber intraocular lenses. <i>Journal of Cataract and Refractive Surgery</i> , 2003 , 29, 301-8	2.3	113
56	The effect of reading and near-work on the development of myopia in emmetropic boys: a prospective, controlled, three-year follow-up study. <i>Vision Research</i> , 2001 , 41, 2511-20	2.1	91
55	Increased nitric oxide production in patients with Behçet's disease: is it a new activity marker?. <i>Journal of the American Academy of Dermatology</i> , 2002 , 46, 50-4	4.5	86
54	Nitric oxide and lipid peroxidation are increased and associated with decreased antioxidant enzyme activities in patients with age-related macular degeneration. <i>Documenta Ophthalmologica</i> , 2003 , 106, 129-36	2.2	71
53	Central corneal thickness is lower in osteogenesis imperfecta and negatively correlates with the presence of blue sclera. <i>Ophthalmic and Physiological Optics</i> , 2002 , 22, 511-5	4.1	60
52	Clinical features of Behçet's disease in children. <i>International Journal of Dermatology</i> , 2006 , 45, 713-6	1.7	55
51	Caffeic acid phenethyl ester to inhibit posterior capsule opacification in rabbits. <i>Journal of Cataract and Refractive Surgery</i> , 1997 , 23, 1572-6	2.3	51
50	Adenosine deaminase enzyme activity is increased and negatively correlates with catalase, superoxide dismutase and glutathione peroxidase in patients with Behçet's disease: original contributions/clinical and laboratory investigations. <i>Mediators of Inflammation</i> , 2003 , 12, 107-16	4.3	46
49	Increased plasma adrenomedullin levels in patients with Behçet's disease. <i>Dermatology</i> , 2000 , 201, 312-5	4.4	44
48	Audio-vestibular evaluation in patients with Behçet's syndrome. <i>Journal of Laryngology and Otology</i> , 2001 , 115, 704-8	1.8	44
47	Use of caffeic acid phenethyl ester to prevent sodium-selenite-induced cataract in rat eyes. <i>Journal of Cataract and Refractive Surgery</i> , 2002 , 28, 1457-62	2.3	39
46	Managing the symptoms of Behçet's disease. <i>Expert Opinion on Pharmacotherapy</i> , 2004 , 5, 317-28	4	38
45	Ingestion of IH636 grape seed proanthocyanidin extract to prevent selenite-induced oxidative stress in experimental cataract. <i>Journal of Cataract and Refractive Surgery</i> , 2006 , 32, 1041-5	2.3	35
44	Autonomic nervous system functions in obese children. <i>Brain and Development</i> , 2000 , 22, 151-3	2.2	32

43	Ocular Behçet disease: current therapeutic approaches. <i>Current Opinion in Ophthalmology</i> , 2011 , 22, 508-16	5.1	28
42	Weill-Marchesani syndrome in three generations. <i>Eye</i> , 1999 , 13 (Pt 6), 773-7	4.4	28
41	Defective visual field tests in chronic heavy smokers. <i>Acta Ophthalmologica</i> , 2001 , 79, 53-6		22
40	Adenosine deaminase enzyme levels, their relation with disease activity, and the effect of colchicine on adenosine deaminase levels in patients with Behçet's disease. <i>Rheumatology International</i> , 2005 , 25, 452-6	3.6	21
39	Craniofacial Anthropometry in a Turkish Population. <i>Cleft Palate-Craniofacial Journal</i> , 2002 , 39, 208-218	1.9	19
38	Lenticular astigmatism in tilted disc syndrome. <i>Journal of Cataract and Refractive Surgery</i> , 2002 , 28, 1836-40	2.4	19
37	Pattern VEP is a useful technique in monitoring the effectiveness of occlusion therapy in amblyopic eyes under occlusion therapy. <i>Documenta Ophthalmologica</i> , 2004 , 109, 223-7	2.2	18
36	Synovial nitric oxide concentrations are increased and correlated with serum levels in patients with active Behçet's disease: a pilot study. <i>Clinical Rheumatology</i> , 2005 , 24, 324-30	3.9	18
35	Decreased Central Corneal Thickness in Children With Down Syndrome. <i>Journal of Pediatric Ophthalmology and Strabismus</i> , 2002 , 39, 274-277	0.9	18
34	Hammer-chisel technique in endoscopic dacryocystorhinostomy. <i>Annals of Otology, Rhinology and Laryngology</i> , 2003 , 112, 444-9	2.1	15
33	Urinary adrenomedullin levels are increased and correlated with plasma concentrations in patients with Behçet's syndrome. <i>International Journal of Urology</i> , 2002 , 9, 296-303	2.3	15
32	The pathophysiological significance of red blood cell nitric oxide concentrations in inflammatory Behçet's disease. <i>Mediators of Inflammation</i> , 2003 , 12, 255-6	4.3	14
31	Determination of nontoxic concentrations of piperacillin/tazobactam for intravitreal application. An electroretinographic, histopathologic and morphometric analysis. <i>Ophthalmic Research</i> , 2004 , 36, 139-44	2.9	14
30	Pattern electroretinogram for monitoring the efficacy of intravitreal triamcinolone injection in diabetic macular edema. <i>Documenta Ophthalmologica</i> , 2004 , 109, 139-45	2.2	14
29	Effect of gestational nicotine treatment on newborn rat retina: a histopathological and morphometric analysis. <i>Ophthalmic and Physiological Optics</i> , 2003 , 23, 527-33	4.1	14
28	Serum levels of soluble P-selectin are increased and associated with disease activity in patients with Behçet's syndrome. <i>Mediators of Inflammation</i> , 2005 , 2005, 237-41	4.3	14
27	The efficacy of piperacillin/tazobactam in experimental <i>Pseudomonas aeruginosa</i> endophthalmitis: a histopathological and microbiological evaluation. <i>Current Eye Research</i> , 2005 , 30, 13-9	2.9	12
26	Aqueous humor adrenomedullin levels differ in patients with different types of glaucoma. <i>Japanese Journal of Ophthalmology</i> , 2002 , 46, 203-8	2.6	10

25	Kite-tailFascia lata strips technique: frontalis suspension using a non-endoscopic minimally invasive single-thigh incision approach. <i>British Journal of Ophthalmology</i> , 2012 , 96, 570-5	5.5	9
24	Invasive squamous cell carcinoma of the conjunctiva first misdiagnosed as a pterygium: a clinicopathologic case report. <i>Eye and Contact Lens</i> , 2008 , 34, 188-90	3.2	9
23	Urinary nitric oxide levels are increased and correlated with plasma concentrations in patients with Behçet's disease: is it a new urinary activity marker?. <i>Nephrology</i> , 2003 , 8, 231-8	2.2	9
22	Do non-steroidal anti-inflammatory drugs delay posterior capsule opacification after phacoemulsification in children? A randomized, prospective controlled trial. <i>Current Eye Research</i> , 2011 , 36, 1139-47	2.9	8
21	mRNA Expression Level of Interleukin Genes in the Determining Phases of Behçet's Disease. <i>Annals of Dermatology</i> , 2015 , 27, 291-7	0.4	7
20	Deep sclerectomy with various implants: an experimental and histopathologic study in a rabbit model. <i>Ophthalmologica</i> , 2004 , 218, 264-9	3.7	7
19	Figure-of-eight vertical mattress suture technique for anterior flap suspension to overlying tissues in external dacryocystorhinostomy. <i>American Journal of Ophthalmology</i> , 2007 , 143, 328-333	4.9	6
18	Normative Values of Craniofacial Measurements in Idiopathic Benign Macrocephalic Children. <i>Cleft Palate-Craniofacial Journal</i> , 2001 , 38, 260-263	1.9	6
17	The migration pattern, patient selection with diagnostic methodological flaw and confusing naming dilemma in Behçet disease. <i>European Journal of Echocardiography</i> , 2007 , 8, 167-73; author reply 174		5
16	The efficacy of intravitreal piperacillin/tazobactam in rabbits with experimental Staphylococcus epidermidis endophthalmitis: a comparison with vancomycin. <i>Ophthalmic Research</i> , 2005 , 37, 168-74	2.9	5
15	Excessive maternal caffeine exposure during pregnancy is cataractogenic for neonatal crystalline lenses in rats: a biomicroscopic and histopathologic study. <i>Acta Ophthalmologica</i> , 2004 , 82, 552-6		5
14	Interpupillary index: a new parameter for hypo-hypertelorism. <i>Journal of Cranio-Maxillo-Facial Surgery</i> , 2001 , 29, 191-4	3.6	5
13	Diagnostic dilemma between intestinal Behçet disease and inflammatory bowel disease with pyoderma gangrenosum. <i>World Journal of Gastroenterology</i> , 2006 , 12, 5748-51	5.6	5
12	High dose of caffeine administered to pregnant rats causes histopathological changes in the cornea of newborn pups. <i>Medical Science Monitor</i> , 2003 , 9, BR168-73	3.2	5
11	Shifting crystalline pseudohypopyon secondary to lens absorption with spontaneous openings in the anterior lens capsule. <i>Journal of Cataract and Refractive Surgery</i> , 2006 , 32, 1971-3	2.3	4
10	Tumour necrosis factor alpha, lipid peroxidation and NO* are increased and associated with decreased free-radical scavenging enzymes in patients with Weill-Marchesani syndrome. <i>Mediators of Inflammation</i> , 2004 , 13, 165-70	4.3	4
9	Tc-99m HMPAO brain SPECT in linear nevus sebaceous syndrome. <i>Annals of Nuclear Medicine</i> , 2003 , 17, 703-6	2.5	4
8	Bilateral epibulbar and posterior scleral choristomas in linear nevus sebaceous syndrome. <i>Acta Ophthalmologica</i> , 2003 , 81, 312-4		4

7	A comparison of intravitreal piperacillin/tazobactam with ceftazidime in experimental Pseudomonas aeruginosa endophthalmitis. <i>Experimental Eye Research</i> , 2005 , 80, 361-7	3.7	3
6	Author's response. <i>Survey of Ophthalmology</i> , 2006 , 51, 174-177	6.1	3
5	Decreased central corneal thickness in children with Down syndrome. <i>Journal of Pediatric Ophthalmology and Strabismus</i> , 2002 , 39, 274-7	0.9	3
4	Primary Probing with and without Monoka Silastic Stent Intubation for Epiphora in Older Children and Adults. <i>Current Eye Research</i> , 2020 , 45, 87-90	2.9	2
3	Regarding neutrophil and lymphocyte responses to oral Streptococcus in Adamantiades-Behçet's disease. <i>FEMS Immunology and Medical Microbiology</i> , 2006 , 47, 311-4		1
2	Internuclear ophthalmoplegia as a symptom associated with Henoch-Schoenlein purpura. <i>Neuro-Ophthalmology</i> , 2002 , 26, 193-196	0.9	1
1	Breakage of the Ritleng Tip Within the Interosseous Nasolacrimal Ductus During Monoka Silastic Stent Intubation and its Management. <i>Journal of Maxillofacial and Oral Surgery</i> , 1	0.9	