## Cem Evereklioglu

## List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1384096/cem-evereklioglu-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

1,851
citations

h-index

42
g-index

2,057
ext. papers

2.9
avg, IF

L-index

#	Paper	IF	Citations
60	Primary Probing with and without Monoka Silastic Stent Intubation for Epiphora in Older Children and Adults. <i>Current Eye Research</i> , <b>2020</b> , 45, 87-90	2.9	2
59	mRNA Expression Level of Interleukin Genes in the Determining Phases of Behāts Disease. <i>Annals of Dermatology</i> , <b>2015</b> , 27, 291-7	0.4	7
58	Kite-tailSfascia lata strips technique: frontalis suspension using a non-endoscopic minimally invasive single-thigh incision approach. <i>British Journal of Ophthalmology</i> , <b>2012</b> , 96, 570-5	5.5	9
57	Ocular Behët disease: current therapeutic approaches. <i>Current Opinion in Ophthalmology</i> , <b>2011</b> , 22, 508-16	5.1	28
56	Do non-steroidal anti-inflammatory drugs delay posterior capsule opacification after phacoemulsification in children? A randomized, prospective controlled trial. <i>Current Eye Research</i> , <b>2011</b> , 36, 1139-47	2.9	8
55	Invasive squamous cell carcinoma of the conjunctiva first misdiagnosed as a pterygium: a clinicopathologic case report. <i>Eye and Contact Lens</i> , <b>2008</b> , 34, 188-90	3.2	9
54	The migration pattern, patient selection with diagnostic methodological flaw and confusing naming dilemma in Behat disease. <i>European Journal of Echocardiography</i> , <b>2007</b> , 8, 167-73; author reply 174		5
53	Figure-of-eight vertical mattress suture technique for anterior flap suspension to overlying tissues in external dacryocystorhinostomy. <i>American Journal of Ophthalmology</i> , <b>2007</b> , 143, 328-333	4.9	6
52	Ingestion of IH636 grape seed proanthocyanidin extract to prevent selenite-induced oxidative stress in experimental cataract. <i>Journal of Cataract and Refractive Surgery</i> , <b>2006</b> , 32, 1041-5	2.3	35
51	Shifting crystalline pseudohypopyon secondary to lens absorption with spontaneous openings in the anterior lens capsule. <i>Journal of Cataract and Refractive Surgery</i> , <b>2006</b> , 32, 1971-3	2.3	4
50	Author's response:. Survey of Ophthalmology, 2006, 51, 174-177	6.1	3
49	Regarding neutrophil and lymphocyte responses to oral Streptococcus in Adamantiades-Behllts disease. <i>FEMS Immunology and Medical Microbiology</i> , <b>2006</b> , 47, 311-4		1
48	Clinical features of Behets disease in children. <i>International Journal of Dermatology</i> , <b>2006</b> , 45, 713-6	1.7	55
47	Diagnostic dilemma between intestinal Behät disease and inflammatory bowel disease with pyoderma gangrenosum. <i>World Journal of Gastroenterology</i> , <b>2006</b> , 12, 5748-51	5.6	5
46	The efficacy of intravitreal piperacillin/tazobactam in rabbits with experimental Staphylococcus epidermidis endophthalmitis: a comparison with vancomycin. <i>Ophthalmic Research</i> , <b>2005</b> , 37, 168-74	2.9	5
45	A comparison of intravitreal piperacillin/tazobactam with ceftazidime in experimental Pseudomonas aeruginosa endophthalmitis. <i>Experimental Eye Research</i> , <b>2005</b> , 80, 361-7	3.7	3
44	Current concepts in the etiology and treatment of Behllt disease. <i>Survey of Ophthalmology</i> , <b>2005</b> , 50, 297-350	6.1	295

## (2003-2005)

43	Adenosine deaminase enzyme levels, their relation with disease activity, and the effect of colchicine on adenosine deaminase levels in patients with Behlets disease. <i>Rheumatology International</i> , <b>2005</b> , 25, 452-6	3.6	21	
42	Synovial nitric oxide concentrations are increased and correlated with serum levels in patients with active Behats disease: a pilot study. <i>Clinical Rheumatology</i> , <b>2005</b> , 24, 324-30	3.9	18	
41	Serum levels of soluble P-selectin are increased and associated with disease activity in patients with Behlets syndrome. <i>Mediators of Inflammation</i> , <b>2005</b> , 2005, 237-41	4.3	14	
40	The efficacy of piperacillin/tazobactam in experimental Pseudomonas aeruginosa endophthalmitis: a histopathological and microbiological evaluation. <i>Current Eye Research</i> , <b>2005</b> , 30, 13-9	2.9	12	
39	Deep sclerectomy with various implants: an experimental and histopathologic study in a rabbit model. <i>Ophthalmologica</i> , <b>2004</b> , 218, 264-9	3.7	7	
38	Determination of nontoxic concentrations of piperacillin/tazobactam for intravitreal application. An electroretinographic, histopathologic and morphometric analysis. <i>Ophthalmic Research</i> , <b>2004</b> , 36, 139-44	2.9	14	
37	Tumour necrosis factor alpha, lipid peroxidation and NO* are increased and associated with decreased free-radical scavenging enzymes in patients with Weill-Marchesani syndrome. <i>Mediators of Inflammation</i> , <b>2004</b> , 13, 165-70	4.3	4	
36	Excessive maternal caffeine exposure during pregnancy is cataractogenic for neonatal crystalline lenses in rats: a biomicroscopic and histopathologic study. <i>Acta Ophthalmologica</i> , <b>2004</b> , 82, 552-6		5	
35	Pattern electroretinogram for monitoring the efficacy of intravitreal triamcinolone injection in diabetic macular edema. <i>Documenta Ophthalmologica</i> , <b>2004</b> , 109, 139-45	2.2	14	
34	Pattern VEP is a useful technique in monitoring the effectiveness of occlusion therapy in amblyopic eyes under occlusion therapy. <i>Documenta Ophthalmologica</i> , <b>2004</b> , 109, 223-7	2.2	18	
33	Managing the symptoms of Beh⊞tS disease. Expert Opinion on Pharmacotherapy, <b>2004</b> , 5, 317-28	4	38	
32	Hammer-chisel technique in endoscopic dacryocystorhinostomy. <i>Annals of Otology, Rhinology and Laryngology</i> , <b>2003</b> , 112, 444-9	2.1	15	
31	Adenosine deaminase enzyme activity is increased and negatively correlates with catalase, superoxide dismutase and glutathione peroxidase in patients with Behats disease: original contributions/clinical and laboratory investigations. <i>Mediators of Inflammation</i> , 2003, 12, 107-16	4.3	46	
30	The pathophysiological significance of red blood cell nitric oxide concentrations in inflammatory BehBtS disease. <i>Mediators of Inflammation</i> , <b>2003</b> , 12, 255-6	4.3	14	
29	Nitric oxide and lipid peroxidation are increased and associated with decreased antioxidant enzyme activities in patients with age-related macular degeneration. <i>Documenta Ophthalmologica</i> , <b>2003</b> , 106, 129-36	2.2	71	
28	Tc-99m HMPAO brain SPECT in linear nevus sebaceous syndrome. <i>Annals of Nuclear Medicine</i> , <b>2003</b> , 17, 703-6	2.5	4	
27	Bilateral epibulbar and posterior scleral choristomas in linear nevus sebaceous syndrome. <i>Acta Ophthalmologica</i> , <b>2003</b> , 81, 312-4		4	
26	Effect of gestational nicotine treatment on newborn rat retina: a histopathological and morphometric analysis. <i>Ophthalmic and Physiological Optics</i> , <b>2003</b> , 23, 527-33	4.1	14	

25	Urinary nitric oxide levels are increased and correlated with plasma concentrations in patients with Behats disease: is it a new urinary activity marker?. <i>Nephrology</i> , <b>2003</b> , 8, 231-8	2.2	9
24	Comparison of secondary implantation of flexible open-loop anterior chamber and scleral-fixated posterior chamber intraocular lenses. <i>Journal of Cataract and Refractive Surgery</i> , <b>2003</b> , 29, 301-8	2.3	113
23	High dose of caffeine administered to pregnant rats causes histopathological changes in the cornea of newborn pups. <i>Medical Science Monitor</i> , <b>2003</b> , 9, BR168-73	3.2	5
22	Internuclear ophthalmoplegia as a symptom associated with Henoch-Schoenlein purpura. <i>Neuro-Ophthalmology</i> , <b>2002</b> , 26, 193-196	0.9	1
21	Aqueous humor adrenomedullin levels differ in patients with different types of glaucoma. <i>Japanese Journal of Ophthalmology</i> , <b>2002</b> , 46, 203-8	2.6	10
20	Central corneal thickness is lower in osteogenesis imperfecta and negatively correlates with the presence of blue sclera. <i>Ophthalmic and Physiological Optics</i> , <b>2002</b> , 22, 511-5	4.1	60
19	Urinary adrenomedullin levels are increased and correlated with plasma concentrations in patients with BehBtS syndrome. <i>International Journal of Urology</i> , <b>2002</b> , 9, 296-303	2.3	15
18	Craniofacial Anthropometry in a Turkish Population. Cleft Palate-Craniofacial Journal, 2002, 39, 208-218	1.9	19
17	Serum levels of TNF-alpha, sIL-2R, IL-6, and IL-8 are increased and associated with elevated lipid peroxidation in patients with Behllts disease. <i>Mediators of Inflammation</i> , <b>2002</b> , 11, 87-93	4.3	163
16	Increased nitric oxide production in patients with Behlets disease: is it a new activity marker?. <i>Journal of the American Academy of Dermatology</i> , <b>2002</b> , 46, 50-4	4.5	86
15	Use of caffeic acid phenethyl ester to prevent sodium-selenite-induced cataract in rat eyes. <i>Journal of Cataract and Refractive Surgery</i> , <b>2002</b> , 28, 1457-62	2.3	39
14	Lenticular astigmatism in tilted disc syndrome. <i>Journal of Cataract and Refractive Surgery</i> , <b>2002</b> , 28, 183	62:40	19
13	Decreased Central Corneal Thickness in Children With Down Syndrome. <i>Journal of Pediatric Ophthalmology and Strabismus</i> , <b>2002</b> , 39, 274-277	0.9	18
12	Decreased central corneal thickness in children with Down syndrome. <i>Journal of Pediatric Ophthalmology and Strabismus</i> , <b>2002</b> , 39, 274-7	0.9	3
11	Normative Values of Craniofacial Measurements in Idiopathic Benign Macrocephalic Children. <i>Cleft Palate-Craniofacial Journal</i> , <b>2001</b> , 38, 260-263	1.9	6
10	Defective visual field tests in chronic heavy smokers. <i>Acta Ophthalmologica</i> , <b>2001</b> , 79, 53-6		22
9	Interpupillary index: a new parameter for hypo-hypertelorism. <i>Journal of Cranio-Maxillo-Facial Surgery</i> , <b>2001</b> , 29, 191-4	3.6	5
8	The effect of reading and near-work on the development of myopia in emmetropic boys: a prospective, controlled, three-year follow-up study. <i>Vision Research</i> , <b>2001</b> , 41, 2511-20	2.1	91

## LIST OF PUBLICATIONS

7	Audio-vestibular evaluation in patients with Beh日tS syndrome. <i>Journal of Laryngology and Otology</i> , <b>2001</b> , 115, 704-8	1.8	44
6	Increased plasma adrenomedullin levels in patients with BehletS disease. Dermatology, 2000, 201, 312-5	5 4.4	44
5	Autonomic nervous system functions in obese children. <i>Brain and Development</i> , <b>2000</b> , 22, 151-3	2.2	32
4	Comparative external versus endoscopic dacryocystorhinostomy: results in 115 patients (130 eyes). Otolaryngology - Head and Neck Surgery, <b>2000</b> , 123, 488-91	5.5	154
3	Weill-Marchesani syndrome in three generations. <i>Eye</i> , <b>1999</b> , 13 ( Pt 6), 773-7	4.4	28
2	Caffeic acid phenethyl ester to inhibit posterior capsule opacification in rabbits. <i>Journal of Cataract and Refractive Surgery</i> , <b>1997</b> , 23, 1572-6	2.3	51
1	Breakage of the Ritleng Tip Within the Interosseous Nasolacrimal Ductus During Monoka Silastic Stent Intubation and its Management. <i>Journal of Maxillofacial and Oral Surgery</i> ,1	0.9	