

Han-Bin Oh

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1398314/han-bin-oh-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

94
papers

2,219
citations

21
h-index

45
g-index

97
ext. papers

2,451
ext. citations

4.2
avg, IF

4.58
L-index

#	Paper	IF	Citations
94	Detailed unfolding and folding of gaseous ubiquitin ions characterized by electron capture dissociation. <i>Journal of the American Chemical Society</i> , 2002 , 124, 6407-20	16.4	273
93	Top-down mass spectrometry of a 29-kDa protein for characterization of any posttranslational modification to within one residue. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 1774-9	11.5	232
92	Secondary and tertiary structures of gaseous protein ions characterized by electron capture dissociation mass spectrometry and photofragment spectroscopy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 15863-8	11.5	210
91	Infrared photodissociation spectroscopy of electrosprayed ions in a Fourier transform mass spectrometer. <i>Journal of the American Chemical Society</i> , 2005 , 127, 4076-83	16.4	149
90	Nonergodic and conformational control of the electron capture dissociation of protein cations. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 14011-6	11.5	149
89	Plasma electron capture dissociation for the characterization of large proteins by top down mass spectrometry. <i>Analytical Chemistry</i> , 2003 , 75, 1599-603	7.8	71
88	Top down characterization of secreted proteins from Mycobacterium tuberculosis by electron capture dissociation mass spectrometry. <i>Journal of the American Society for Mass Spectrometry</i> , 2003 , 14, 253-61	3.5	70
87	Optimized Automatic Noise Level Calculations for Broadband FT-ICR Mass Spectra of Petroleum Give More Reliable and Faster Peak Picking Results. <i>Bulletin of the Korean Chemical Society</i> , 2009 , 30, 2665-2668	1.2	57
86	Gas-phase peptide sequencing by TEMPO-mediated radical generation. <i>Analyst, The</i> , 2009 , 134, 1706-12	5	55
85	Proposed nomenclature for peptide ion fragmentation. <i>International Journal of Mass Spectrometry</i> , 2015 , 390, 24-27	1.9	51
84	Hydrogen Atom Loss in Electron-Capture Dissociation: A Fourier Transform-Ion Cyclotron Resonance Study with Single Isotopomeric Ubiquitin Ions. <i>European Journal of Mass Spectrometry</i> , 2002 , 8, 177-180	1.1	48
83	Disulfide bond cleavage in TEMPO-free radical initiated peptide sequencing mass spectrometry. <i>Journal of Mass Spectrometry</i> , 2011 , 46, 830-9	2.2	37
82	Radical-driven peptide backbone dissociation tandem mass spectrometry. <i>Mass Spectrometry Reviews</i> , 2015 , 34, 116-32	11	35
81	Numerous isomers of serine octamer ions characterized by infrared photodissociation spectroscopy. <i>ChemPhysChem</i> , 2009 , 10, 2603-6	3.2	35
80	Observation of pronounced b*,y cleavages in the electron capture dissociation mass spectrometry of polyamidoamine (PAMAM) dendrimer ions with amide functionalities. <i>Journal of the American Society for Mass Spectrometry</i> , 2006 , 17, 536-543	3.5	34
79	Multivariate analysis of electron detachment dissociation and infrared multiphoton dissociation mass spectra of heparan sulfate tetrasaccharides differing only in hexuronic acid stereochemistry. <i>Journal of the American Society for Mass Spectrometry</i> , 2011 , 22, 582-90	3.5	31
78	Dynamics of harpooning studied by transition state spectroscopy. II. Li??FH. <i>Journal of Chemical Physics</i> , 2000 , 113, 9897-9900	3.9	28

77	One-step peptide backbone dissociations in negative-ion free radical initiated peptide sequencing mass spectrometry. <i>Analytical Chemistry</i> , 2013 , 85, 7044-51	7.8	27
76	Ultraviolet photodissociation at 266 nm of phosphorylated peptide cations. <i>Rapid Communications in Mass Spectrometry</i> , 2009 , 23, 3609-20	2.2	25
75	Theoretical study of the ionic hydrogen bond in the isolated proton-bound dimer of cytosine. <i>Chemical Physics Letters</i> , 2006 , 432, 269-274	2.5	25
74	Photoelectron spectroscopy of pyridine cluster anions, (Py) _n ⁻ (n=4-13). <i>Journal of Chemical Physics</i> , 1999 , 111, 4041-4050	3.9	24
73	Base-pair interactions in the gas-phase proton-bonded complexes of C+G and C+GC. <i>Journal of Chemical Physics</i> , 2007 , 127, 245102	3.9	21
72	Characterization of permethylated β-cyclodextrin-peptide noncovalently bound complexes using electron capture dissociation mass spectrometry (ECD MS). <i>International Journal of Mass Spectrometry</i> , 2009 , 279, 47-52	1.9	20
71	Reliable screening and confirmation of 156 multi-class illegal adulterants in dietary supplements based on extracted common ion chromatograms by ultra-high-performance liquid chromatography-quadrupole/time of flight-mass spectrometry. <i>Journal of Chromatography A</i> , 2017 , 1491, 43-56	4.5	18
70	Electron capture dissociation mass spectrometry of peptide cations containing a lysine homologue: a mobile proton model for explaining the observation of b-type product ions. <i>Rapid Communications in Mass Spectrometry</i> , 2006 , 20, 3167-75	2.2	18
69	Photodissociation of TEMPO-modified peptides: new approaches to radical-directed dissociation of biomolecules. <i>Physical Chemistry Chemical Physics</i> , 2014 , 16, 4871-9	3.6	17
68	Dynamic modeling of lactic acid fermentation metabolism with <i>Lactococcus lactis</i> . <i>Journal of Microbiology and Biotechnology</i> , 2011 , 21, 162-9	3.3	16
67	Chiral differentiation of d- and l-alanine by permethylated β-cyclodextrin: IRMPD spectroscopy and DFT methods. <i>Physical Chemistry Chemical Physics</i> , 2017 , 19, 14729-14737	3.6	15
66	Tautomerism and isomerism of guanine-cytosine DNA base pair: Ab initio and density functional theory approaches. <i>Computational and Theoretical Chemistry</i> , 2005 , 730, 241-249		15
65	Quantitative analysis of polyhexamethylene guanidine (PHMG) oligomers via matrix-assisted laser desorption/ionization time-of-flight mass spectrometry with an ionic-liquid matrix. <i>Rapid Communications in Mass Spectrometry</i> , 2015 , 29, 213-9	2.2	14
64	Infrared multiple photon dissociation spectroscopy and density functional theory (DFT) studies of protonated permethylated β-cyclodextrin-water non-covalent complexes. <i>Physical Chemistry Chemical Physics</i> , 2014 , 16, 8376-83	3.6	14
63	Determination of the Intracellular Concentrations of Metabolites in <i>Escherichia coli</i> Collected during the Exponential and Stationary Growth Phases using Liquid Chromatography-Mass Spectrometry. <i>Bulletin of the Korean Chemical Society</i> , 2011 , 32, 524-530	1.2	14
62	An anomalous dissociation of protonated cluster ions of DNA guanine-cytosine base-pair. <i>Mass Spectrometry Letters</i> , 2011 , 2, 73-75		14
61	Biofilm development of <i>Bacillus siamensis</i> ATKU1 on pristine short chain low-density polyethylene: A case study on microbe-microplastics interaction. <i>Journal of Hazardous Materials</i> , 2021 , 409, 124516	12.8	14
60	Chiral differentiation of d- and l-isoleucine using permethylated β-cyclodextrin: infrared multiple photon dissociation spectroscopy, ion-mobility mass spectrometry, and DFT calculations. <i>Physical Chemistry Chemical Physics</i> , 2018 , 20, 30428-30436	3.6	14

59	Application of paper EWOD (electrowetting-on-dielectrics) chip: Protein tryptic digestion and its detection using MALDI-TOF mass spectrometry. <i>Biochip Journal</i> , 2017 , 11, 146-152	4	13
58	Synergistic effects of urban tributary mixing on dissolved organic matter biodegradation in an impounded river system. <i>Science of the Total Environment</i> , 2019 , 676, 105-119	10.2	12
57	Structural and functional characterization of an Isd-type haem-degradation enzyme from <i>Listeria monocytogenes</i> . <i>Acta Crystallographica Section D: Biological Crystallography</i> , 2014 , 70, 615-26		12
56	Dynamics of harpooning studied by transition state spectroscopy. Part III. Li...FCH3. <i>Faraday Discussions</i> , 2001 , 191-207; discussion 233-45	3.6	12
55	MALDI-TOF Analysis of Polyhexamethylene Guanidine (PHMG) Oligomers Used as a Commercial Antibacterial Humidifier Disinfectant. <i>Bulletin of the Korean Chemical Society</i> , 2013 , 34, 1708-1714	1.2	12
54	A Liquid Chromatography-Quadrupole-Time of Flight Mass Spectrometry (LC-Q-TOF MS) Study for Analyzing 35 Corticosteroid Compounds: Elucidation of MS/MS Fragmentation Pathways. <i>Bulletin of the Korean Chemical Society</i> , 2016 , 37, 1029-1038	1.2	11
53	LC-MS/MS Software for Screening Unknown Erectile Dysfunction Drugs and Analogues: Artificial Neural Network Classification, Peak-Count Scoring, Simple Similarity Search, and Hybrid Similarity Search Algorithms. <i>Analytical Chemistry</i> , 2019 , 91, 9119-9128	7.8	10
52	Host-guest chemistry in the gas phase: complex formation with 18-crown-6 enhances helicity of alanine-based peptides. <i>Journal of Physical Chemistry A</i> , 2011 , 115, 14215-20	2.8	10
51	Evaluation of the internal temperatures of an 8.6 kDa protein cation exposed to a hot dispenser cathode employed in electron capture dissociation mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2006 , 20, 1918-24	2.2	10
50	Charge-Directed Peptide Backbone Dissociations of o-TEMPO-Bz-C(O)-Peptides. <i>Mass Spectrometry Letters</i> , 2013 , 4, 71-74		10
49	Properties of Polyhexamethylene Guanidine (PHMG) Associated with Fatal Lung Injury in Korea. <i>Molecules</i> , 2020 , 25,	4.8	10
48	Noncovalent Complexes of Cyclodextrin with Small Organic Molecules: Applications and Insights into Host-Guest Interactions in the Gas Phase and Condensed Phase. <i>Molecules</i> , 2020 , 25,	4.8	10
47	Rapid screening of sulfonamides in dietary supplements based on extracted common ion chromatogram and neutral loss scan by LC-Q/TOF-mass spectrometry. <i>Journal of Food and Drug Analysis</i> , 2019 , 27, 164-174	7	10
46	Ionization of Gas-Phase Polycyclic Aromatic Hydrocarbons in Electrospray Ionization Coupled with Gas Chromatography. <i>Analytical Chemistry</i> , 2018 , 90, 4203-4211	7.8	9
45	MALDI in-source decay mass spectrometry of polyamidoamine dendrimers. <i>Journal of the American Society for Mass Spectrometry</i> , 2012 , 23, 1821-5	3.5	9
44	Online Simultaneous Hydrogen/Deuterium Exchange of Multitarget Gas-Phase Molecules by Electrospray Ionization Mass Spectrometry Coupled with Gas Chromatography. <i>Analytical Chemistry</i> , 2017 , 89, 12284-12292	7.8	8
43	Design and synthesis of new mass tags for matrix-free laser desorption ionization mass spectrometry (LDI-MS) based on 6,11-dihydrothiochromeno[4,3-b]indole. <i>Tetrahedron</i> , 2016 , 72, 5612-5619	2.4	8
42	Reliable screening and classification of phosphodiesterase type 5 inhibitors in dietary supplements using gas chromatography / mass spectrometry combined with specific common ions. <i>Journal of Chromatography A</i> , 2020 , 1623, 461210	4.5	7

41	Multiresidue analysis of 85 persistent organic pollutants in small human serum samples by modified QuEChERS preparation with different ionization sources in mass spectrometry. <i>Journal of Chromatography A</i> , 2020 , 1623, 461170	4.5	7
40	TEMPO-Assisted Free Radical-Initiated Peptide Sequencing Mass Spectrometry (FRIPS MS) in Q-TOF and Orbitrap Mass Spectrometers: Single-Step Peptide Backbone Dissociations in Positive Ion Mode. <i>Journal of the American Society for Mass Spectrometry</i> , 2017 , 28, 154-163	3.5	7
39	Bromine isotopic signature facilitates de novo sequencing of peptides in free-radical-initiated peptide sequencing (FRIPS) mass spectrometry. <i>Journal of Mass Spectrometry</i> , 2015 , 50, 378-87	2.2	7
38	Zinc-finger motif noncovalent interactions with double-stranded DNA characterized by negative-ion electrospray ionization mass spectrometry. <i>Analyst, The</i> , 2011 , 136, 3739-46	5	7
37	Synergetic and antagonistic role of natural antioxidant in the autoxidation of soybean oil. <i>Journal of Industrial and Engineering Chemistry</i> , 2011 , 17, 537-542	6.3	7
36	Structural and biochemical study of <i>Bacillus subtilis</i> HmoB in complex with heme. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 446, 286-91	3.4	6
35	Sample preparation of chemical warfare agent simulants on a digital microfluidic (DMF) device using magnetic bead-based solid-phase extraction. <i>Microfluidics and Nanofluidics</i> , 2017 , 21, 1	2.8	6
34	Infrared Multiple Photon Depletion of the Gas-phase Proton-bound Cytosine Dimer. <i>Chemistry Letters</i> , 2015 , 44, 1756-1758	1.7	6
33	Guanidination of lysine residue improves the sensitivity and facilitates the interpretation of free radical initiated peptide sequencing (FRIPS) mass spectrometry results. <i>International Journal of Mass Spectrometry</i> , 2015 , 390, 110-117	1.9	6
32	Investigation of the MALDI process used to characterize self-assembled monolayers of alkanethiolates on gold. <i>Langmuir</i> , 2009 , 25, 3692-7	4	6
31	Increased incorporation of gaseous CO into succinate by <i>Escherichia coli</i> overexpressing carbonic anhydrase and phosphoenolpyruvate carboxylase genes. <i>Journal of Biotechnology</i> , 2017 , 241, 101-107	3.7	5
30	Characteristics of Exposure to Chloromethylisothiazolinone (CMIT) and Methylisothiazolinone (MIT) among Humidifier Disinfectant-Associated Lung Injury (HDLI) Patients in South Korea. <i>Molecules</i> , 2020 , 25,	4.8	5
29	Quantitative MALDI-TOF mass spectrometric analysis of biocidal polyhexamethylene guanidine (PHMG) oligomers in consumer products. <i>International Journal of Mass Spectrometry</i> , 2019 , 435, 298-304 ^{1.9}	1.9	5
28	Untargeted Metabolomics and Steroid Signatures in Urine of Male Pattern Baldness Patients after Finasteride Treatment for a Year. <i>Metabolites</i> , 2020 , 10,	5.6	5
27	Enhanced Incorporation of Gaseous CO ₂ to Succinate by a Recombinant <i>Escherichia coli</i> W3110. <i>Biotechnology and Bioprocess Engineering</i> , 2019 , 24, 103-108	3.1	4
26	Fragmentation Pathways of Tadalafil and Its Analogues in Electrospray Ionization Tandem Mass Spectrometry. <i>Bulletin of the Korean Chemical Society</i> , 2018 , 39, 190-196	1.2	4
25	Tandem mass spectrometric analysis of isosorbide-1,4-cyclohexane-dicarboxylic acid polyester oligomer cations using ion-trap mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2013 , 27, 1913-8	2.2	4
24	Prediction of liquid chromatography retention times of erectile dysfunction drugs and analogues using chemometric approaches. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2017 , 40, 790-797	1.3	4

23	MALDI-TOF Mass Spectrometric Analysis of Chemical Warfare Nerve Agent Simulants. <i>Bulletin of the Korean Chemical Society</i> , 2016 , 37, 316-320	1.2	4
22	Inactivation of the [(1, 2)-xylosyltransferase and the [(1, 3)-fucosyltransferase gene in rice (<i>Oryza sativa</i>) by multiplex CRISPR/Cas9 strategy. <i>Plant Cell Reports</i> , 2021 , 40, 1025-1035	5.1	4
21	Protein Analysis Using a Combination of an Online Monolithic Trypsin Immobilized Enzyme Reactor and Collisionally-Activated Dissociation/Electron Transfer Dissociation Dual Tandem Mass Spectrometry. <i>Bulletin of the Korean Chemical Society</i> , 2012 , 33, 3233-3240	1.2	3
20	A Multidimensional System for Phosphopeptide Analysis Using TiO ₂ Enrichment and Ion-exchange Chromatography with Mass Spectrometry. <i>Bulletin of the Korean Chemical Society</i> , 2012 , 33, 3298-3302	1.2	3
19	Collisionally-Activated Dissociation of Peptides with a Disulfide Bond: Confirmation of the Mobile-Proton Model Based Explanation 2010 , 1, 5-8		3
18	Photo-Oxidative Protection of Chlorophyll in C-Phycocyanin Aqueous Medium. <i>Antioxidants</i> , 2020 , 9,	7.1	3
17	Comprehensive screening of multiclass illegal adulterants in herbal supplements and Spice-type drugs using specific MS/MS fragmentations by UHPLC-Q/TOF-MS. <i>Analytical Methods</i> , 2019 , 11, 5260-5273	3.2	2
16	Graph theory-based reaction pathway searches and DFT calculations for the mechanism studies of free radical-initiated peptide sequencing mass spectrometry (FRIPS MS): a model gas-phase reaction of GGR tri-peptide. <i>Physical Chemistry Chemical Physics</i> , 2020 , 22, 5057-5069	3.6	2
15	Collisional Activation Dissociation Mass Spectrometry Studies of Oligosaccharides Conjugated with Na ⁺ -Encapsulated Dibenzo-18-Crown-6 Ether. <i>Mass Spectrometry Letters</i> , 2016 , 7, 96-101		2
14	Examination of Various Metal Ion Sources for Reducing Nonspecific Zinc finger-Zn ²⁺ Complex Formation in ESI Mass Spectrometry. <i>Mass Spectrometry Letters</i> , 2012 , 3, 82-85		2
13	Characteristics of the Molecular Weight of Polyhexamethylene Guanidine (PHMG) Used as a Household Humidifier Disinfectant. <i>Molecules</i> , 2021 , 26,	4.8	2
12	Production of uracil from methane by a newly isolated <i>Methylomonas</i> sp. SW1. <i>Journal of Biotechnology</i> , 2016 , 240, 43-47	3.7	2
11	Free Radical-Initiated Peptide Sequencing Mass Spectrometry for Phosphopeptide Post-translational Modification Analysis. <i>Journal of the American Society for Mass Spectrometry</i> , 2019 , 30, 538-547	3.5	2
10	Unveiling host-guest-solvent interactions in solution by identifying highly unstable host-guest configurations in thermal non-equilibrium gas phase.. <i>Scientific Reports</i> , 2022 , 12, 8169	4.9	2
9	Sex-related differences in urinary immune-related metabolic profiling of alopecia areata patients. <i>Metabolomics</i> , 2020 , 16, 15	4.7	1
8	Microbial production of uracil by an isolated <i>Methylobacterium</i> sp. WJ4 using methanol. <i>Enzyme and Microbial Technology</i> , 2018 , 111, 63-66	3.8	1
7	High Accuracy Mass Measurement Approach in the Identification of Phospholipids in Lipid Extracts: 7 T Fourier-transform Mass Spectrometry and MS/MS Validation. <i>Bulletin of the Korean Chemical Society</i> , 2011 , 32, 1170-1178	1.2	1
6	Density Functional Theory (DFT) Study of Gas-phase O.C Bond Dissociation Energy of Models for o-TEMPO-Bz-C(O)-Peptide: A Model Study for Free Radical Initiated Peptide Sequencing. <i>Bulletin of the Korean Chemical Society</i> , 2014 , 35, 770-774	1.2	1

5	Liposome leakage and increased cellular permeability induced by guanidine-based oligomers: effects of liposome composition on liposome leakage and human lung epithelial barrier permeability.. <i>RSC Advances</i> , 2021 , 11, 32000-32011	3.7	1
4	GCMS Method for the Quantitative Analysis of Limonene from Genetically Engineered <i>Saccharomyces cerevisiae</i> . <i>Bulletin of the Korean Chemical Society</i> , 2018 , 39, 1368-1372	1.2	0
3	A solvent-free headspace GC/MS method for sensitive screening of N-nitrosodimethylamine in drug products. <i>Analytical Methods</i> , 2021 , 13, 3402-3409	3.2	0
2	Structural and functional characterization of an Isd-type haem-degradation enzyme from <i>Listeria monocytogenes</i> . Corrigendum. <i>Acta Crystallographica Section D: Biological Crystallography</i> , 2014 , 70, 1498-1498		
1	Abnormal behaviors in the calibration curves of liquid chromatography-tandem mass spectrometry occurring in the quantitative analysis of surfactants near critical micelle concentrations. <i>Journal of Mass Spectrometry</i> , 2020 , 56, e4599	2.2	