List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/140058/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. New England Journal of Medicine, 2017, 377, 2531-2544.	13.9	3,865
2	Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1–2 trial. Lancet Oncology, The, 2019, 20, 31-42.	5.1	1,467
3	Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2015, 373, 2425-2437.	13.9	1,261
4	Lisocabtagene maraleucel for patients with relapsed or refractory large B-cell lymphomas (TRANSCEND NHL 001): a multicentre seamless design study. Lancet, The, 2020, 396, 839-852.	6.3	1,224
5	Phase 1 Results of ZUMA-1: A Multicenter Study of KTE-C19 Anti-CD19 CAR T Cell Therapy in Refractory Aggressive Lymphoma. Molecular Therapy, 2017, 25, 285-295.	3.7	498
6	lbrutinib for patients with relapsed or refractory chronic lymphocytic leukaemia with 17p deletion (RESONATE-17): a phase 2, open-label, multicentre study. Lancet Oncology, The, 2016, 17, 1409-1418.	5.1	290
7	A randomized phase 3 trial of zanubrutinib vs ibrutinib in symptomatic Waldenström macroglobulinemia: the ASPEN study. Blood, 2020, 136, 2038-2050.	0.6	281
8	Objective responses in relapsed T-cell lymphomas with single-agent brentuximab vedotin. Blood, 2014, 123, 3095-3100.	0.6	280
9	Phase 1 studies of central memory–derived CD19 CAR T–cell therapy following autologous HSCT in patients with B-cell NHL. Blood, 2016, 127, 2980-2990.	0.6	264
10	Moxetumomab pasudotox in relapsed/refractory hairy cell leukemia. Leukemia, 2018, 32, 1768-1777.	3.3	184
11	NCCN Guidelines Insights: Cancer-Associated Venous Thromboembolic Disease, Version 2.2018. Journal of the National Comprehensive Cancer Network: JNCCN, 2018, 16, 1289-1303.	2.3	168
12	Relapsed or Refractory Double-Expressor and Double-Hit Lymphomas Have Inferior Progression-Free Survival After Autologous Stem-Cell Transplantation. Journal of Clinical Oncology, 2017, 35, 24-31.	0.8	152
13	Results of a Multicenter Phase II Trial of Brentuximab Vedotin as Second-Line Therapy before Autologous Transplantation in Relapsed/Refractory Hodgkin Lymphoma. Biology of Blood and Marrow Transplantation, 2015, 21, 2136-2140.	2.0	131
14	Ibrutinib Plus Venetoclax for First-Line Treatment of Chronic Lymphocytic Leukemia: Primary Analysis Results From the Minimal Residual Disease Cohort of the Randomized Phase II CAPTIVATE Study. Journal of Clinical Oncology, 2021, 39, 3853-3865.	0.8	115
15	Cancer-Associated Venous Thromboembolic Disease, Version 1.2015. Journal of the National Comprehensive Cancer Network: JNCCN, 2015, 13, 1079-1095.	2.3	100
16	Pivotal Safety and Efficacy Results from Transcend NHL 001, a Multicenter Phase 1 Study of Lisocabtagene Maraleucel (liso-cel) in Relapsed/Refractory (R/R) Large B Cell Lymphomas. Blood, 2019, 134, 241-241.	0.6	89
17	Fixed-duration ibrutinib plus venetoclax for first-line treatment of CLL: primary analysis of the CAPTIVATE FD cohort. Blood, 2022, 139, 3278-3289.	0.6	83
18	Venous Thromboembolic Disease. Journal of the National Comprehensive Cancer Network: JNCCN, 2013, 11, 1402-1429	2.3	80

#	Article	IF	CITATIONS
19	Post-Marketing Use Outcomes of an Anti-CD19 Chimeric Antigen Receptor (CAR) T Cell Therapy, Axicabtagene Ciloleucel (Axi-Cel), for the Treatment of Large B Cell Lymphoma (LBCL) in the United States (US). Blood, 2019, 134, 764-764.	0.6	77
20	Ublituximab and umbralisib in relapsed/refractory B-cell non-Hodgkin lymphoma and chronic lymphocytic leukemia. Blood, 2019, 134, 1811-1820.	0.6	75
21	A phase II study of vorinostat and rituximab for treatment of newly diagnosed and relapsed/refractory indolent non-Hodgkin lymphoma. Haematologica, 2015, 100, 357-362.	1.7	66
22	Phase 1 TRANSCEND CLL 004 study of lisocabtagene maraleucel in patients with relapsed/refractory CLL or SLL. Blood, 2022, 139, 1794-1806.	0.6	66
23	Tolerability and activity of ublituximab, umbralisib, and ibrutinib in patients with chronic lymphocytic leukaemia and non-Hodgkin lymphoma: a phase 1 dose escalation and expansion trial. Lancet Haematology,the, 2019, 6, e100-e109.	2.2	65
24	CD19-directed CAR T-cell therapy for treatment of primary CNS lymphoma. Blood Advances, 2021, 5, 4059-4063.	2.5	62
25	Real-World Evidence of Axicabtagene Ciloleucel for the Treatment of Large B Cell Lymphoma in the United States. Transplantation and Cellular Therapy, 2022, 28, 581.e1-581.e8.	0.6	61
26	Zanubrutinib for the treatment of MYD88 wild-type Waldenström macroglobulinemia: a substudy of the phase 3 ASPEN trial. Blood Advances, 2020, 4, 6009-6018.	2.5	57
27	Brentuximab Vedotin Is Associated with Improved Progression-Free Survival after Allogeneic Transplantation for Hodgkin Lymphoma. Biology of Blood and Marrow Transplantation, 2014, 20, 1864-1868.	2.0	56
28	Venetoclax in Previously Treated Waldenström Macroglobulinemia. Journal of Clinical Oncology, 2022, 40, 63-71.	0.8	53
29	NCCN Guidelines Insights: Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma, Version 2.2019. Journal of the National Comprehensive Cancer Network: JNCCN, 2019, 17, 12-20.	2.3	52
30	Safety and Preliminary Efficacy in Patients with Relapsed/Refractory Mantle Cell Lymphoma Receiving Lisocabtagene Maraleucel in Transcend NHL 001. Blood, 2020, 136, 10-11.	0.6	52
31	Long-Term (≥4 Year and ≥5 Year) Overall Survival (OS) By 12- and 24-Month Event-Free Survival (EFS): An Updated Analysis of ZUMA-1, the Pivotal Study of Axicabtagene Ciloleucel (Axi-Cel) in Patients (Pts) with Refractory Large B-Cell Lymphoma (LBCL). Blood, 2021, 138, 1764-1764.	0.6	48
32	Apoptosis induced by JAK2 inhibition is mediated by Bim and enhanced by the BH3 mimetic ABT-737 in JAK2 mutant human erythroid cells. Blood, 2010, 115, 2901-2909.	0.6	46
33	Brentuximab vedotin activity in diffuse large B-cell lymphoma with CD30 undetectable by visual assessment of conventional immunohistochemistry. Leukemia and Lymphoma, 2017, 58, 1607-1616.	0.6	46
34	Cancer-Associated Venous Thromboembolic Disease, Version 2.2021, NCCN Clinical Practice Guidelines in Oncology. Journal of the National Comprehensive Cancer Network: JNCCN, 2021, 19, 1181-1201.	2.3	43
35	Comparison of 2-year outcomes with CAR T cells (ZUMA-1) vs salvage chemotherapy in refractory large B-cell lymphoma. Blood Advances, 2021, 5, 4149-4155.	2.5	42
36	A Case of HIV-Negative Primary Effusion Lymphoma Treated with Bortezomib, Pegylated Liposomal Doxorubicin, and Rituximab. Clinical Lymphoma and Myeloma, 2008, 8, 300-304.	1.4	41

#	Article	IF	CITATIONS
37	lbrutinib (lbr) Plus Venetoclax (Ven) for First-Line Treatment of Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Results from the MRD Cohort of the Phase 2 CAPTIVATE Study. Blood, 2019, 134, 35-35.	0.6	40
38	Transcend CLL 004: Phase 1 Cohort of Lisocabtagene Maraleucel (liso-cel) in Combination with Ibrutinib for Patients with Relapsed/Refractory (R/R) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL). Blood, 2020, 136, 39-40.	0.6	40
39	Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma, Version 4.2020, NCCN Clinical Practice Guidelines in Oncology. Journal of the National Comprehensive Cancer Network: JNCCN, 2020, 18, 185-217.	2.3	40
40	Impact of Pretransplantation 18F-fluorodeoxy Glucose–Positron Emission Tomography Status on Outcomes after Allogeneic Hematopoietic Cell Transplantation for Non-Hodgkin Lymphoma. Biology of Blood and Marrow Transplantation, 2015, 21, 1605-1611.	2.0	39
41	Umbralisib Plus Ublituximab (U2) Is Superior to Obinutuzumab Plus Chlorambucil (O+Chl) in Patients with Treatment Na÷ve (TN) and Relapsed/Refractory (R/R) Chronic Lymphocytic Leukemia (CLL): Results from the Phase 3 Unity-CLL Study. Blood, 2020, 136, 37-39.	0.6	37
42	Understanding Caregiver Quality of Life in Caregivers of Hospitalized Older Adults With Cancer. Journal of the American Geriatrics Society, 2019, 67, 978-986.	1.3	36
43	Two-Year Follow-up of Transcend NHL 001, a Multicenter Phase 1 Study of Lisocabtagene Maraleucel (liso-cel) in Relapsed or Refractory (R/R) Large B-Cell Lymphomas (LBCL). Blood, 2021, 138, 2840-2840.	0.6	34
44	Efficacy and Safety of Ibrutinib in Patients with Relapsed or Refractory Chronic Lymphocytic Leukemia or Small Lymphocytic Leukemia with 17p Deletion: Results from the Phase II RESONATEâ,,¢-17 Trial. Blood, 2014, 124, 327-327.	0.6	33
45	NCCN Guidelines® Insights: Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma, Version 3.2022. Journal of the National Comprehensive Cancer Network: JNCCN, 2022, 20, 622-634.	2.3	33
46	Ibrutinib (Ibr) Plus Venetoclax (Ven) for First-Line Treatment of Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): 1-Year Disease-Free Survival (DFS) Results From the MRD Cohort of the Phase 2 CAPTIVATE Study. Blood, 2020, 136, 16-17.	0.6	32
47	Outcomes after Allogeneic Stem Cell Transplantation in Patients with Double-Hit and Double-Expressor Lymphoma. Biology of Blood and Marrow Transplantation, 2018, 24, 514-520.	2.0	31
48	Kte-C19 (anti-CD19 CAR T Cells) Induces Complete Remissions in Patients with Refractory Diffuse Large B-Cell Lymphoma (DLBCL): Results from the Pivotal Phase 2 Zuma-1. Blood, 2016, 128, LBA-6-LBA-6.	0.6	30
49	Updated Follow-up of Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Treated with Lisocabtagene Maraleucel in the Phase 1 Monotherapy Cohort of Transcend CLL 004, Including High-Risk and Ibrutinib-Treated Patients. Blood, 2020, 136, 40-41.	0.6	26
50	A Phase 2 Multicenter Trial of KTE-C19 (anti-CD19 CAR T Cells) in Patients With Chemorefractory Primary Mediastinal B-Cell Lymphoma (PMBCL) and Transformed Follicular Lymphoma (TFL): Interim Results From ZUMA-1. Blood, 2016, 128, 998-998.	0.6	26
51	Transcend NHL 001: Immunotherapy with the CD19-Directed CAR T-Cell Product JCAR017 Results in High Complete Response Rates in Relapsed or Refractory B-Cell Non-Hodgkin Lymphoma. Blood, 2016, 128, 4192-4192.	0.6	25
52	Hairy Cell Leukemia, Version 2.2018, NCCN Clinical Practice Guidelines in Oncology. Journal of the National Comprehensive Cancer Network: JNCCN, 2017, 15, 1414-1427.	2.3	24
53	The Cerebroventricular Environment Modifies CAR T Cells for Potent Activity against Both Central Nervous System and Systemic Lymphoma. Cancer Immunology Research, 2021, 9, 75-88.	1.6	24
54	Rapid Undetectable MRD (uMRD) Responses in Patients with Relapsed/Refractory (R/R) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) Treated with Lisocabtagene Maraleucel (liso-cel), a CD19-Directed CAR T Cell Product: Updated Results from Transcend CLL 004, a Phase 1/2 Study Including Patients with High-Risk Disease Previously Treated with Ibrutinib. Blood, 2019, 134, 503-503.	0.6	24

#	Article	IF	CITATIONS
55	The Chronic Lymphocytic Leukemia Comorbidity Index (CLL-CI): A Three-Factor Comorbidity Model. Clinical Cancer Research, 2021, 27, 4814-4824.	3.2	23
56	Phase 1 study of the Aurora kinase A inhibitor alisertib (MLN8237) combined with the histone deacetylase inhibitor vorinostat in lymphoid malignancies. Leukemia and Lymphoma, 2020, 61, 309-317.	0.6	22
57	Multicenter Prospective Phase II Study of Venetoclax in Patients with Previously Treated Waldenstrom Macroglobulinemia. Blood, 2018, 132, 2888-2888.	0.6	22
58	Long-Term Results of High-Dose Therapy and Autologous Stem Cell Transplantation for Mantle Cell Lymphoma: Effectiveness of Maintenance Rituximab. Biology of Blood and Marrow Transplantation, 2017, 23, 1861-1869.	2.0	19
59	Cytokine release syndrome and neurological event costs in lisocabtagene maraleucel–treated patients in the TRANSCEND NHL 001 trial. Blood Advances, 2021, 5, 1695-1705.	2.5	18
60	Developmental partitioning of SYK and ZAP70 prevents autoimmunity and cancer. Molecular Cell, 2021, 81, 2094-2111.e9.	4.5	17
61	Rapid MRD-Negative Responses in Patients with Relapsed/Refractory CLL Treated with Liso-Cel, a CD19-Directed CAR T-Cell Product: Preliminary Results from Transcend CLL 004, a Phase 1/2 Study Including Patients with High-Risk Disease Previously Treated with Ibrutinib. Blood, 2018, 132, 300-300.	0.6	17
62	Fixed-duration (FD) first-line treatment (tx) with ibrutinib (I) plus venetoclax (V) for chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL): Primary analysis of the FD cohort of the phase 2 captivate study Journal of Clinical Oncology, 2021, 39, 7501-7501.	0.8	16
63	Brentuximab Vedotin As First Line Salvage Therapy in Relapsed/Refractory HL. Blood, 2012, 120, 3699-3699.	0.6	16
64	Cost-effectiveness of polatuzumab vedotin combined with chemoimmunotherapy in untreated diffuse large B-cell lymphoma. Blood, 2022, 140, 2697-2708.	0.6	15
65	ASPEN: Long-term follow-up results of a phase 3 randomized trial of zanubrutinib (ZANU) versus ibrutinib (IBR) in patients with Waldenstr¶m macroglobulinemia (WM) Journal of Clinical Oncology, 2022, 40, 7521-7521.	0.8	14
66	2-Year Follow-up and High-Risk Subset Analysis of Zuma-1, the Pivotal Study of Axicabtagene Ciloleucel (Axi-Cel) in Patients with Refractory Large B Cell Lymphoma. Blood, 2018, 132, 2967-2967.	0.6	13
67	TRANSCEND CLL 004: Minimal residual disease (MRD) negative responses after lisocabtagene maraleucel (Liso-Cel; JCAR017), a CD19-directed CAR T cell product, in patients (pts) with relapsed/refractory chronic lymphocytic leukemia or small lymphocytic lymphoma (CLL/SLL) Journal of Clinical Oncology, 2019, 37, 7501-7501.	0.8	13
68	ASPEN: Results of a phase III randomized trial of zanubrutinib versus ibrutinib for patients with Waldenström macroglobulinemia (WM) Journal of Clinical Oncology, 2020, 38, 8007-8007.	0.8	13
69	Role of brentuximab vedotin in the treatment of relapsed or refractory Hodgkin lymphoma. Pharmacogenomics and Personalized Medicine, 2014, 7, 79.	0.4	12
70	Multi-center phase II trial of bortezomib and rituximab maintenance combination therapy in patients with mantle cell lymphoma after consolidative autologous stem cell transplantation. Journal of Hematology and Oncology, 2018, 11, 87.	6.9	12
71	A phase 2, open-label study of brentuximab vedotin in patients with CD30-expressing solid tumors. Investigational New Drugs, 2019, 37, 738-747.	1.2	12
72	Long-term follow-up of the PI3Kδ inhibitor TGR-1202 to demonstrate a differentiated safety profile and high response rates in CLL and NHL: Integrated-analysis of TGR-1202 monotherapy and combined with ublituximab Journal of Clinical Oncology, 2016, 34, 7512-7512.	0.8	12

#	Article	IF	CITATIONS
73	Outcomes by prior lines of therapy (LoT) in ZUMA-1, the pivotal phase 2 study of axicabtagene ciloleucel (Axi-Cel) in patients (Pts) with refractory large B cell lymphoma Journal of Clinical Oncology, 2018, 36, 3039-3039.	0.8	12
74	Phase 1/2 study of cirmtuzumab and ibrutinib in mantle cell lymphoma (MCL) or chronic lymphocytic leukemia (CLL) Journal of Clinical Oncology, 2021, 39, 7556-7556.	0.8	11
75	Lisocabtagene maraleucel (liso-cel) treatment of patients (pts) with relapsed/refractory (R/R) B-cell non-Hodgkin lymphoma (NHL) and secondary CNS lymphoma: Initial results from TRANSCEND NHL 001 Journal of Clinical Oncology, 2019, 37, 7515-7515.	0.8	11
76	CD19-Targeting CAR-T Cell Therapy in CNS Lymphoma. Blood, 2019, 134, 4075-4075.	0.6	10
77	Targeted Therapies in CLL: Monotherapy Versus Combination Approaches. Current Hematologic Malignancy Reports, 2018, 13, 525-533.	1.2	9
78	A Phase 2 Study of Brentuximab Vedotin in Patients with Relapsed or Refractory CD30-Positive Non-Hodgkin Lymphomas: Interim Results Blood, 2012, 120, 2746-2746.	0.6	9
79	Results of a Phase II Trial of Brentuximab Vedotin As First Line Salvage Therapy in Relapsed/Refractory HL Prior to AHCT. Blood, 2014, 124, 501-501.	0.6	9
80	Phase 1 Clinical Results of the ZUMA-1 (KTE-C19-101) Study: A Phase 1-2 Multi-Center Study Evaluating the Safety and Efficacy of Anti-CD19 CAR T Cells (KTE-C19) in Subjects with Refractory Aggressive Non-Hodgkin Lymphoma (NHL). Blood, 2015, 126, 3991-3991.	0.6	9
81	Post Transplant Outcome of a Multicenter Phase II Study of Brentuximab Vedotin As First Line Salvage Therapy in Relapsed/Refractory HL Prior to AHCT. Blood, 2015, 126, 519-519.	0.6	9
82	Real-World Outcomes of Axicabtagene Ciloleucel (Axi-cel) for the Treatment of Large B-Cell Lymphoma (LBCL): Impact of Age and Specific Organ Dysfunction. Blood, 2021, 138, 530-530.	0.6	9
83	Total marrow and lymphoid irradiation as conditioning in haploidentical transplant with posttransplant cyclophosphamide. Blood Advances, 2022, 6, 4098-4106.	2.5	9
84	Model of a Queuing Approach for Patient Accrual in Phase 1 Oncology Studies. JAMA Network Open, 2020, 3, e204787.	2.8	8
85	A Comparison of Two-Year Outcomes in ZUMA-1 (Axicabtagene Ciloleucel) and SCHOLAR-1 in Patients with Refractory Large B Cell Lymphoma. Blood, 2019, 134, 4095-4095.	0.6	8
86	Long-Term Survival and Gradual Recovery of B Cells in Patients with Refractory Large B Cell Lymphoma Treated with Axicabtagene Ciloleucel (Axi-Cel). Blood, 2020, 136, 40-42.	0.6	8
87	Does antithymocyte globulin have a place in reduced-intensity conditioning for allogeneic hematopoietic stem cell transplantation?. Hematology American Society of Hematology Education Program, 2012, 2012, 246-50.	0.9	8
88	Using ibrutinib in earlier lines of treatment results in better outcomes for patients with chronic lymphocytic leukemia/small lymphocytic lymphoma. Leukemia and Lymphoma, 2021, 62, 3278-3282.	0.6	7
89	Toxicities Associated With Metformin/Ritonavir Combination Treatment in Relapsed/Refractory Multiple Myeloma. Clinical Lymphoma, Myeloma and Leukemia, 2020, 20, e667-e672.	0.2	6
90	Ublituximab + TGR-1202 Demonstrates Activity and a Favorable Safety Profile in Relapsed/Refractory B-Cell NHL and High-Risk CLL: Phase I Results. Blood, 2015, 126, 1538-1538.	0.6	6

#	Article	IF	CITATIONS
91	Sarcopenia Is a Clinically Relevant and Independent Predictor of Health Outcomes after Chimeric Antigen Receptor T-Cell Therapy for Lymphoma. Blood, 2021, 138, 2502-2502.	0.6	6
92	Real-world outcomes of axicabtagene ciloleucel (Axi-cel) for the treatment of large B-cell lymphoma (LBCL) by race and ethnicity Journal of Clinical Oncology, 2022, 40, 7571-7571.	0.8	6
93	Burkitt leukemia limited to the bone marrow has a better prognosis than Burkitt lymphoma with bone marrow involvement in adults. Leukemia and Lymphoma, 2016, 57, 866-871.	0.6	5
94	A phase Ib, open label, dose escalation trial of the anti-CD37 monoclonal antibody, BI 836826, in combination with ibrutinib in patients with relapsed/refractory chronic lymphocytic leukemia. Investigational New Drugs, 2021, 39, 1099-1105.	1.2	5
95	Real-world evidence of axicabtagene ciloleucel (Axi-cel) for the treatment of large B-cell lymphoma (LBCL) in the United States (US) Journal of Clinical Oncology, 2021, 39, 7552-7552.	0.8	5
96	Estimation of the Resource Utilization and Costs of Cytokine Release Syndrome Observed in the Transcend-NHL Clinical Trial: A Micro-Costing Study. Blood, 2018, 132, 319-319.	0.6	5
97	Cirmtuzumab, an Anti-ROR1 Antibody, in Combination with Ibrutinib: Clinical Activity in Mantle Cell Lymphoma (MCL) or Chronic Lymphocytic Leukemia (CLL) from a Phase 1/2 Study. Blood, 2020, 136, 45-46.	0.6	5
98	Brentuximab Vedotin Monotherapy in DLBCL Patients with Undetectable CD30: Preliminary Results from a Phase 2 Study. Blood, 2014, 124, 629-629.	0.6	5
99	Burden of cytokine release syndrome (CRS) and neurologic events (NE) in patients (Pts) with relapsed/refractory non-Hodgkin lymphoma (NHL) receiving lisocabtagene maraleucel (Liso-cel;) Tj ETQq1 1 0.78	343 0. &rgB	T /@verlock 1
100	Ublituximab, a Novel Glycoengineered Anti-CD20 Monoclonal Antibody (mAb), in Combination with TGR-1202, a Next Generation Once Daily PI3kî´Inhibitor, Demonstrates Activity in Heavily Pre-Treated and High-Risk Chronic Lymphocytic Leukemia (CLL) and B-Cell Lymphoma. Blood, 2014, 124, 801-801.	0.6	4
101	Updated results of the ASPEN trial from a cohort of patients with <i>MYD88</i> wild-type (<i>MYD88</i> ^{WT}) Waldenström macroglobulinemia (WM) Journal of Clinical Oncology, 2020, 38, e20056-e20056.	0.8	4
102	Fixed-duration (FD) ibrutinib (I) + venetoclax (V) for first-line (1L) treatment (tx) of chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL): Three-year follow-up from the FD cohort of the phase 2 CAPTIVATE study Journal of Clinical Oncology, 2022, 40, 7519-7519.	0.8	4
103	Brentuximab Vedotin Improves HCT-CI, CR Status, and Peri-Transplant Toxicity In Patients With Relapsed/Refractory Hodgkin Lymphoma Heading To RIC Allo-HCT. Blood, 2013, 122, 3374-3374.	0.6	3
104	Double-Hit and Double-Expressor Lymphomas Are Not Associated with an Adverse Outcome after Allogeneic Stem Cell Transplantation. Blood, 2016, 128, 830-830.	0.6	3
105	ZUMA-2: Phase 2 multicenter study evaluating efficacy of kte-C19 in patients with relapsed/refractory mantle cell lymphoma Journal of Clinical Oncology, 2018, 36, TPS3102-TPS3102.	0.8	3
106	A Phase I Trial of PI3Kαδ Inhibitor Copanlisib in Combination with Nivolumab in Patients with Richter's Transformation (RT) or Transformed Non-Hodgkin Lymphoma (tNHL). Blood, 2021, 138, 3558-3558.	0.6	3
107	The Cerebroventricular Environment Reprograms Locally Infused CAR T Cells for Superior Activity Against Both CNS and Systemic B Cell Lymphoma. Blood, 2018, 132, 965-965.	0.6	2
108	Results from the International, Randomized Phase 3 Study of Ibrutinib Versus Chlorambucil in Patients 65 Years and Older with Treatment-NaÃ-ve CLL/SLL (RESONATE-2TM). Blood, 2015, 126, 495-495.	0.6	2

#	Article	IF	CITATIONS
109	Phase I Studies of Cellular Immunotherapy Using Central Memory Derived-CD19-Specific T Cells Following Autologous Stem Cell Transplantation for Patients with High-Risk Intermediate Grade B-Lineage Non-Hodgkin Lymphoma. Blood, 2015, 126, 930-930.	0.6	2
110	Ongoing complete remissions (CR) in the phase 1 of ZUMA-1: a phase 1-2 multicenter study evaluating the safety and efficacy of KTE-C19 (anti-CD19 CAR T cells) in subjects with refractory aggressive B-cell Non-Hodgkin Lymphoma (NHL) Journal of Clinical Oncology, 2016, 34, 7559-7559.	0.8	2
111	Phase 1 Biomarker Analysis of the ZUMA-1 (KTE-C19-101) Study: A Phase 1-2 Multi-Center Study Evaluating the Safety and Efficacy of Anti-CD19 CAR T Cells (KTE-C19) in Subjects with Refractory Aggressive Non-Hodgkin Lymphoma (NHL). Blood, 2015, 126, 2730-2730.	0.6	2
112	Incidence and Causes of Prolonged Hematologic Toxicity after Chimeric Antigen Receptor T Cell Therapy: A City of Hope (COH) Experience. Blood, 2020, 136, 40-41.	0.6	2
113	Phase 1/2 study of zilovertamab and ibrutinib in mantle cell lymphoma (MCL) or chronic lymphocytic leukemia (CLL) Journal of Clinical Oncology, 2022, 40, 7520-7520.	0.8	2
114	Combination of Umbralisib, Ublituximab, and Bendamustine Is Safe and Highly Active in Patients with Advanced Diffuse Large B-Cell Lymphoma and Follicular Lymphoma. Blood, 2018, 132, 4197-4197.	0.6	1
115	Final Results Of A Phase 2 Study Of Vorinostat Plus Rituximab In Newly Diagnosed, Relapsed Or Refractory Indolent Non-Hodgkin's Lymphoma. Blood, 2013, 122, 4398-4398.	0.6	1
116	Phase 1 Study of MLN8237, an Aurora KinaseA (AURKA) Inhibitor, Combined with Vorinostat, a Histone Deacetylase (HDAC) Inhibitor, in Lymphoid Malignancies. Blood, 2014, 124, 4483-4483.	0.6	1
117	Factors That Influence Treatment Decision-Making: Perspectives of 1147 Chronic Lymphocytic Leukemia (CLL) Patients in the United States. Blood, 2018, 132, 4414-4414.	0.6	1
118	Cirmtuzumab, a ROR1 Targeted Mab, Reverses Cancer Stemness, and Its Combination with Ibrutinib Is Safe and Effective: Planned Analysis of the Cirll Phase 1/2 Trial for CLL and MCL. Blood, 2019, 134, 1755-1755.	0.6	1
119	Phase 1b/2 Study of Cirmtuzumab and Ibrutinib in Mantle Cell Lymphoma (MCL) or Chronic Lymphocytic Leukemia (CLL). Blood, 2021, 138, 3534-3534.	0.6	1
120	MGA deletion Leads to Richter's Transformation Via NME1. Blood, 2021, 138, 252-252.	0.6	1
121	Atezolizumab Combined with Immunogenic Salvage Chemoimmunotherapy (R-GemOx+Atezo) in Patients with Transformed Diffuse Large B-Cell Lymphoma. Blood, 2021, 138, 1407-1407.	0.6	1
122	Chronic Lymphocytic Leukemia (CLL): Biology and Therapy. Cancer Treatment and Research, 2021, 181, 133-149.	0.2	0
123	A Phase 2 Study of Vorinostat (Suberoylanilide Hydroxamic Acid, SAHA) Plus Rituximab in Newly Diagnosed, Relapsed or Refractory Indolent Non-Hodgkin's Lymphoma. Blood, 2012, 120, 3698-3698.	0.6	0
124	Results from a Phase 1 Study and Expanded Cohort of an Interrupted Dosing Schedule of the Aurora Kinase a Inhibitor MLN8237 Combined with Vorinostat in Lymphoid Malignancies. Blood, 2015, 126, 2731-2731.	0.6	0
125	Interim Analysis of a Phase 2 Study of Bortezomib Plus Rituximab Maintenance Therapy in Patients with Mantle Cell Lymphoma Status Post Autologous Stem Cell Transplantation. Blood, 2015, 126, 1961-1961.	0.6	0

Small B-cell lymphocytic lymphoma and chronic lymphocytic leukemia. , 2017, , 47-73.

#	Article	IF	CITATIONS
127	Lymphoplasmacytic lymphoma in the era of next generation sequencing. , 2017, , 97-109.		Ο
128	Characterization of cytokine release syndrome (CRS) and neurological events (NEs) in the phase I TRANSCEND NHL 001 trial of lisocabtagene maraleucel (liso-cel) for patients (pts) with relapsed/refractory (R/R) large B-cell lymphoma (LBCL) Journal of Clinical Oncology, 2020, 38, e20046-e20046.	0.8	0
129	Pharmacologic Inhibition of SUMO-Activating Enzyme Potentiates Interferon Response and T Cell-Mediated Anti-Tumor Immunity in Chronic Lymphocytic Leukemia (CLL) and Lymphoma Models. Blood, 2021, 138, 3719-3719.	0.6	Ο
130	Interviews of Patients (Pts) with Relapsed or Refractory (R/R) Large B-Cell Lymphoma (LBCL) Enrolled in 2 Lisocabtagene Maraleucel (liso-cel) Trials on Their Treatment Experience for 3 Months after Infusion. Blood, 2021, 138, 4098-4098.	0.6	0
131	Nedd8-Activating Enzyme Inhibition Enhances Anti-Tumor Immunity and PD1 Blockade in <i>In Vivo</i> lymphoma Models. Blood, 2021, 138, 2414-2414.	0.6	Ο
132	Fixed Duration Combination Therapy with Ibrutinib (ibr) and Venetoclax (ven) Leads to Deep Responses in Relapsed/Refractory (rel/ref) Chronic Lymphocytic Leukemia (CLL): Results of a Phase 2 Study. Blood, 2021, 138, 3754-3754.	0.6	0
133	Integrative Transcriptome and Quantitative Proteome Analyses Identify METTL3 As a Key Regulator for Aberrant RNA Processing in Chronic Lymphocytic Leukemia. Blood, 2020, 136, 12-12.	0.6	Ο
134	Qualitative Analysis of the Treatment Experience and Well-Being of Patients with Relapsed/Refractory (R/R) Large B-Cell Lymphoma (LBCL) Enrolled in 2 Trials of Lisocabtagene Maraleucel (liso-cel) during the Initial Stages of Therapy. Blood, 2020, 136, 22-23.	0.6	0
135	Management of Relapsed/Refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma in the Era of Targeted Therapies. Current Hematologic Malignancy Reports, 2022, 17, 39-45.	1.2	0
136	Characterization of low-grade arthralgia, myalgia, and musculoskeletal pain with ibrutinib therapy: pooled analysis of clinical trials in patients with chronic lymphocytic leukemia and mantle cell lymphoma. Leukemia and Lymphoma, 2022, 63, 1580-1588.	0.6	0