
## **Tingting Zhou**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1405180/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Connecting the dots: Targeting the microbiome in drug toxicity. Medicinal Research Reviews, 2022, 42, 83-111.	5.0	8
2	Butyrate emerges as a crucial effector of Zhi-Zi-Chi decoctions to ameliorate depression via multiple pathways of brain-gut axis. Biomedicine and Pharmacotherapy, 2022, 149, 112861.	2.5	14
3	Soy protein degradation drives diversity of amino-containing compounds via Bacillus subtilis natto fermentation. Food Chemistry, 2022, 388, 133034.	4.2	10
4	Physcion 8-O-β-glucopyranoside ameliorates liver fibrosis through inflammation inhibition by regulating SIRT3-mediated NF-κB P65 nuclear expression. International Immunopharmacology, 2021, 90, 107206.	1.7	19
5	Metabolomics based comprehensive investigation of Gardeniae Fructus induced hepatotoxicity. Food and Chemical Toxicology, 2021, 153, 112250.	1.8	17
6	Compatibility with Semen Sojae Praeparatum attenuates hepatotoxicity of Gardeniae Fructus by regulating the microbiota, promoting butyrate production and activating antioxidant response. Phytomedicine, 2021, 90, 153656.	2.3	12
7	Isoflavones' effects on pharmacokinetic profiles of main iridoids from Gardeniae Fructus in rats. Journal of Pharmaceutical Analysis, 2020, 10, 571-580.	2.4	7
8	Predicting a Potential Link to Antidepressant Effect: Neuroprotection of Zhi-zi-chi Decoction on Glutamate-induced Cytotoxicity in PC12 Cells. Frontiers in Pharmacology, 2020, 11, 625108.	1.6	2
9	Combination of cell metabolomics and pharmacology: A novel strategy to investigate the neuroprotective effect of Zhi-zi-chi decoction. Journal of Ethnopharmacology, 2019, 236, 302-315.	2.0	11
10	A new strategy for statistical analysis-based fingerprint establishment: Application to quality assessment of Semen sojae praeparatum. Food Chemistry, 2018, 258, 189-198.	4.2	38
11	Non-isoflavones Diet Incurred Metabolic Modifications Induced by Constipation in Rats via Targeting Gut Microbiota. Frontiers in Microbiology, 2018, 9, 3002.	1.5	25
12	Quantification of isoflavone glycosides and aglycones in rat plasma by LC–MS/MS: Troubleshooting of interference from food and its application to pharmacokinetic study of Semen Sojae Praeparatum extract. Journal of Pharmaceutical and Biomedical Analysis, 2018, 161, 444-454.	1.4	15
13	Simultaneous fingerprint, quantitative analysis and anti-oxidative based screening of components in Rhizoma Smilacis Glabrae using liquid chromatography coupled with Charged Aerosol and Coulometric array Detection. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences. 2017. 1049-1050. 41-50.	1.2	9
14	Determination of Vancomycin in Human Serum by Cyclodextrin-Micellar Electrokinetic Capillary Chromatography (CD-MEKC) and Application for PDAP Patients. Molecules, 2017, 22, 538.	1.7	9
15	Quantification of a Novel Photosensitizer of Chlorin e6-C15-Monomethyl Ester in Beagle Dog Plasma Using HPLC: Application to Pharmacokinetic Studies. Molecules, 2017, 22, 693.	1.7	3
16	On-Line Organic Solvent Field Enhanced Sample Injection in Capillary Zone Electrophoresis for Analysis of Quetiapine in Beagle Dog Plasma. Molecules, 2016, 21, 121.	1.7	7
17	A comprehensive strategy using chromatographic profiles combined with chemometric methods: Application to quality control of Polygonum cuspidatum Sieb. et Zucc. Journal of Chromatography A, 2016, 1466, 67-75.	1.8	12
18	Cyclodextrin-based ultrasonic-assisted microwave extraction and HPLC-PDA-ESI-ITMSn separation and identification of hydrophilic and hydrophobic components of Polygonum cuspidatum: A green, rapid and effective process. Industrial Crops and Products, 2016, 80, 59-69.	2.5	30

Тінстінс Zhou

#	Article	IF	CITATIONS
19	Optimization of an accelerated solvent extraction dispersive liquid–liquid microextraction method for the separation and determination of essential oil from <i>Ligusticum chuanxiong</i> Hort by gas chromatography with mass spectrometry. Journal of Separation Science, 2015, 38, 3588-3598.	1.3	10
20	Chemical fingerprinting of Gardenia jasminoides Ellis by HPLC–DAD–ESI-MS combined with chemometrics methods. Food Chemistry, 2015, 188, 648-657.	4.2	68
21	Impurities preparation of sodium tanshinone IIA sulfonate by high-speed counter-current chromatography and identification by liquid chromatography/multistage tandem mass spectrometry. Journal of Chromatography A, 2013, 1288, 28-34.	1.8	15
22	Fragmentation study of iridoid glycosides including epimers by liquid chromatographyâ€diode array detection/electrospray ionization mass spectrometry and its application in metabolic fingerprint analysis of <i>Gardenia jasminoides</i> Ellis. Rapid Communications in Mass Spectrometry, 2010, 24, 2520-2528.	0.7	50
23	An efficient strategy based on MAE, HPLCâ€DADâ€ESIâ€MS/MS and 2Dâ€prepâ€HPLCâ€DAD for the rapid extract separation, identification and purification of five active coumarin components from radix angelicae dahuricae. Phytochemical Analysis, 2010, 21, 473-482.	tion, 1.2	54
24	Isolation and Purification of Three Flavonoids from the Hawthorn Leaves by High Speed Countercurrent Chromatography, Combined with Isocratic Preparative Reversed-Phase High Performance Liquid Chromatography. Journal of Liquid Chromatography and Related Technologies, 2009, 32, 2216-2231.	0.5	8
25	Isolation and purification of isoflavonoids from Rhizoma Belamcandae by twoâ€dimensional preparative highâ€performance liquid chromatography with column switch technology. Biomedical Chromatography, 2009, 23, 1064-1072.	0.8	18
26	Optimization and validation of an ionâ€pair RPâ€HPLCâ€UV method for the determination of total free iodine in rabbit plasma: application to a pharmacokinetic study. Biomedical Chromatography, 2009, 23, 1151-1159.	0.8	2
27	Rapid determination of telmisartan in human plasma by HPLC using a monolithic column with fluorescence detection and its application to a bioequivalence study. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2009, 877, 3729-3733.	1.2	29
28	Separation and determination of coumarins in Fructus cnidii extracts by pressurized capillary electrochromatography using a packed column with a monolithic outlet frit. Journal of Pharmaceutical and Biomedical Analysis, 2009, 50, 695-702.	1.4	54
29	Isolation and purification of lignans from <i>Magnolia biondii </i> Pamp by isocratic reversedâ€phase twoâ€dimensional liquid chromatography following microwaveâ€assisted extraction. Journal of Separation Science, 2007, 30, 2370-2381.	1.3	25
30	Isolation and purification of iridoid glycosides from Gardenia jasminoides Ellis by isocratic reversed-phase two-dimensional preparative high-performance liquid chromatography with column switch technology. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2007, 858, 296-301.	1.2	45
31	On-line purity monitoring in high-speed counter-current chromatography: Application of HSCCC-HPLC-DAD for the preparation of 5-HMF, neomangiferin and mangiferin from Anemarrhena asphodeloides Bunge. Journal of Pharmaceutical and Biomedical Analysis, 2007, 44, 96-100.	1.4	38
32	Qualitative and quantitative determination of ten alkaloids in traditional Chinese medicine Corydalis yanhusuo W.T. Wang by LC–MS/MS and LC–DAD. Journal of Pharmaceutical and Biomedical Analysis, 2007, 45, 219-226.	1.4	102
33	Application of high-speed counter-current chromatography coupled with high-performance liquid chromatography–diode array detection for the preparative isolation and purification of hyperoside from Hypericum perforatum with online purity monitoring. Journal of Chromatography A, 2006, 1116, 97-101.	1.8	52
34	Large-scale isolation and purification of geniposide from the fruit of Gardenia jasminoides Ellis by high-speed counter-current chromatography. Journal of Chromatography A, 2005, 1100, 76-80.	1.8	63