Smadar Gertel

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1413078/publications.pdf

Version: 2024-02-01

17 papers	318 citations	8 h-index	940134 16 g-index
18	18	18	622
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Smoke and autoimmunity: The fire behind the disease. Autoimmunity Reviews, 2016, 15, 354-374.	2.5	143
2	Hypothyroidism among SLE patients: Case–control study. Autoimmunity Reviews, 2016, 15, 484-486.	2.5	47
3	Immune Tolerance Induction with Multiepitope Peptide Derived from Citrullinated Autoantigens Attenuates Arthritis Manifestations in Adjuvant Arthritis Rats. Journal of Immunology, 2015, 194, 5674-5680.	0.4	29
4	Tolerogenic dendritic cells specific for \hat{l}^2 2-glycoprotein-l Domain-l, attenuate experimental antiphospholipid syndrome. Journal of Autoimmunity, 2014, 54, 72-80.	3.0	25
5	Tofacitinib attenuates arthritis manifestations and reduces the pathogenic CD4 T cells in adjuvant arthritis rats. Clinical Immunology, 2017, 184, 77-81.	1.4	17
6	Anti-citrullinated peptide antibodies is more than an accurate tool for diagnosis of rheumatoid arthritis. Israel Medical Association Journal, 2013, 15, 516-9.	0.1	11
7	Soluble ST2 and CXCL-10 may serve as biomarkers of subclinical diastolic dysfunction in SLE and correlate with disease activity and damage. Lupus, 2020, 29, 1430-1437.	0.8	10
8	Lymphocyte activation gene-3 (LAG-3) regulatory T cells: An evolving biomarker for treatment response in autoimmune diseases. Autoimmunity Reviews, 2022, 21, 103085.	2.5	9
9	Immunomodulation of RA Patients' PBMC with a Multiepitope Peptide Derived from Citrullinated Autoantigens. Mediators of Inflammation, 2017, 2017, 1-9.	1.4	6
10	CD4+LAG-3+ T cells are decreased in active psoriatic arthritis patients and their restoration (i) in vitro (i) is mediated by TNF inhibitors. Clinical and Experimental Immunology, 2021, 206, 173-183.	1.1	5
11	Tolerogenic citrullinated peptide for arthritis. Oncotarget, 2015, 6, 19344-19345.	0.8	4
12	T cell functions of psoriatic arthritis patients are regulated differently by TNF, IL-17A and IL-6 receptor blockades in vitro. Clinical and Experimental Rheumatology, 2022, 40, 120-128.	0.4	4
13	Reduced levels of Coco in sera of multiple sclerosis patients: A potential role in neuro-regeneration failure. Journal of Neuroimmunology, 2019, 327, 36-40.	1.1	3
14	Anticitrullinated Protein Antibodies Induce Inflammatory Gene Expression Profile in Peripheral Blood Cells from CCP–positive Patients with RA. Journal of Rheumatology, 2018, 45, 310-319.	1.0	2
15	The role of synthetic manufactured peptides containing common citrullinated epitopes in rheumatoid arthritis diagnosis. Clinical Immunology, 2019, 199, 7-11.	1.4	2
16	T cell functions of psoriatic arthritis patients are regulated differently by TNF, IL-17A and IL-6 receptor blockades in vitro. Clinical and Experimental Rheumatology, 2021, , .	0.4	1
17	THU0030â€THE DIFFERENTIAL EFFECT OF TNF-α AND IL-6R BLOCKERS ON THE EXPRESSION OF IL-17 AND ACTIVATED CD4+CD25+ T CELLS IN PATIENTS WITH PSORIATIC ARTHRITIS. , 2019, , .		0