

# Qinghua Han

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1415754/publications.pdf>

Version: 2024-02-01

12  
papers

293  
citations

1040056

9  
h-index

1199594

12  
g-index

12  
all docs

12  
docs citations

12  
times ranked

534  
citing authors

#	ARTICLE	IF	CITATIONS
1	LncRNA Tincr regulates PKC $\epsilon$ expression in a miR-31-5p-dependent manner in cardiomyocyte hypertrophy. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2020, 393, 2495-2506.	3.0	9
2	Activation of microRNA-378a-3p biogenesis promotes hepatic secretion of VLDL and hyperlipidemia by modulating ApoB100-Sortilin1 axis. <i>Theranostics</i> , 2020, 10, 3952-3966.	10.0	14
3	Trimethylamine-N-oxide (TMAO) increased aquaporin-2 expression in spontaneously hypertensive rats. <i>Clinical and Experimental Hypertension</i> , 2019, 41, 312-322.	1.3	28
4	MiR-195-5p Promotes Cardiomyocyte Hypertrophy by Targeting MFN2 and FBXW7. <i>BioMed Research International</i> , 2019, 2019, 1-10.	1.9	37
5	<p>Analysis Of Re-Hospitalizations For Patients With Heart Failure Caused By Coronary Heart Disease: Data Of First Event And Recurrent Event</p>. <i>Therapeutics and Clinical Risk Management</i> , 2019, Volume 15, 1333-1341.	2.0	5
6	CHF-PROM: validation of a patient-reported outcome measure for patients with chronic heart failure. <i>Health and Quality of Life Outcomes</i> , 2018, 16, 51.	2.4	13
7	MicroRNA-148b-3p is involved in regulating hypoxia/reoxygenation-induced injury of cardiomyocytes in vitro through modulating SIRT7/p53 signaling. <i>Chemico-Biological Interactions</i> , 2018, 296, 211-219.	4.0	33
8	Inhibition of leukotriene B4 receptor 1 attenuates lipopolysaccharide-induced cardiac dysfunction: role of AMPK-regulated mitochondrial function. <i>Scientific Reports</i> , 2017, 7, 44352.	3.3	20
9	Role of PKA in the process of neonatal cardiomyocyte hypertrophy induced by urotensin II. <i>International Journal of Molecular Medicine</i> , 2017, 40, 499-504.	4.0	10
10	MicroRNA-223-5p and -3p Cooperatively Suppress Necroptosis in Ischemic/Reperfused Hearts. <i>Journal of Biological Chemistry</i> , 2016, 291, 20247-20259.	3.4	109
11	Urotensin II induction of neonatal cardiomyocyte hypertrophy involves the CaMKII/PLN/SERCA 2a signaling pathway. <i>Gene</i> , 2016, 583, 8-14.	2.2	10
12	An investigation into the expression and mechanism of action of urotensin II in chronic pressure-overloaded rat hearts. <i>Molecular Medicine Reports</i> , 2015, 12, 6626-6634.	2.4	5