

Jens Flensted Lassen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1424141/publications.pdf>

Version: 2024-02-01

265
papers

17,292
citations

16411

64
h-index

16127

124
g-index

272
all docs

272
docs citations

272
times ranked

12586
citing authors

#	ARTICLE	IF	CITATIONS
1	Diagnostic Performance of Noninvasive Fractional Flow Reserve Derived From Coronary Computed Tomography Angiography in Suspected Coronary Artery Disease. <i>Journal of the American College of Cardiology</i> , 2014, 63, 1145-1155.	1.2	1,240
2	Remote ischaemic conditioning before hospital admission, as a complement to angioplasty, and effect on myocardial salvage in patients with acute myocardial infarction: a randomised trial. <i>Lancet, The</i> , 2010, 375, 727-734.	6.3	885
3	Randomized Study on Simple Versus Complex Stenting of Coronary Artery Bifurcation Lesions. <i>Circulation</i> , 2006, 114, 1955-1961.	1.6	666
4	Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. <i>Lancet, The</i> , 2016, 388, 2743-2752.	6.3	620
5	Prehospital Ticagrelor in ST-Segment Elevation Myocardial Infarction. <i>New England Journal of Medicine</i> , 2014, 371, 1016-1027.	13.9	538
6	System Delay and Mortality Among Patients With STEMI Treated With Primary Percutaneous Coronary Intervention. <i>JAMA - Journal of the American Medical Association</i> , 2010, 304, 763.	3.8	519
7	2-Year Clinical Outcomes After Implantation of Sirolimus-Eluting, Paclitaxel-Eluting, and Bare-Metal Coronary Stents. <i>Journal of the American College of Cardiology</i> , 2009, 53, 658-664.	1.2	316
8	Percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: updated 5-year outcomes from the randomised, non-inferiority NOBLE trial. <i>Lancet, The</i> , 2020, 395, 191-199.	6.3	280
9	Randomized Comparison of Final Kissing Balloon Dilatation Versus No Final Kissing Balloon Dilatation in Patients With Coronary Bifurcation Lesions Treated With Main Vessel Stenting. <i>Circulation</i> , 2011, 123, 79-86.	1.6	269
10	Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. <i>European Heart Journal</i> , 2005, 26, 18-26.	1.0	262
11	Classification of coronary artery bifurcation lesions and treatments: Time for a consensus!. <i>Catheterization and Cardiovascular Interventions</i> , 2008, 71, 175-183.	0.7	260
12	Coronary plaque quantification and fractional flow reserve by coronary computed tomography angiography identify ischaemia-causing lesions. <i>European Heart Journal</i> , 2016, 37, 1220-1227.	1.0	257
13	Antiplatelet Therapy for Stable Coronary Artery Disease in Atrial Fibrillation Patients Taking an Oral Anticoagulant. <i>Circulation</i> , 2014, 129, 1577-1585.	1.6	256
14	Routine Thrombectomy in Percutaneous Coronary Intervention for Acute ST-Segment Elevation Myocardial Infarction. <i>Circulation</i> , 2006, 114, 40-47.	1.6	242
15	Stent Thrombosis, Myocardial Infarction, and Death After Drug-Eluting and Bare-Metal Stent Coronary Interventions. <i>Journal of the American College of Cardiology</i> , 2007, 50, 463-470.	1.2	229
16	Effect of remote ischaemic conditioning on clinical outcomes in patients with acute myocardial infarction (CONDI-2/ERIC-PPCI): a single-blind randomised controlled trial. <i>Lancet, The</i> , 2019, 394, 1415-1424.	6.3	223
17	Invasive coronary treatment strategies for out-of-hospital cardiac arrest: a consensus statement from the European Association for Percutaneous Cardiovascular Interventions (EAPCI)/Stent for Life (SFL) groups. <i>EuroIntervention</i> , 2014, 10, 31-37.	1.4	221
18	Reduction of treatment delay in patients with ST-elevation myocardial infarction: impact of pre-hospital diagnosis and direct referral to primary percutaneous coronary intervention. <i>European Heart Journal</i> , 2005, 26, 770-777.	1.0	220

#	ARTICLE	IF	CITATIONS
19	Percutaneous coronary intervention for coronary bifurcation disease: consensus from the first 10 years of the European Bifurcation Club meetings. <i>EuroIntervention</i> , 2014, 10, 545-560.	1.4	213
20	Identification of vulnerable plaques and patients by intracoronary near-infrared spectroscopy and ultrasound (PROSPECT II): a prospective natural history study. <i>Lancet, The</i> , 2021, 397, 985-995.	6.3	208
21	Efficacy and safety of zotarolimus-eluting and sirolimus-eluting coronary stents in routine clinical care (SORT OUT III): a randomised controlled superiority trial. <i>Lancet, The</i> , 2010, 375, 1090-1099.	6.3	198
22	Percutaneous coronary intervention for the left main stem and other bifurcation lesions: 12th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2018, 13, 1540-1553.	1.4	185
23	Percutaneous coronary intervention for coronary bifurcation disease: 11th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2016, 12, 38-46.	1.4	181
24	Biolimus-eluting biodegradable polymer-coated stent versus durable polymer-coated sirolimus-eluting stent in unselected patients receiving percutaneous coronary intervention (SORT OUT V): a randomised non-inferiority trial. <i>Lancet, The</i> , 2013, 381, 661-669.	6.3	173
25	Long-Term Results After Simple Versus Complex Stenting of Coronary Artery Bifurcation Lesions. <i>Journal of the American College of Cardiology</i> , 2013, 62, 30-34.	1.2	168
26	Urban and rural implementation of pre-hospital diagnosis and direct referral for primary percutaneous coronary intervention in patients with acute ST-elevation myocardial infarction. <i>European Heart Journal</i> , 2011, 32, 430-436.	1.0	163
27	Evaluation of Coronary Artery Stenosis by Quantitative Flow Ratio During Invasive Coronary Angiography. <i>Circulation: Cardiovascular Imaging</i> , 2018, 11, e007107.	1.3	157
28	Randomized Comparison of Coronary Bifurcation Stenting With the Crush Versus the Culotte Technique Using Sirolimus Eluting Stents. <i>Circulation: Cardiovascular Interventions</i> , 2009, 2, 27-34.	1.4	156
29	Simple or Complex Stenting for Bifurcation Coronary Lesions. <i>Circulation: Cardiovascular Interventions</i> , 2011, 4, 57-64.	1.4	152
30	Randomized Comparison of Everolimus-Eluting and Sirolimus-Eluting Stents in Patients Treated With Percutaneous Coronary Intervention. <i>Circulation</i> , 2012, 125, 1246-1255.	1.6	149
31	Percutaneous coronary intervention for bifurcation coronary lesions: the 15 th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2021, 16, 1307-1317.	1.4	147
32	Cardiac arrest in the catheterisation laboratory: A 5-year experience of using mechanical chest compressions to facilitate PCI during prolonged resuscitation efforts. <i>Resuscitation</i> , 2010, 81, 383-387.	1.3	146
33	Influence of Coronary Calcification on the Diagnostic Performance of CT Angiography Derived FFR in Coronary Artery Disease. <i>JACC: Cardiovascular Imaging</i> , 2015, 8, 1045-1055.	2.3	145
34	Consensus from the 5th European Bifurcation Club meeting. <i>EuroIntervention</i> , 2010, 6, 34-38.	1.4	138
35	Randomized Comparison of Distal Protection Versus Conventional Treatment in Primary Percutaneous Coronary Intervention. <i>Journal of the American College of Cardiology</i> , 2008, 51, 899-905.	1.2	135
36	Integrated prediction of lesion-specific ischaemia from quantitative coronary CT angiography using machine learning: a multicentre study. <i>European Radiology</i> , 2018, 28, 2655-2664.	2.3	135

#	ARTICLE	IF	CITATIONS
37	Comparison of Paclitaxel- and Sirolimus-Eluting Stents in Everyday Clinical Practice. JAMA - Journal of the American Medical Association, 2008, 299, 409-16.	3.8	130
38	The European bifurcation club Left Main Coronary Stent study: a randomized comparison of stepwise provisional vs. systematic dual stenting strategies (EBC MAIN). European Heart Journal, 2021, 42, 3829-3839.	1.0	119
39	Safety and Efficacy of Everolimus- Versus Sirolimus-Eluting Stents. Journal of the American College of Cardiology, 2016, 67, 751-762.	1.2	116
40	The Stenting Coronary Arteries in Non-stress/benestent Disease (SCANDSTENT) Trial. Journal of the American College of Cardiology, 2006, 47, 449-455.	1.2	107
41	Zotarolimus-eluting durable-polymer-coated stent versus a biolimus-eluting biodegradable-polymer-coated stent in unselected patients undergoing percutaneous coronary intervention (SORT OUT VI): a randomised non-inferiority trial. Lancet, The, 2015, 385, 1527-1535.	6.3	107
42	Graphical interpretation of analytical data from comparison of a field method with a Reference Method by use of difference plots. Clinical Chemistry, 1997, 43, 2039-2046.	1.5	104
43	Randomized Comparison of a Biodegradable Polymer Ultrathin Strut Sirolimus-Eluting Stent With a Biodegradable Polymer Biolimus-Eluting Stent in Patients Treated With Percutaneous Coronary Intervention. Circulation: Cardiovascular Interventions, 2016, 9, .	1.4	104
44	Coronary bifurcation lesions treated with simple or complex stenting: 5-year survival from patient-level pooled analysis of the Nordic Bifurcation Study and the British Bifurcation Coronary Study. European Heart Journal, 2016, 37, 1923-1928.	1.0	103
45	The EBC TWO Study (European Bifurcation Coronary TWO). Circulation: Cardiovascular Interventions, 2016, 9, .	1.4	102
46	Percutaneous coronary intervention for bifurcation lesions: 2008 consensus document from the fourth meeting of the European Bifurcation Club. EuroIntervention, 2009, 5, 39-49.	1.4	102
47	Consensus from the 7th European Bifurcation Club meeting. EuroIntervention, 2013, 9, 36-45.	1.4	102
48	Renal denervation in treatment-resistant essential hypertension. A randomized, SHAM-controlled, double-blinded 24-h blood pressure-based trial. Journal of Hypertension, 2016, 34, 1639-1647.	0.3	101
49	Percutaneous coronary intervention for obstructive bifurcation lesions: the 14th consensus document from the European Bifurcation Club. EuroIntervention, 2019, 15, 90-98.	1.4	99
50	P2Y12 receptor inhibition and effect of morphine in patients undergoing primary PCI for ST-segment elevation myocardial infarction. Thrombosis and Haemostasis, 2016, 116, 369-378.	1.8	97
51	Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. Lancet, The, 2014, 383, 2047-2056.	6.3	96
52	Telemedicine used for remote prehospital diagnosing in patients suspected of acute myocardial infarction. Journal of Internal Medicine, 2002, 252, 412-420.	2.7	95
53	Percutaneous coronary intervention in left main coronary artery disease: the 13th consensus document from the European Bifurcation Club. EuroIntervention, 2018, 14, 112-120.	1.4	94
54	Infarct size and myocardial salvage after primary angioplasty in patients presenting with symptoms for <12 h vs. 12-72 h. European Heart Journal, 2009, 30, 1322-1330.	1.0	89

#	ARTICLE	IF	CITATIONS
55	Health Care System Delay and Heart Failure in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention: Follow-up of Population-Based Medical Registry Data. <i>Annals of Internal Medicine</i> , 2011, 155, 361.	2.0	81
56	Clinical Outcomes Following Coronary Bifurcation PCI Techniques. <i>JACC: Cardiovascular Interventions</i> , 2020, 13, 1432-1444.	1.1	78
57	Distance to invasive heart centre, performance of acute coronary angiography, and angioplasty and associated outcome in out-of-hospital cardiac arrest: a nationwide study. <i>European Heart Journal</i> , 2017, 38, 1645-1652.	1.0	77
58	Comparison of sirolimus-eluting and bare metal stents in coronary bifurcation lesions: Subgroup analysis of the Stenting Coronary Arteries in Non-Stress/Benestent Disease Trial (SCANDSTENT). <i>American Heart Journal</i> , 2006, 152, 1140-1145.	1.2	76
59	Long-Term Outcome After Drug-Eluting Versus Bare-Metal Stent Implantation in Patients With ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2010, 56, 641-645.	1.2	75
60	Impact of Side Branch Modeling on Computation of Endothelial Shear Stress in Coronary Artery Disease. <i>Journal of the American College of Cardiology</i> , 2015, 66, 125-135.	1.2	75
61	Electromechanical Mapping for Detection of Myocardial Viability in Patients With Ischemic Cardiomyopathy. <i>Circulation</i> , 2001, 103, 1631-1637.	1.6	74
62	Outcomes after primary percutaneous coronary intervention in octogenarians and nonagenarians with ST-segment elevation myocardial infarction: From the Western Denmark heart registry. <i>Catheterization and Cardiovascular Interventions</i> , 2013, 81, 912-919.	0.7	68
63	Quality of cardiopulmonary resuscitation in out-of-hospital cardiac arrest is hampered by interruptions in chest compressions—a nationwide prospective feasibility study. <i>Resuscitation</i> , 2011, 82, 263-269.	1.3	67
64	Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). <i>European Heart Journal</i> , 2006, 27, 267-275.	1.0	66
65	Drug-Eluting Versus Bare Metal Stents in Patients With ST-Segment Elevation Myocardial Infarction. <i>Circulation</i> , 2008, 118, 1155-1162.	1.6	66
66	International normalized ratio for prothrombin times in patients taking oral anticoagulants: critical difference and probability of significant change in consecutive measurements. <i>Clinical Chemistry</i> , 1995, 41, 444-447.	1.5	65
67	Primary PCI as the preferred reperfusion therapy in STEMI: it is a matter of time. <i>Heart</i> , 2008, 95, 362-369.	1.2	64
68	Rationale and design of the HeartFlowNXT (HeartFlow analysis of coronary blood flow using CT) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 22	0.7	64
69	FFR Derived From Coronary CT Angiography in Nonculprit Lesions of Patients With Recent STEMI. <i>JACC: Cardiovascular Imaging</i> , 2017, 10, 424-433.	2.3	64
70	Moderate overweight is beneficial and severe obesity detrimental for patients with documented atherosclerotic heart disease. <i>Heart</i> , 2013, 99, 655-660.	1.2	62
71	10-Year Clinical Outcome After Randomization to Treatment by Sirolimus- or Paclitaxel-Eluting Coronary Stents. <i>Journal of the American College of Cardiology</i> , 2017, 69, 616-624.	1.2	60
72	Intravascular ultrasound in the evaluation and treatment of left main coronary artery disease: a consensus statement from the European Bifurcation Club. <i>EuroIntervention</i> , 2018, 14, e467-e474.	1.4	60

#	ARTICLE	IF	CITATIONS
73	Side branch fractional flow reserve measurements after main vessel stenting: a Nordic-Baltic Bifurcation Study III substudy. <i>EuroIntervention</i> , 2012, 7, 1155-1161.	1.4	59
74	Timely and optimal treatment of patients with STEMI. <i>Nature Reviews Cardiology</i> , 2013, 10, 41-48.	6.1	57
75	European Bifurcation Club white paper on stenting techniques for patients with bifurcated coronary artery lesions. <i>Catheterization and Cardiovascular Interventions</i> , 2020, 96, 1067-1079.	0.7	57
76	Safety in simple versus complex stenting of coronary artery bifurcation lesions. The Nordic Bifurcation Study 14-month follow-up results. <i>EuroIntervention</i> , 2008, 4, 229-233.	1.4	56
77	Comparison of the Sirolimus-Eluting Versus Paclitaxel-Eluting Coronary Stent in Patients With Diabetes Mellitus: The Diabetes and Drug-Eluting Stent (DiabeDES) Randomized Angiography Trial—A list of participating centers and investigators appears in the Appendix.. <i>American Journal of Cardiology</i> , 2009, 103, 345-349.	0.7	55
78	Influence of Diabetes Mellitus on Clinical Outcomes Following Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. <i>American Journal of Cardiology</i> , 2012, 109, 629-635.	0.7	54
79	Impact of cardiovascular risk factors and medication use on the efficacy of remote ischaemic conditioning: post hoc subgroup analysis of a randomised controlled trial. <i>BMJ Open</i> , 2015, 5, e006923-e006923.	0.8	54
80	Randomised comparison of manual compression and FemoSeal [®] vascular closure device for closure after femoral artery access coronary angiography: the CLOSure dEvices Used in everyday Practice (CLOSE-UP) study. <i>EuroIntervention</i> , 2014, 10, 183-190.	1.4	54
81	Near-Patient Test for C-Reactive Protein in General Practice: Assessment of Clinical, Organizational, and Economic Outcomes. <i>Clinical Chemistry</i> , 1999, 45, 478-485.	1.5	53
82	Prehospital Troponin T Testing in the Diagnosis and Triage of Patients With Suspected Acute Myocardial Infarction. <i>American Journal of Cardiology</i> , 2011, 107, 1436-1440.	0.7	53
83	Prevalence and Significance of Accelerated Idioventricular Rhythm in Patients With ST-Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. <i>American Journal of Cardiology</i> , 2009, 104, 1641-1646.	0.7	52
84	Long-Term Outcome in Patients Treated With Sirolimus-Eluting Stents in Complex Coronary Artery Lesions. <i>Journal of the American College of Cardiology</i> , 2008, 51, 2011-2016.	1.2	51
85	Clinical Outcome After Crush Versus Culotte Stenting of Coronary Artery Bifurcation Lesions. <i>JACC: Cardiovascular Interventions</i> , 2013, 6, 1160-1165.	1.1	51
86	Joint consensus on the use of OCT in coronary bifurcation lesions by the European and Japanese bifurcation clubs. <i>EuroIntervention</i> , 2019, 14, e1568-e1577.	1.4	51
87	Comparison of Outcomes in Patients With Versus Without Diabetes Mellitus After Revascularization With Everolimus- and Sirolimus-Eluting Stents (from the SORT OUT IV Trial). <i>American Journal of Cardiology</i> , 2012, 110, 1585-1591.	0.7	48
88	Randomized Comparison of the Polymer-Free Biolimus-Coated BioFreedom Stent With the Ultrathin Strut Biodegradable Polymer Sirolimus-Eluting Orsiro Stent in an All-Comers Population Treated With Percutaneous Coronary Intervention. <i>Circulation</i> , 2020, 141, 2052-2063.	1.6	48
89	Dimensions of Socioeconomic Status and Clinical Outcome After Primary Percutaneous Coronary Intervention. <i>Circulation: Cardiovascular Interventions</i> , 2012, 5, 641-648.	1.4	46
90	Markers of Coagulation and Fibrinolysis as Measures of Disease Activity in Inflammatory Bowel Disease. <i>Scandinavian Journal of Gastroenterology</i> , 1998, 33, 637-643.	0.6	45

#	ARTICLE	IF	CITATIONS
91	Quantitative Point-of-Care Troponin T Measurement for Diagnosis and Prognosis in Patients With a Suspected Acute Myocardial Infarction. <i>American Journal of Cardiology</i> , 2013, 112, 1361-1366.	0.7	45
92	Fractional flow reserve derived from coronary CT angiography: Variation of repeated analyses. <i>Journal of Cardiovascular Computed Tomography</i> , 2014, 8, 307-314.	0.7	45
93	Temporal Course of Pregnancy-Associated Plasma Protein-A in Angioplasty-Treated ST-Elevation Myocardial Infarction Patients and Potential Significance of Concomitant Heparin Administration. <i>American Journal of Cardiology</i> , 2009, 103, 29-35.	0.7	44
94	Prevention of Contrast-Induced Nephropathy With N-Acetylcysteine or Sodium Bicarbonate in Patients With ST-Segment Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2014, 7, 216-224.	1.4	44
95	Interpretation of serial measurements of international normalized ratio for prothrombin times in monitoring oral anticoagulant therapy. <i>Clinical Chemistry</i> , 1995, 41, 1171-1176.	1.5	43
96	The platelet polymorphism PIA2 is a genetic risk factor for myocardial infarction. <i>Journal of Internal Medicine</i> , 2004, 255, 637-644.	2.7	43
97	3-Year Clinical Outcomes in the Randomized SORT OUT III Superiority Trial Comparing Zotarolimus- and Sirolimus-Eluting Coronary Stents. <i>JACC: Cardiovascular Interventions</i> , 2012, 5, 812-818.	1.1	43
98	Ambulance or in-catheterization laboratory administration of ticagrelor for primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: Rationale and design of the randomized, double-blind Administration of Ticagrelor in the cath Lab or in the Ambulance for New ST elevation myocardial infarction to open the Coronary artery (ATLANTIC) study. <i>American Heart Journal</i> , 2013, 165, 515-522.	1.2	43
99	Treatment of coronary bifurcation lesions, part I: implanting the first stent in the provisional pathway. The 16th expert consensus document of the European Bifurcation Club. <i>EuroIntervention</i> , 2022, 18, e362-e376.	1.4	43
100	2-Year Patient-Related Versus Stent-Related Outcomes. <i>Journal of the American College of Cardiology</i> , 2012, 60, 1140-1147.	1.2	42
101	Paclitaxel and sirolimus eluting stents versus bare metal stents: long-term risk of stent thrombosis and other outcomes. From the Western Denmark Heart Registry. <i>EuroIntervention</i> , 2010, 5, 898-905.	1.4	42
102	Treatment of coronary bifurcation lesions, part II: implanting two stents. The 16th expert consensus document of the European Bifurcation Club. <i>EuroIntervention</i> , 2022, 18, 457-470.	1.4	42
103	The Arg506Gln Mutation (FV Leiden) Among a Cohort of 4188 Unselected Danish Newborns. <i>Thrombosis Research</i> , 1998, 89, 211-215.	0.8	41
104	Long-Term Outcomes After Percutaneous Coronary Intervention in Patients With and Without Diabetes Mellitus in Western Denmark. <i>American Journal of Cardiology</i> , 2010, 105, 1513-1519.	0.7	41
105	Long-Term Outcome After Drug-Eluting Versus Bare-Metal Stent Implantation in Patients With ST-Segment Elevation Myocardial Infarction. <i>JACC: Cardiovascular Interventions</i> , 2013, 6, 548-553.	1.1	41
106	Co-registration of optical coherence tomography and X-ray angiography in percutaneous coronary intervention. The Does Optical Coherence Tomography Optimize Revascularization (DOCTOR) fusion study. <i>International Journal of Cardiology</i> , 2015, 182, 272-278.	0.8	41
107	Neointimal hyperplasia after sirolimus-eluting and paclitaxel-eluting stent implantation in diabetic patients: The Randomized Diabetes and Drug-Eluting Stent (DiabeDES) Intravascular Ultrasound Trial. <i>European Heart Journal</i> , 2008, 29, 2733-2741.	1.0	39
108	Outcome of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stent Implantation in Patients With and Without Diabetes Mellitus (a SORT OUT III Substudy). <i>American Journal of Cardiology</i> , 2011, 108, 1232-1237.	0.7	39

#	ARTICLE	IF	CITATIONS
109	Prehospital evaluation in ST-elevation myocardial infarction patients treated with primary percutaneous coronary intervention. <i>Journal of Electrocardiology</i> , 2005, 38, 187-192.	0.4	38
110	Three-dimensional multidetector computed tomography versus conventional two-dimensional transesophageal echocardiography for annular sizing in transcatheter aortic valve replacement: Influence on postprocedural paravalvular aortic regurgitation. <i>Catheterization and Cardiovascular Interventions</i> , 2013, 82, 977-986.	0.7	38
111	Increased Rate of Stent Thrombosis and Target Lesion Revascularization After Filter Protection in Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2010, 55, 867-871.	1.2	37
112	Primary Percutaneous Coronary Intervention as a National Reperfusion Strategy in Patients With ST-Segment Elevation Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2011, 4, 570-576.	1.4	37
113	Culprit only or multivessel percutaneous coronary interventions in patients with ST-segment elevation myocardial infarction and multivessel disease. <i>EuroIntervention</i> , 2012, 8, 456-464.	1.4	37
114	Quality of cardiopulmonary resuscitation in out-of-hospital cardiac arrest before and after introduction of a mechanical chest compression device, LUCAS-2; a prospective, observational study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2015, 23, 37.	1.1	36
115	Evaluation of a near-patient test for C-reactive protein used in daily routine in primary healthcare by use of difference plots. <i>Clinical Chemistry</i> , 1997, 43, 2064-2075.	1.5	35
116	Quantitative angiography methods for bifurcation lesions: a consensus statement update from the European Bifurcation Club. <i>EuroIntervention</i> , 2017, 13, 115-123.	1.4	35
117	Randomised comparison of provisional side branch stenting versus a two-stent strategy for treatment of true coronary bifurcation lesions involving a large side branch: the Nordic-Baltic Bifurcation Study IV. <i>Open Heart</i> , 2020, 7, e000947.	0.9	34
118	Comparison of Stent Thrombosis, Myocardial Infarction, and Mortality Following Drug-Eluting Versus Bare-Metal Stent Coronary Intervention in Patients With Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2008, 102, 165-172.	0.7	31
119	The Danish Heart Registry. <i>Clinical Epidemiology</i> , 2016, Volume 8, 503-508.	1.5	31
120	Effect of Pre-Hospital Ticagrelor During the First 24 h After Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. <i>JACC: Cardiovascular Interventions</i> , 2016, 9, 646-656.	1.1	31
121	Differences between the left main and other bifurcations. <i>EuroIntervention</i> , 2015, 11, V106-V110.	1.4	31
122	Sex- and age-related differences in clinical outcome after primary percutaneous coronary intervention. <i>EuroIntervention</i> , 2012, 8, 904-911.	1.4	31
123	Clinical Outcome After Primary Percutaneous Coronary Intervention With Drug-Eluting and Bare Metal Stents in Patients With ST-Segment Elevation Myocardial Infarction. <i>Circulation: Cardiovascular Interventions</i> , 2008, 1, 176-184.	1.4	30
124	The potential of optimizing prehospital triage of patients with suspected acute myocardial infarction using high-sensitivity cardiac troponin T and copeptin. <i>Biomarkers</i> , 2017, 22, 351-360.	0.9	30
125	Three-Year Outcomes After Revascularization With Everolimus- and Sirolimus-Eluting Stents From the SORT OUT IV Trial. <i>JACC: Cardiovascular Interventions</i> , 2014, 7, 840-848.	1.1	28
126	ST changes before and during primary percutaneous coronary intervention predict final infarct size in patients with ST elevation myocardial infarction. <i>Journal of Electrocardiology</i> , 2009, 42, 64-72.	0.4	27

#	ARTICLE	IF	CITATIONS
127	Intravascular ultrasound assessed incomplete stent apposition and stent fracture in stent thrombosis after bare metal versus drug-eluting stent treatment the Nordic Intravascular Ultrasound Study (NIVUS). <i>International Journal of Cardiology</i> , 2013, 168, 1010-1016.	0.8	27
128	Clopidogrel discontinuation within the first year after coronary drug-eluting stent implantation: an observational study. <i>BMC Cardiovascular Disorders</i> , 2014, 14, 100.	0.7	27
129	Everolimus-Eluting Versus Biolimus-Eluting Stents With Biodegradable Polymers in Unselected Patients Undergoing Percutaneous Coronary Intervention. <i>JACC: Cardiovascular Interventions</i> , 2019, 12, 624-633.	1.1	27
130	Gains and losses of warfarin therapy as performed in an anticoagulation clinic. <i>Journal of Internal Medicine</i> , 2006, 259, 296-304.	2.7	26
131	Percutaneous coronary intervention of bifurcation lesions: state-of-the-art. Insights from the second meeting of the European Bifurcation Club. <i>EuroIntervention</i> , 2007, 3, 44-9.	1.4	26
132	Biological variation of International Normalized Ratio for prothrombin times, and consequences in monitoring oral anticoagulant therapy: computer simulation of serial measurements with goal-setting for analytical quality. <i>Clinical Chemistry</i> , 1997, 43, 2175-2182.	1.5	25
133	Influence of a Pressure Gradient Distal to Implanted Bare-Metal Stent on In-Stent Restenosis After Percutaneous Coronary Intervention. <i>Circulation</i> , 2007, 116, 2802-2808.	1.6	25
134	Impact of Health Care System Delay in Patients With ST-Elevation Myocardial Infarction on Return to Labor Market and Work Retirement. <i>American Journal of Cardiology</i> , 2014, 114, 1810-1816.	0.7	25
135	A randomized controlled trial of shared care versus routine care for patients receiving oral anticoagulant therapy. <i>Journal of Internal Medicine</i> , 2002, 252, 322-331.	2.7	23
136	Does Postsystolic Motion or Shortening Predict Recovery of Myocardial Function After Primary Percutaneous Coronary Intervention?. <i>Journal of the American Society of Echocardiography</i> , 2007, 20, 505-511.	1.2	23
137	Outcome in high risk patients with unprotected left main coronary artery stenosis treated with percutaneous coronary intervention. <i>Catheterization and Cardiovascular Interventions</i> , 2010, 75, 101-108.	0.7	23
138	Support with intra-aortic balloon pump vs. Impella2.5® and blood flow to the heart, brain and kidneys – An experimental porcine model of ischaemic heart failure. <i>International Journal of Cardiology</i> , 2015, 178, 153-158.	0.8	23
139	Myocardial Perfusion Imaging Versus Computed Tomography Angiography – Derived Fractional Flow Reserve Testing in Stable Patients With Intermediate-Range Coronary Lesions: Influence on Downstream Diagnostic Workflows and Invasive Angiography Findings. <i>Journal of the American Heart Association</i> , 2017, 6, .	1.6	23
140	Morphine and Ticagrelor Interaction in Primary Percutaneous Coronary Intervention in ST-Segment Elevation Myocardial Infarction: ATLANTIC-Morphine. <i>American Journal of Cardiovascular Drugs</i> , 2019, 19, 173-183.	1.0	23
141	Late lumen loss and intima hyperplasia after sirolimus-eluting and zotarolimus-eluting stent implantation in diabetic patients: the diabetes and drug-eluting stent (DiabeDES III) angiography and intravascular ultrasound trial. <i>EuroIntervention</i> , 2011, 7, 323-331.	1.4	23
142	Earlier reperfusion in patients with ST-elevation Myocardial infarction by use of helicopter. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2012, 20, 70.	1.1	21
143	Event detection using population-based health care databases in randomized clinical trials: a novel research tool in interventional cardiology. <i>Clinical Epidemiology</i> , 2013, 5, 357.	1.5	21
144	Influence of multivessel disease with or without additional revascularization on mortality in patients with ST-segment elevation myocardial infarction. <i>American Heart Journal</i> , 2015, 170, 70-78.	1.2	21

#	ARTICLE	IF	CITATIONS
145	Intravenous magnesium does not influence the activity of the coagulation cascade. <i>Blood Coagulation and Fibrinolysis</i> , 2001, 12, 223-228.	0.5	20
146	Is there a hypercoagulable phase during initiation of antithrombotic therapy with oral anticoagulants in patients with atrial fibrillation?. <i>Thrombosis Research</i> , 2003, 109, 241-246.	0.8	20
147	One-year clinical and angiographic results of hybrid coronary revascularization. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2015, 150, 1181-1186.	0.4	20
148	Danegaptide for primary percutaneous coronary intervention in acute myocardial infarction patients: a phase 2 randomised clinical trial. <i>Heart</i> , 2018, 104, 1593-1599.	1.2	20
149	Diabetes and risk of peripheral artery disease in patients undergoing first-time coronary angiography between 2000 and 2012 – a nationwide study. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 234.	0.7	20
150	Randomized clinical trial. <i>European Journal of Gastroenterology and Hepatology</i> , 2017, 29, 1118-1125.	0.8	19
151	Rationale and Design of a Randomized Clinical Comparison of Everolimus-Eluting (Xience V/Promus) and Sirolimus-Eluting (Cypher Select+) Coronary Stents in Unselected Patients with Coronary Heart Disease. <i>Cardiology</i> , 2010, 116, 73-78.	0.6	18
152	Impact of presentation and transfer delays on complete ST-segment resolution before primary percutaneous coronary intervention: insights from the ATLANTIC trial. <i>EuroIntervention</i> , 2017, 13, 69-77.	1.4	18
153	HbA1c in an unselected population of 4438 people with Type 2 diabetes in a Danish county. <i>Scandinavian Journal of Primary Health Care</i> , 2001, 19, 241-246.	0.6	17
154	Recruitable collateral blood flow index predicts coronary in-stent restenosis after percutaneous coronary intervention. <i>European Heart Journal</i> , 2007, 28, 1820-1826.	1.0	17
155	Comparison of Outcomes of Patients ≥ 80 Years of Age Having Percutaneous Coronary Intervention According to Presentation (Stable vs Unstable Angina Pectoris/Non-ST-Segment Elevation Myocardial Infarction). <i>Journal of the American College of Cardiology</i> , 2014, 64, 1395-1400.	0.7	17
156	Randomized comparison of a sirolimus-eluting Orsiro stent with a biolimus-eluting Nobori stent in patients treated with percutaneous coronary intervention: Rationale and study design of the Scandinavian Organization for Randomized Trials with Clinical Outcome VII trial. <i>American Heart Journal</i> , 2015, 170, 210-215.	1.2	17
157	The impact of distal embolization and distal protection on long-term outcome in patients with ST elevation myocardial infarction randomized to primary percutaneous coronary intervention – results from a randomized study. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015, 4, 180-188.	0.4	17
158	Two-year outcome after biodegradable polymer sirolimus- and biolimus-eluting coronary stents (from the OVERLOCK II trial). <i>Journal of the American College of Cardiology</i> , 2014, 64, 1395-1400.	1.4	17
159	The risk and prognostic impact of definite stent thrombosis or in-stent restenosis after coronary stent implantation. <i>EuroIntervention</i> , 2012, 8, 591-598.	1.4	17
160	Target lesion revascularisation in patients treated with a sirolimus-eluting or paclitaxel-eluting stent. <i>Heart</i> , 2007, 93, 694-697.	1.2	16
161	Serial Intravascular Ultrasound Analysis of Peri-Stent Remodeling and Proximal and Distal Edge Effects After Sirolimus-Eluting or Paclitaxel-Eluting Stent Implantation in Patients With Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2009, 103, 1083-1088.	0.7	16
162	2012 ESC STEMI guidelines and reperfusion therapy. <i>Heart</i> , 2013, 99, 1154-1156.	1.2	16

#	ARTICLE	IF	CITATIONS
163	Quantitative coronary analysis in the Nordic Bifurcation studies. International Journal of Cardiovascular Imaging, 2011, 27, 175-180.	0.7	15
164	Long-Term Outcome of Sirolimus-Eluting and Zotarolimus-Eluting Coronary Stent Implantation in Patients With and Without Diabetes Mellitus (A Danish Organization for Randomized Trials on) Tj ETQq0 0 0 rgBT /0verlock 10 Tf 50 69	0.7	15
165	Influence of preinfarction angina and coronary collateral blood flow on the efficacy of remote ischaemic conditioning in patients with ST segment elevation myocardial infarction: post hoc subgroup analysis of a randomised controlled trial. BMJ Open, 2016, 6, e013314.	0.8	15
166	C-reactive protein in general practice - how commonly is it used and why?. Scandinavian Journal of Primary Health Care, 1997, 15, 35-38.	0.6	14
167	Intravascular Ultrasound Assessment of Expansion of the Sirolimus-Eluting (Cypher Select) and Paclitaxel-Eluting (Taxus Express-2) Stent in Patients With Diabetes Mellitus. American Journal of Cardiology, 2008, 102, 19-26.	0.7	14
168	Comparison of Outcome of Patients With ST-Segment Elevation Myocardial Infarction and Complete Versus Incomplete ST-Resolution Before Primary Percutaneous Coronary Intervention. American Journal of Cardiology, 2016, 117, 1735-1740.	0.7	14
169	Quality of lipid-lowering therapy in patients with ischaemic heart disease: a register-based study in 3477 patients. Journal of Internal Medicine, 2004, 255, 367-372.	2.7	13
170	Percutaneous treatment of coronary bifurcation lesions: a novel "extended Y" technique with complete lesion stent coverage. Heart, 2006, 92, 981-982.	1.2	13
171	Incidence of definite stent thrombosis or inâ€stent restenosis after drugâ€eluting stent implantation for treatment of coronary inâ€stent restenosis: From Western Denmark heart registry. Catheterization and Cardiovascular Interventions, 2013, 81, 260-265.	0.7	13
172	Patient-specific computational simulation of coronary artery bifurcation stenting. Scientific Reports, 2021, 11, 16486.	1.6	13
173	Cholesterol reduction following health screening in general practice. Scandinavian Journal of Primary Health Care, 2002, 20, 219-223.	0.6	12
174	Pain and discomfort in closure of femoral access coronary angiography. The CLOSuredEvices Used in everyday Practice (CLOSE-UP) pain sub study. European Journal of Cardiovascular Nursing, 2014, 13, 221-226.	0.4	12
175	Implantable cardioverter defibrillator and survival after out-of-hospital cardiac arrest due to acute myocardial infarction in Denmark in the years 2001â€2012, a nationwide study. European Heart Journal: Acute Cardiovascular Care, 2017, 6, 144-154.	0.4	12
176	Assessing the cardiology community position on transradial intervention and the use of bivalirudin in patients with acute coronary syndrome undergoing invasive management: results of an EAPCI survey. EuroIntervention, 2016, 12, 1154-1163.	1.4	12
177	Evaluation of a Simple Dosage Scheme for Transition from Phenprocoumon to Warfarin in Oral Anticoagulation. Thrombosis Research, 2000, 98, 157-163.	0.8	11
178	Prospective evaluation of the quality of oral anticoagulation management in an outpatient clinic and in general practices. Thrombosis Research, 2002, 105, 103-108.	0.8	11
179	Similar five-year outcome with paclitaxel- and sirolimus-eluting coronary stents. Scandinavian Cardiovascular Journal, 2014, 48, 148-155.	0.4	11
180	Longâ€term outcome following percutaneous coronary intervention with drugâ€eluting stents compared with bareâ€metal stents in saphenous vein graft lesions: From Western Denmark heart registry. Catheterization and Cardiovascular Interventions, 2014, 83, 1035-1042.	0.7	11

#	ARTICLE	IF	CITATIONS
181	A nationwide investigation of CPR courses, books, and skill retention. <i>Resuscitation</i> , 2019, 134, 110-121.	1.3	11
182	Interaction of ischaemic postconditioning and thrombectomy in patients with ST-elevation myocardial infarction. <i>Heart</i> , 2020, 106, 24-32.	1.2	11
183	Efficacy and Safety of Glycoprotein IIb/IIIa Inhibitors on Top of Ticagrelor in STEMI: A Subanalysis of the ATLANTIC Trial. <i>Thrombosis and Haemostasis</i> , 2020, 120, 065-074.	1.8	11
184	Final five-year outcomes after implantation of biodegradable polymer-coated biolimus-eluting stents versus durable polymer-coated sirolimus-eluting stents. <i>EuroIntervention</i> , 2017, 13, 1336-1344.	1.4	11
185	Outcome of unprotected left main percutaneous coronary intervention in surgical low-risk, surgical high-risk, and acute myocardial infarction patients. <i>EuroIntervention</i> , 2006, 1, 403-8.	1.4	11
186	Are we underestimating the full potential of early thrombolytic treatment in patients with acute myocardial infarction?. <i>British Heart Journal</i> , 2003, 89, 483-484.	2.2	10
187	Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. <i>Journal of Nuclear Cardiology</i> , 2009, 16, 784-791.	1.4	10
188	Zotarolimus-eluting vs. sirolimus-eluting coronary stents in patients with and without acute coronary syndromes: a SORT OUT III substudy. <i>European Journal of Clinical Investigation</i> , 2012, 42, 1047-1054.	1.7	10
189	Safety of therapeutic hypothermia combined with primary percutaneous coronary intervention after out-of-hospital cardiac arrest. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015, 4, 60-63.	0.4	10
190	Duration of ischemia and treatment effects of pre- versus in-hospital ticagrelor in patients with ST-segment elevation myocardial infarction: Insights from the ATLANTIC study. <i>American Heart Journal</i> , 2018, 196, 56-64.	1.2	10
191	Association between socioeconomic factors and ICD implantation in a publicly financed health care system: a Danish nationwide study. <i>Europace</i> , 2018, 20, 1129-1137.	0.7	10
192	Comparison of the polymer-free biolimus-coated BioFreedom stent with the thin-strut biodegradable polymer sirolimus-eluting Orsiro stent in an all-comers population treated with percutaneous coronary intervention: Rationale and design of the randomized SORT OUT IX trial. <i>American Heart Journal</i> , 2019, 213, 1-7.	1.2	10
193	Stepwise visualisation of a provisional bifurcation stenting procedure – multimodal visualisation within a reanimated human heart utilising Visible Heart methodologies. <i>EuroIntervention</i> , 2020, 16, e734-e737.	1.4	10
194	Clinical outcomes three-year after revascularization with biodegradable polymer stents: ultrathin-strut sirolimus-eluting stent versus biolimus-eluting stent: from the Scandinavian organization for randomized trials with clinical outcome VII trial. <i>Coronary Artery Disease</i> , 2020, 31, 485-492.	0.3	9
195	Feasibility and early safety of hybrid coronary revascularisation combining off-pump coronary surgery through J-hemisternotomy with percutaneous coronary intervention. <i>EuroIntervention</i> , 2015, 10, e1-e6.	1.4	9
196	Use of three-dimensional optical coherence tomography to verify correct wire position in a jailed side branch after main vessel stent implantation. <i>EuroIntervention</i> , 2011, 7, 528-529.	1.4	9
197	Myocardial infarction centres: the way forward. <i>Heart</i> , 2005, 91, iii12-iii15.	1.2	8
198	Nonsteroidal Antiinflammatory Drug Use and Cardiovascular Risks After Coronary Stent Implantation. <i>Pharmacotherapy</i> , 2011, 31, 458-468.	1.2	8

#	ARTICLE	IF	CITATIONS
199	Intimal hyperplasia and vascular remodeling after everolimus-eluting and sirolimus-eluting stent implantation in diabetic patients: The randomized diabetes and drug-eluting stent (DiabeDES) IV intravascular ultrasound trial. <i>Catheterization and Cardiovascular Interventions</i> , 2014, 83, 864-872.	0.7	8
200	Use of renal replacement therapy after out-of-hospital cardiac arrest in Denmark 2005–2013. <i>Scandinavian Cardiovascular Journal</i> , 2018, 52, 238-243.	0.4	8
201	Ten-year clinical outcome of patients treated with a drug-eluting stent in the proximal left anterior descending artery segment compared with patients stented in other non-left main coronary segments. <i>EuroIntervention</i> , 2018, 14, 764-771.	1.4	8
202	Asia Pacific consensus document on coronary bifurcation interventions. <i>EuroIntervention</i> , 2020, 16, e706-e714.	1.4	8
203	Outcomes after revascularisation with everolimus- and sirolimus-eluting stents in patients with acute coronary syndromes and stable angina pectoris: a substudy of the SORT OUT IV trial. <i>EuroIntervention</i> , 2014, 10, 212-223.	1.4	8
204	Hallucinogenic mushroom use by Danish students: pattern of consumption. <i>Journal of Internal Medicine</i> , 1993, 233, 111-112.	2.7	7
205	Should patients with acute ST elevation MI be transferred for primary PCI?. <i>Heart</i> , 2004, 90, 1358-1363.	1.2	7
206	Haemostatic Activity in Patients with Atrial Fibrillation Treated with Low-Molecular-Weight Heparin Before and After Electrical Cardioversion. <i>Journal of Thrombosis and Thrombolysis</i> , 2004, 17, 185-189.	1.0	7
207	Treatment delays in ST elevation myocardial infarction. <i>BMJ: British Medical Journal</i> , 2008, 336, 401-402.	2.4	7
208	Comparison between patients included in randomized controlled trials of ischemic heart disease and real-world data. A nationwide study. <i>American Heart Journal</i> , 2018, 204, 128-138.	1.2	7
209	Mechanical performance and healing patterns of the novel sirolimus-eluting bioresorbable Fantom scaffold: 6-month and 9-month follow-up by optical coherence tomography in the FANTOM II study. <i>Open Heart</i> , 2019, 6, e000941.	0.9	7
210	Assessment of single and double coronary bifurcation stenting techniques using multimodal imaging and 3D modeling in reanimated swine hearts using Visible Heart® methodologies. <i>International Journal of Cardiovascular Imaging</i> , 2021, 37, 2591-2601.	0.7	7
211	Research, technology assessment, and quality assurance. <i>European Journal of General Practice</i> , 1996, 2, 162-165.	0.9	6
212	Effect of Anticoagulant Therapy on the Hypercoagulable State in Patients Carrying the Factor V Arg506Gln Mutation. <i>Thrombosis Research</i> , 1998, 92, 157-162.	0.8	6
213	Left ventricular contractile function after distal protection in primary percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2011, 146, 395-398.	0.8	6
214	A “normal”-invasive coronary angiogram may not be normal. <i>Journal of Cardiovascular Computed Tomography</i> , 2015, 9, 264-266.	0.7	6
215	Ten-year outcomes from a randomised comparison of zotarolimus-eluting and sirolimus-eluting stents: the SORT OUT III study. <i>EuroIntervention</i> , 2019, 15, e1022-e1024.	1.4	6
216	Technical aspects of the culotte technique. <i>EuroIntervention</i> , 2015, 11, V99-V101.	1.4	6

#	ARTICLE	IF	CITATIONS
217	When and how to use BRS in bifurcations?. <i>EuroIntervention</i> , 2015, 11, V185-V187.	1.4	6
218	Coronary Edema Demonstrated by Cardiovascular Magnetic Resonance in Patients With Peri-Stent Inflammation and Aneurysm Formation After Treatment by Drug-Eluting Stents. <i>Circulation: Cardiovascular Imaging</i> , 2013, 6, 352-354.	1.3	5
219	A response to a misrepresentation of the STEMI guidelines: the response. <i>Heart</i> , 2013, 99, 1787-1788.	1.2	5
220	The Extent of Myocardial Injury During Prolonged Targeted Temperature Management After Out-of-Hospital Cardiac Arrest. <i>American Journal of Medicine</i> , 2017, 130, 37-46.	0.6	5
221	Randomized comparison of sirolimus eluting, and biolimus eluting bioresorbable polymer stents: the SORT-OUT VII optical coherence tomography study. <i>European Heart Journal Cardiovascular Imaging</i> , 2018, 19, 329-338.	0.5	5
222	Unreported exclusion and sampling bias in interpretation of randomized controlled trials in patients with STEMI. <i>International Journal of Cardiology</i> , 2019, 289, 1-5.	0.8	5
223	Ten-Year Outcomes of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stents in Patients With Versus Without Diabetes Mellitus (SORT OUT III). <i>American Journal of Cardiology</i> , 2020, 125, 349-353.	0.7	5
224	The association between self-reported health status and adverse events: a comparison among coronary artery bypass grafting (CABG) versus percutaneous coronary intervention (PCI). <i>Quality of Life Research</i> , 2020, 29, 3017-3029.	1.5	5
225	Clinical outcome following late reperfusion with percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2020, , .	0.4	5
226	Trends in Use of Cardioprotective Medication in Peripheral Artery Disease: A Nationwide Study. <i>Journal of the American Heart Association</i> , 2021, 10, e020333.	1.6	5
227	Intravascular ultrasound findings of the Fantom sirolimus-eluting bioresorbable scaffold at six- and nine-month follow-up: the FANTOM II study. <i>EuroIntervention</i> , 2018, 14, e1215-e1223.	1.4	5
228	One or two stents for coronary bifurcation lesions?. <i>EuroIntervention</i> , 2010, 6, J61-J64.	1.4	5
229	Definitions and Standardized Endpoints for Treatment of Coronary Bifurcations. <i>EuroIntervention</i> , 2023, 19, e807-e831.	1.4	5
230	From science to everyday clinical practice. Need for systematic evaluation of research findings. <i>Scandinavian Journal of Primary Health Care</i> , 1999, 17, 6-10.	0.6	4
231	Invasive angiography and revascularization in patients with stable angina following prior coronary artery bypass grafting: Results from the East Denmark heart registry. <i>Catheterization and Cardiovascular Interventions</i> , 2017, 89, 341-349.	0.7	4
232	Thrombus aspiration and prehospital ticagrelor administration in ST-elevation myocardial infarction: Findings from the ATLANTIC trial. <i>American Heart Journal</i> , 2018, 196, 1-8.	1.2	4
233	Pre-hospital administration of ticagrelor in diabetic patients with ST-elevation myocardial infarction undergoing primary angioplasty: A sub-analysis of the ATLANTIC trial. <i>Catheterization and Cardiovascular Interventions</i> , 2019, 93, E369-E377.	0.7	4
234	Direct Visualization of TAVR-Related Coronary Artery Management Techniques in Reanimated Beating Hearts. <i>JACC: Cardiovascular Interventions</i> , 2021, 14, e87-e91.	1.1	4

#	ARTICLE	IF	CITATIONS
235	Thin- versus thick-strut polymer-free biolimus-eluting stents: the BioFreedom QCA randomised trial. <i>EuroIntervention</i> , 2021, 17, 233-239.	1.4	4
236	Denmark: coronary and structural heart interventions from 2010 to 2015. <i>EuroIntervention</i> , 2017, 13, Z17-Z20.	1.4	4
237	Impact of age on the effect of pre-hospital P2Y12 receptor inhibition in primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: the ATLANTIC-Elderly analysis. <i>EuroIntervention</i> , 2018, 14, 789-797.	1.4	4
238	Coronary bifurcation treatment revisited. <i>EuroIntervention</i> , 2015, 11, 850-851.	1.4	4
239	Reduction of treatment delay in patients with ST-elevation myocardial infarction: impact of pre-hospital diagnosis and direct referral to primary percutaneous intervention. <i>European Heart Journal</i> , 2005, 26, 1343-1343.	1.0	3
240	Acquired peri-stent evaginations in a second generation durable polymer drug eluting stent. <i>Cardiovascular Revascularization Medicine</i> , 2013, 14, 246-247.	0.3	3
241	Pre-hospital ticagrelor in ST-segment elevation myocardial infarction in the French ATLANTIC population. <i>International Journal of Cardiology</i> , 2017, 244, 49-53.	0.8	3
242	Sub-acute cardiac magnetic resonance to predict irreversible reduction in left ventricular ejection fraction after ST-segment elevation myocardial infarction: A DANAMI-3 sub-study. <i>International Journal of Cardiology</i> , 2020, 301, 215-219.	0.8	3
243	The <i>EuroIntervention</i> coronary bifurcation treatment supplement. <i>EuroIntervention</i> , 2015, 11, V9-V11.	1.4	3
244	Zotarolimus-eluting versus sirolimus-eluting coronary stent implantation. <i>Interventional Cardiology</i> , 2010, 2, 807-812.	0.0	2
245	Long-term safety and efficacy of drug-eluting stents. <i>Lancet, The</i> , 2011, 377, 1213-1214.	6.3	2
246	Comparison of zotarolimus-eluting and sirolimus-eluting coronary stents: a study from the Western Denmark Heart Registry. <i>BMC Cardiovascular Disorders</i> , 2012, 12, 84.	0.7	2
247	TCT-10 Physiological testing of coronary artery stenosis by computation of invasive coronary angiography. The wire-free functional imaging (WIFI-II) study. <i>Journal of the American College of Cardiology</i> , 2016, 68, B4-B5.	1.2	2
248	Neurological prognostication tools in out-of-hospital cardiac arrest patients in Danish intensive care units from 2005 to 2013. <i>Acta Anaesthesiologica Scandinavica</i> , 2018, 62, 1412-1420.	0.7	2
249	Time trends in the risk of atrial fibrillation and ischaemic stroke in patients with peripheral artery disease between 1997 and 2015. <i>Open Heart</i> , 2020, 7, e001185.	0.9	2
250	When is a two-stent technique needed and which technique should then be used for bifurcation coronary lesions?. <i>EuroIntervention</i> , 2015, 11, V96-V98.	1.4	2
251	Optical Coherence Tomography- Versus Angiography-Guided Magnesium Bioresorbable Scaffold Implantation in NSTEMI Patients. <i>Cardiovascular Revascularization Medicine</i> , 2022, 40, 101-110.	0.3	2
252	Two stent or not two stent--that is the question. <i>European Heart Journal</i> , 2008, 29, 2829-2830.	1.0	1

#	ARTICLE	IF	CITATIONS
253	Sirolimus-eluting versus bare-metal stent implantation in patients with ostial lesions. International Journal of Cardiology, 2010, 145, 162-163.	0.8	1
254	TCT-297 Stent edge dissections detected by Optical Coherence Tomography, incidence, predictors and 12-month outcome. Journal of the American College of Cardiology, 2012, 60, B84.	1.2	1
255	TCT-318 Ten-year All-cause Mortality after Simple versus Complex Stenting of Coronary Artery Bifurcation Lesions in the Randomized Nordic Bifurcation Study. Journal of the American College of Cardiology, 2016, 68, B131-B132.	1.2	1
256	When analyses are invalidated by erroneous assumptions. International Journal of Technology Assessment in Health Care, 2005, 21, 414-416.	0.2	0
257	TCT-347 Clinical Outcome Of Biolimus-Eluting Versus Sirolimus-Eluting Coronary Stent Implantation In Patients With And Without Diabetes Mellitus: A SORT OUT V Substudy. Journal of the American College of Cardiology, 2012, 60, B98-B99.	1.2	0
258	Clinical outcomes after treatment of multiple lesions with zotarolimus-eluting versus sirolimus-eluting coronary stents (a SORT OUT III substudy). BMC Cardiovascular Disorders, 2012, 12, 18.	0.7	0
259	TCT-321 Definite and probable stent thrombosis after revascularization with drug-eluting stents with a biodegradable polymer. From the randomized SORT OUT VII Trial. Journal of the American College of Cardiology, 2016, 68, B133.	1.2	0
260	INTERACTION BETWEEN MORPHINE AND TICAGRELOR IN THE ATLANTIC STUDY. Journal of the American College of Cardiology, 2016, 67, 545.	1.2	0
261	Coronary Artery Healing Process after Bioresorbable Scaffold in Patients with Non-ST-Segment Elevation Myocardial Infarction: Rationale, Design, and Methodology of the HONEST Study. Cardiology, 2021, 146, 1-11.	0.6	0
262	Treatment of Bifurcation Lesions by Bail-Out TAP or Culotte: Lost in Translation?. Reviews on Recent Clinical Trials, 2017, 12, 212-215.	0.4	0
263	Applications of computational simulation in bifurcation stenting: past, present and future. EuroIntervention, 2020, 16, e698-e700.	1.4	0
264	"The significant other": Evaluation of side branch ostial compromise in bifurcation stenting. Cardiology Journal, 2020, 27, 474-477.	0.5	0
265	MO445: Implications of Cardiac Biomarkers in Patients With Renal Insufficiency on Probability of Coronary Angiography and Subsequent Cardiovascular Outcomes. Nephrology Dialysis Transplantation, 2022, 37, .	0.4	0