

Jens Flensted Lassen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1424141/jens-flensted-lassen-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

252
papers

13,133
citations

57
h-index

108
g-index

271
ext. papers

15,582
ext. citations

6.2
avg, IF

5.69
L-index

#	Paper	IF	Citations
252	Percutaneous coronary intervention for bifurcation coronary lesions: the 15 consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2021 , 16, 1307-1317	3.1	45
251	Identification of vulnerable plaques and patients by intracoronary near-infrared spectroscopy and ultrasound (PROSPECT II): a prospective natural history study. <i>Lancet, The</i> , 2021 , 397, 985-995	4.0	45
250	Direct Visualization of TAVR-Related Coronary Artery Management Techniques in Reanimated Beating Hearts. <i>JACC: Cardiovascular Interventions</i> , 2021 , 14, e87-e91	5	2
249	Assessment of single and double coronary bifurcation stenting techniques using multimodal imaging and 3D modeling in reanimated swine hearts using Visible Heart [®] methodologies. <i>International Journal of Cardiovascular Imaging</i> , 2021 , 37, 2591-2601	2.5	1
248	The European bifurcation club Left Main Coronary Stent study: a randomized comparison of stepwise provisional vs. systematic dual stenting strategies (EBC MAIN). <i>European Heart Journal</i> , 2021 , 42, 3829-3839	9.5	20
247	Thin- versus thick-strut polymer-free biolimus-eluting stents: the BioFreedom QCA randomised trial. <i>EuroIntervention</i> , 2021 , 17, 233-239	3.1	1
246	Coronary Artery Healing Process after Bioresorbable Scaffold in Patients with Non-ST-Segment Elevation Myocardial Infarction: Rationale, Design, and Methodology of the HONEST Study. <i>Cardiology</i> , 2021 , 146, 161-171	1.6	
245	Trends in Use of Cardioprotective Medication in Peripheral Artery Disease: A Nationwide Study. <i>Journal of the American Heart Association</i> , 2021 , 10, e020333	6	1
244	Patient-specific computational simulation of coronary artery bifurcation stenting. <i>Scientific Reports</i> , 2021 , 11, 16486	4.9	2
243	Clinical outcome following late reperfusion with percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2020 ,	4.3	1
242	Randomized Comparison of the Polymer-Free Biolimus-Coated BioFreedom Stent With the Ultrathin Strut Biodegradable Polymer Sirolimus-Eluting Orsiro Stent in an All-Comers Population Treated With Percutaneous Coronary Intervention: The SORT OUT IX Trial. <i>Circulation</i> , 2020 , 141, 2052-2063	16.7	23
241	European Bifurcation Club white paper on stenting techniques for patients with bifurcated coronary artery lesions. <i>Catheterization and Cardiovascular Interventions</i> , 2020 , 96, 1067-1079	2.7	17
240	Clinical Outcomes Following Coronary Bifurcation PCI Techniques: A Systematic Review and Network Meta-Analysis Comprising 5,711 Patients. <i>JACC: Cardiovascular Interventions</i> , 2020 , 13, 1432-1444	5.4	19
239	Randomised comparison of provisional side branch stenting versus a two-stent strategy for treatment of true coronary bifurcation lesions involving a large side branch: the Nordic-Baltic Bifurcation Study IV. <i>Open Heart</i> , 2020 , 7, e000947	3	5
238	Applications of computational simulation in bifurcation stenting: past, present and future. <i>EuroIntervention</i> , 2020 , 16, e698-e700	3.1	
237	Stepwise visualisation of a provisional bifurcation stenting procedure - multimodal visualisation within a reanimated human heart utilising Visible Heart methodologies. <i>EuroIntervention</i> , 2020 , 16, e734-e737	3.1	3
236	Asia Pacific consensus document on coronary bifurcation interventions. <i>EuroIntervention</i> , 2020 , 16, e706-e714	3.1	5

235	Percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: updated 5-year outcomes from the randomised, non-inferiority NOBLE trial. <i>Lancet, The</i> , 2020 , 395, 191-199	40	136
234	Ten-Year Outcomes of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stents in Patients With Versus Without Diabetes Mellitus (SORT OUT III). <i>American Journal of Cardiology</i> , 2020 , 125, 349-353	3	5
233	Efficacy and Safety of Glycoprotein IIb/IIIa Inhibitors on Top of Ticagrelor in STEMI: A Subanalysis of the ATLANTIC Trial. <i>Thrombosis and Haemostasis</i> , 2020 , 120, 65-74	7	7
232	Clinical outcomes three-year after revascularization with biodegradable polymer stents: ultrathin-strut sirolimus-eluting stent versus biolimus-eluting stent: from the Scandinavian organization for randomized trials with clinical outcome VII trial. <i>Coronary Artery Disease</i> , 2020 , 31, 485-492	1.4	6
231	The association between self-reported health status and adverse events: a comparison among coronary artery bypass grafting (CABG) versus percutaneous coronary intervention (PCI). <i>Quality of Life Research</i> , 2020 , 29, 3017-3029	3.7	1
230	Sub-acute cardiac magnetic resonance to predict irreversible reduction in left ventricular ejection fraction after ST-segment elevation myocardial infarction: A DANAMI-3 sub-study. <i>International Journal of Cardiology</i> , 2020 , 301, 215-219	3.2	1
229	Interaction of ischaemic postconditioning and thrombectomy in patients with ST-elevation myocardial infarction. <i>Heart</i> , 2020 , 106, 24-32	5.1	7
228	Effect of remote ischaemic conditioning on clinical outcomes in patients with acute myocardial infarction (CONDI-2/ERIC-PPCI): a single-blind randomised controlled trial. <i>Lancet, The</i> , 2019 , 394, 1415-1424	40	125
227	Unreported exclusion and sampling bias in interpretation of randomized controlled trials in patients with STEMI. <i>International Journal of Cardiology</i> , 2019 , 289, 1-5	3.2	3
226	Comparison of the polymer-free biolimus-coated BioFreedom stent with the thin-strut biodegradable polymer sirolimus-eluting Orsiro stent in an all-comers population treated with percutaneous coronary intervention: Rationale and design of the randomized SORT OUT IX trial. <i>European Heart Journal</i> , 2019 , 40, 1-7	4.9	5
225	Mechanical performance and healing patterns of the novel sirolimus-eluting bioresorbable Fantom scaffold: 6-month and 9-month follow-up by optical coherence tomography in the FANTOM II study. <i>Open Heart</i> , 2019 , 6, e000941	3	3
224	Everolimus-Eluting Versus Biolimus-Eluting Stents With Biodegradable Polymers in Unselected Patients Undergoing Percutaneous Coronary Intervention: A Randomized Noninferiority Trial With 1-Year Follow-Up (SORT OUT VIII Trial). <i>JACC: Cardiovascular Interventions</i> , 2019 , 12, e014102	5	18
223	Diabetes and risk of peripheral artery disease in patients undergoing first-time coronary angiography between 2000 and 2012 - a nationwide study. <i>BMC Cardiovascular Disorders</i> , 2019 , 19, 234	2.3	12
222	Joint consensus on the use of OCT in coronary bifurcation lesions by the European and Japanese bifurcation clubs. <i>EuroIntervention</i> , 2019 , 14, e1568-e1577	3.1	32
221	Percutaneous coronary intervention for obstructive bifurcation lesions: the 14th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2019 , 15, 90-98	3.1	62
220	Ten-year outcomes from a randomised comparison of zotarolimus-eluting and sirolimus-eluting stents: the SORT OUT III study. <i>EuroIntervention</i> , 2019 , 15, e1022-e1024	3.1	4
219	Morphine and Ticagrelor Interaction in Primary Percutaneous Coronary Intervention in ST-Segment Elevation Myocardial Infarction: ATLANTIC-Morphine. <i>American Journal of Cardiovascular Drugs</i> , 2019 , 19, 173-183	4	15
218	A nationwide investigation of CPR courses, books, and skill retention. <i>Resuscitation</i> , 2019 , 134, 110-121	4	6

217	Pre-hospital administration of ticagrelor in diabetic patients with ST-elevation myocardial infarction undergoing primary angioplasty: A sub-analysis of the ATLANTIC trial. <i>Catheterization and Cardiovascular Interventions</i> , 2019 , 93, E369-E377	2.7	3
216	Integrated prediction of lesion-specific ischaemia from quantitative coronary CT angiography using machine learning: a multicentre study. <i>European Radiology</i> , 2018 , 28, 2655-2664	8	85
215	Danegaptide for primary percutaneous coronary intervention in acute myocardial infarction patients: a phase 2 randomised clinical trial. <i>Heart</i> , 2018 , 104, 1593-1599	5.1	11
214	Evaluation of Coronary Artery Stenosis by Quantitative Flow Ratio During Invasive Coronary Angiography: The WIFI II Study (Wire-Free Functional Imaging II). <i>Circulation: Cardiovascular Imaging</i> , 2018 , 11, e007107	3.9	92
213	Randomized comparison of sirolimus eluting, and biolimus eluting bioresorbable polymer stents: the SORT-OUT VII optical coherence tomography study. <i>European Heart Journal Cardiovascular Imaging</i> , 2018 , 19, 329-338	4.1	3
212	Thrombus aspiration and prehospital ticagrelor administration in ST-elevation myocardial infarction: Findings from the ATLANTIC trial. <i>American Heart Journal</i> , 2018 , 196, 1-8	4.9	4
211	Neurological prognostication tools in out-of-hospital cardiac arrest patients in Danish intensive care units from 2005 to 2013. <i>Acta Anaesthesiologica Scandinavica</i> , 2018 , 62, 1412-1420	1.9	1
210	Comparison between patients included in randomized controlled trials of ischemic heart disease and real-world data. A nationwide study. <i>American Heart Journal</i> , 2018 , 204, 128-138	4.9	7
209	Percutaneous coronary intervention for the left main stem and other bifurcation lesions: 12th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2018 , 13, 1540-1553	3.1	127
208	Two-year outcome after biodegradable polymer sirolimus- and biolimus-eluting coronary stents (from the randomised SORT OUT VII trial). <i>EuroIntervention</i> , 2018 , 13, 1587-1590	3.1	13
207	Impact of age on the effect of pre-hospital P2Y12 receptor inhibition in primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: the ATLANTIC-Elderly analysis. <i>EuroIntervention</i> , 2018 , 14, 789-797	3.1	2
206	Intravascular ultrasound in the evaluation and treatment of left main coronary artery disease: a consensus statement from the European Bifurcation Club. <i>EuroIntervention</i> , 2018 , 14, e467-e474	3.1	45
205	Percutaneous coronary intervention in left main coronary artery disease: the 13th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2018 , 14, 112-120	3.1	61
204	Ten-year clinical outcome of patients treated with a drug-eluting stent in the proximal left anterior descending artery segment compared with patients stented in other non-left main coronary segments. <i>EuroIntervention</i> , 2018 , 14, 764-771	3.1	5
203	Intravascular ultrasound findings of the Fantom sirolimus-eluting bioresorbable scaffold at six- and nine-month follow-up: the FANTOM II study. <i>EuroIntervention</i> , 2018 , 14, e1215-e1223	3.1	4
202	Duration of ischemia and treatment effects of pre- versus in-hospital ticagrelor in patients with ST-segment elevation myocardial infarction: Insights from the ATLANTIC study. <i>American Heart Journal</i> , 2018 , 196, 56-64	4.9	6
201	Use of renal replacement therapy after out-of-hospital cardiac arrest in Denmark 2005-2013. <i>Scandinavian Cardiovascular Journal</i> , 2018 , 52, 238-243	2	3
200	Association between socioeconomic factors and ICD implantation in a publicly financed health care system: a Danish nationwide study. <i>Europace</i> , 2018 , 20, 1129-1137	3.9	9

199	Invasive angiography and revascularization in patients with stable angina following prior coronary artery bypass grafting: Results from the East Denmark heart registry. <i>Catheterization and Cardiovascular Interventions</i> , 2017 , 89, 341-349	2.7	1
198	Implantable cardioverter defibrillator and survival after out-of-hospital cardiac arrest due to acute myocardial infarction in Denmark in the years 2001-2012, a nationwide study. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2017 , 6, 144-154	4.3	6
197	10-Year Clinical Outcome After Randomization to Treatment by Sirolimus- or Paclitaxel-Eluting Coronary Stents. <i>Journal of the American College of Cardiology</i> , 2017 , 69, 616-624	15.1	38
196	Pre-hospital ticagrelor in ST-segment elevation myocardial infarction in the French ATLANTIC population. <i>International Journal of Cardiology</i> , 2017 , 244, 49-53	3.2	2
195	Distance to invasive heart centre, performance of acute coronary angiography, and angioplasty and associated outcome in out-of-hospital cardiac arrest: a nationwide study. <i>European Heart Journal</i> , 2017 , 38, 1645-1652	9.5	62
194	The potential of optimizing prehospital triage of patients with suspected acute myocardial infarction using high-sensitivity cardiac troponin T and copeptin. <i>Biomarkers</i> , 2017 , 22, 351-360	2.6	22
193	Myocardial Perfusion Imaging Versus Computed Tomography Angiography-Derived Fractional Flow Reserve Testing in Stable Patients With Intermediate-Range Coronary Lesions: Influence on Downstream Diagnostic Workflows and Invasive Angiography Findings. <i>Journal of the American Heart Association</i> , 2017 , 6	6	18
192	Randomized clinical trial: the impact of gastrointestinal risk factor screening and prophylactic proton pump inhibitor therapy in patients receiving dual antiplatelet therapy. <i>European Journal of Gastroenterology and Hepatology</i> , 2017 , 29, 1118-1125	2.2	13
191	FFR Derived From Coronary CT Angiography in Nonculprit Lesions of Patients With Recent STEMI. <i>JACC: Cardiovascular Imaging</i> , 2017 , 10, 424-433	8.4	44
190	The Extent of Myocardial Injury During Prolonged Targeted Temperature Management After Out-of-Hospital Cardiac Arrest. <i>American Journal of Medicine</i> , 2017 , 130, 37-46	2.4	3
189	Denmark: coronary and structural heart interventions from 2010 to 2015. <i>EuroIntervention</i> , 2017 , 13, Z17-Z20	3.1	3
188	Quantitative angiography methods for bifurcation lesions: a consensus statement update from the European Bifurcation Club. <i>EuroIntervention</i> , 2017 , 13, 115-123	3.1	26
187	Impact of presentation and transfer delays on complete ST-segment resolution before primary percutaneous coronary intervention: insights from the ATLANTIC trial. <i>EuroIntervention</i> , 2017 , 13, 69-77	3.1	13
186	Final five-year outcomes after implantation of biodegradable polymer-coated biolimus-eluting stents versus durable polymer-coated sirolimus-eluting stents. <i>EuroIntervention</i> , 2017 , 13, 1336-1344	3.1	10
185	Treatment of Bifurcation Lesions by Bail-Out TAP or Culotte: Lost in Translation?. <i>Reviews on Recent Clinical Trials</i> , 2017 , 12, 212-215	1.2	
184	The EBC TWO Study (European Bifurcation Coronary TWO): A Randomized Comparison of Provisional T-Stenting Versus a Systematic 2 Stent Culotte Strategy in Large Caliber True Bifurcations. <i>Circulation: Cardiovascular Interventions</i> , 2016 , 9,	6	57
183	Randomized Comparison of a Biodegradable Polymer Ultrathin Strut Sirolimus-Eluting Stent With a Biodegradable Polymer Biolimus-Eluting Stent in Patients Treated With Percutaneous Coronary Intervention: The SORT OUT VII Trial. <i>Circulation: Cardiovascular Interventions</i> , 2016 , 9,	6	78
182	Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. <i>Lancet, The</i> , 2016 , 388, 2743-2752	40	446

181	P2Y12 receptor inhibition and effect of morphine in patients undergoing primary PCI for ST-segment elevation myocardial infarction. The PRIVATE-ATLANTIC study. <i>Thrombosis and Haemostasis</i> , 2016 , 116, 369-78	7	83
180	Influence of preinfarction angina and coronary collateral blood flow on the efficacy of remote ischaemic conditioning in patients with ST segment elevation myocardial infarction: post hoc subgroup analysis of a randomised controlled trial. <i>BMJ Open</i> , 2016 , 6, e013314	3	12
179	Comparison of Outcome of Patients With ST-Segment Elevation Myocardial Infarction and Complete Versus Incomplete ST-Resolution Before Primary Percutaneous Coronary Intervention. <i>American Journal of Cardiology</i> , 2016 , 117, 1735-40	3	9
178	Safety and Efficacy of Everolimus- Versus Sirolimus-Eluting Stents: 5-Year Results From SORT OUT IV. <i>Journal of the American College of Cardiology</i> , 2016 , 67, 751-62	15.1	77
177	Effect of Pre-Hospital Ticagrelor During the First 24 h After Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction: The ATLANTIC-HII Analysis. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 646-56	5	25
176	Percutaneous coronary intervention for coronary bifurcation disease: 11th consensus document from the European Bifurcation Club. <i>EuroIntervention</i> , 2016 , 12, 38-46	3.1	140
175	Assessing the cardiology community position on transradial intervention and the use of bivalirudin in patients with acute coronary syndrome undergoing invasive management: results of an EAPCI survey. <i>EuroIntervention</i> , 2016 , 12, 1154-1163	3.1	10
174	The Danish Heart Registry. <i>Clinical Epidemiology</i> , 2016 , 8, 503-508	5.9	23
173	Renal denervation in treatment-resistant essential hypertension. A randomized, SHAM-controlled, double-blinded 24-h blood pressure-based trial. <i>Journal of Hypertension</i> , 2016 , 34, 1639-47	1.9	68
172	TCT-10 Physiological testing of coronary artery stenosis by computation of invasive coronary angiography. The wire-free functional imaging (WIFI-II) study. <i>Journal of the American College of Cardiology</i> , 2016 , 68, B4-B5	15.1	2
171	Coronary plaque quantification and fractional flow reserve by coronary computed tomography angiography identify ischaemia-causing lesions. <i>European Heart Journal</i> , 2016 , 37, 1220-7	9.5	184
170	Coronary bifurcation lesions treated with simple or complex stenting: 5-year survival from patient-level pooled analysis of the Nordic Bifurcation Study and the British Bifurcation Coronary Study. <i>European Heart Journal</i> , 2016 , 37, 1923-8	9.5	67
169	Zotarolimus-eluting durable-polymer-coated stent versus a biolimus-eluting biodegradable-polymer-coated stent in unselected patients undergoing percutaneous coronary intervention (SORT OUT VI): a randomised non-inferiority trial. <i>Lancet, The</i> , 2015 , 385, 1527-35	40	92
168	Co-registration of optical coherence tomography and X-ray angiography in percutaneous coronary intervention. the Does Optical Coherence Tomography Optimize Revascularization (DOCTOR) fusion study. <i>International Journal of Cardiology</i> , 2015 , 182, 272-8	3.2	30
167	Safety of therapeutic hypothermia combined with primary percutaneous coronary intervention after out-of-hospital cardiac arrest. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015 , 4, 60-3	4.3	8
166	Impact of Side Branch Modeling on Computation of Endothelial Shear Stress in Coronary Artery Disease: Coronary Tree Reconstruction by Fusion of 3D Angiography and OCT. <i>Journal of the American College of Cardiology</i> , 2015 , 66, 125-35	15.1	54
165	Influence of multivessel disease with or without additional revascularization on mortality in patients with ST-segment elevation myocardial infarction. <i>American Heart Journal</i> , 2015 , 170, 70-8	4.9	16
164	One-year clinical and angiographic results of hybrid coronary revascularization. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2015 , 150, 1181-6	1.5	13

163	Influence of Coronary Calcification on the Diagnostic Performance of CT Angiography Derived FFR in Coronary Artery Disease: A Substudy of the NXT Trial. <i>JACC: Cardiovascular Imaging</i> , 2015 , 8, 1045-1055	8.4	102
162	Support with intra-aortic balloon pump vs. Impella2.5 and blood flow to the heart, brain and kidneys - an experimental porcine model of ischaemic heart failure. <i>International Journal of Cardiology</i> , 2015 , 178, 153-8	3.2	15
161	The impact of distal embolization and distal protection on long-term outcome in patients with ST elevation myocardial infarction randomized to primary percutaneous coronary intervention--results from a randomized study. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015 , 4, 180-8	4.3	14
160	Quality of cardiopulmonary resuscitation in out-of-hospital cardiac arrest before and after introduction of a mechanical chest compression device, LUCAS-2; a prospective, observational study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2015 , 23, 37	3.6	26
159	Impact of cardiovascular risk factors and medication use on the efficacy of remote ischaemic conditioning: post hoc subgroup analysis of a randomised controlled trial. <i>BMJ Open</i> , 2015 , 5, e006923	3	45
158	Randomized comparison of a sirolimus-eluting Orsiro stent with a biolimus-eluting Nobori stent in patients treated with percutaneous coronary intervention: Rationale and study design of the Scandinavian Organization for Randomized Trials with Clinical Outcome VII trial. <i>American Heart Journal</i> , 2015 , 170, 210-5	4.9	16
157	A "normal" invasive coronary angiogram may not be normal. <i>Journal of Cardiovascular Computed Tomography</i> , 2015 , 9, 264-6	2.8	4
156	Long-term outcome of sirolimus-eluting and zotarolimus-eluting coronary stent implantation in patients with and without diabetes mellitus (a Danish organization for randomized trials on clinical outcome III substudy). <i>American Journal of Cardiology</i> , 2015 , 115, 298-302	3	12
155	Feasibility and early safety of hybrid coronary revascularisation combining off-pump coronary surgery through J-hemisternotomy with percutaneous coronary intervention. <i>EuroIntervention</i> , 2015 , 10, e1-6	3.1	8
154	The EuroIntervention coronary bifurcation treatment supplement. <i>EuroIntervention</i> , 2015 , 11 Suppl V, V9-11	3.1	3
153	When is a two-stent technique needed and which technique should then be used for bifurcation coronary lesions?. <i>EuroIntervention</i> , 2015 , 11 Suppl V, V96-8	3.1	2
152	Technical aspects of the culotte technique. <i>EuroIntervention</i> , 2015 , 11 Suppl V, V99-101	3.1	5
151	Differences between the left main and other bifurcations. <i>EuroIntervention</i> , 2015 , 11 Suppl V, V106-10	3.1	25
150	When and how to use BRS in bifurcations?. <i>EuroIntervention</i> , 2015 , 11 Suppl V, V185-7	3.1	4
149	Diagnostic performance of noninvasive fractional flow reserve derived from coronary computed tomography angiography in suspected coronary artery disease: the NXT trial (Analysis of Coronary Blood Flow Using CT Angiography: Next Steps). <i>Journal of the American College of Cardiology</i> , 2014 , 63, 1145-1155	15.1	871
148	Prevention of contrast-induced nephropathy with N-acetylcysteine or sodium bicarbonate in patients with ST-segment-myocardial infarction: a prospective, randomized, open-labeled trial. <i>Circulation: Cardiovascular Interventions</i> , 2014 , 7, 216-24	6	39
147	Antiplatelet therapy for stable coronary artery disease in atrial fibrillation patients taking an oral anticoagulant: a nationwide cohort study. <i>Circulation</i> , 2014 , 129, 1577-85	16.7	204
146	Intimal hyperplasia and vascular remodeling after everolimus-eluting and sirolimus-eluting stent implantation in diabetic patients: the randomized Diabetes and Drug-Eluting Stent (DiabeDES) IV Intravascular Ultrasound trial. <i>Catheterization and Cardiovascular Interventions</i> , 2014 , 83, 864-72	2.7	8

145	Fractional flow reserve derived from coronary CT angiography: variation of repeated analyses. <i>Journal of Cardiovascular Computed Tomography</i> , 2014 , 8, 307-14	2.8	39
144	Prehospital ticagrelor in ST-segment elevation myocardial infarction. <i>New England Journal of Medicine</i> , 2014 , 371, 1016-27	59.2	409
143	Clopidogrel discontinuation within the first year after coronary drug-eluting stent implantation: an observational study. <i>BMC Cardiovascular Disorders</i> , 2014 , 14, 100	2.3	22
142	Three-year outcomes after revascularization with everolimus- and sirolimus-eluting stents from the SORT OUT IV trial. <i>JACC: Cardiovascular Interventions</i> , 2014 , 7, 840-8	5	27
141	Long-term outcome following percutaneous coronary intervention with drug-eluting stents compared with bare-metal stents in saphenous vein graft lesions: from Western Denmark Heart Registry. <i>Catheterization and Cardiovascular Interventions</i> , 2014 , 83, 1035-42	2.7	9
140	Similar five-year outcome with paclitaxel- and sirolimus-eluting coronary stents. <i>Scandinavian Cardiovascular Journal</i> , 2014 , 48, 148-55	2	8
139	Impact of health care system delay in patients with ST-elevation myocardial infarction on return to labor market and work retirement. <i>American Journal of Cardiology</i> , 2014 , 114, 1810-6	3	19
138	Pain and discomfort in closure of femoral access coronary angiography. The CLOSuredEVICES Used in everyday Practice (CLOSE-UP) pain sub study. <i>European Journal of Cardiovascular Nursing</i> , 2014 , 13, 221-6	3.3	11
137	Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. <i>Lancet, The</i> , 2014 , 383, 2047-2056	40	80
136	Invasive coronary treatment strategies for out-of-hospital cardiac arrest: a consensus statement from the European association for percutaneous cardiovascular interventions (EAPCI)/stent for life (SFL) groups. <i>EuroIntervention</i> , 2014 , 10, 31-7	3.1	164
135	Randomised comparison of manual compression and FemoSeal [®] vascular closure device for closure after femoral artery access coronary angiography: the CLOSure dEVICES Used in everyday Practice (CLOSE-UP) study. <i>EuroIntervention</i> , 2014 , 10, 183-90	3.1	40
134	Outcomes after revascularisation with everolimus- and sirolimus-eluting stents in patients with acute coronary syndromes and stable angina pectoris: a substudy of the SORT OUT IV trial. <i>EuroIntervention</i> , 2014 , 10, 212-23	3.1	8
133	Percutaneous coronary intervention for coronary bifurcation disease: consensus from the first 10 years of the European Bifurcation Club meetings. <i>EuroIntervention</i> , 2014 , 10, 545-60	3.1	177
132	Long-term outcome after drug-eluting versus bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: 5 years follow-up from the randomized DEDICATION trial (Drug Elution and Distal Protection in Acute Myocardial Infarction). <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 516-23	5	32
131	Long-term results after simple versus complex stenting of coronary artery bifurcation lesions: Nordic Bifurcation Study 5-year follow-up results. <i>Journal of the American College of Cardiology</i> , 2013 , 62, 30-4	15.1	127
130	Timely and optimal treatment of patients with STEMI. <i>Nature Reviews Cardiology</i> , 2013 , 10, 41-8	14.8	45
129	Rationale and design of the HeartFlowNXT (HeartFlow analysis of coronary blood flow using CT angiography: NeXt sTEps) study. <i>Journal of Cardiovascular Computed Tomography</i> , 2013 , 7, 279-88	2.8	47
128	Acquired peri-stent evaginations in a second generation durable polymer drug eluting stent. <i>Cardiovascular Revascularization Medicine</i> , 2013 , 14, 246-7	1.6	2

127	Clinical outcome after crush versus culotte stenting of coronary artery bifurcation lesions: the Nordic Stent Technique Study 36-month follow-up results. <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 1160-5	5	42
126	Quantitative point-of-care troponin T measurement for diagnosis and prognosis in patients with a suspected acute myocardial infarction. <i>American Journal of Cardiology</i> , 2013 , 112, 1361-6	3	37
125	Biolimus-eluting biodegradable polymer-coated stent versus durable polymer-coated sirolimus-eluting stent in unselected patients receiving percutaneous coronary intervention (SORT OUT V): a randomised non-inferiority trial. <i>Lancet, The</i> , 2013 , 381, 661-9	40	152
124	Intravascular ultrasound assessed incomplete stent apposition and stent fracture in stent thrombosis after bare metal versus drug-eluting stent treatment the Nordic Intravascular Ultrasound Study (NIVUS). <i>International Journal of Cardiology</i> , 2013 , 168, 1010-6	3.2	24
123	Ambulance or in-catheterization laboratory administration of ticagrelor for primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: rationale and design of the randomized, double-blind Administration of Ticagrelor in the cath Lab or in the Ambulance for New ST elevation myocardial infarction to open the Coronary artery (ATLANTIC) study. <i>American Heart</i> 2012 ESC STEMI guidelines and reperfusion therapy: Evidence base ignored, threatening optimal patient management. Heart , 2013 , 99, 1154-6	4.9	39
122	Coronary edema demonstrated by cardiovascular magnetic resonance in patients with peri-stent inflammation and aneurysm formation after treatment by drug-eluting stents. <i>Circulation: Cardiovascular Imaging</i> , 2013 , 6, 352-4	5.1	14
121	A response to a misrepresentation of the STEMI guidelines: the response. <i>Heart</i> , 2013 , 99, 1787-8	3.9	5
120	Three-dimensional multidetector computed tomography versus conventional 2-dimensional transesophageal echocardiography for annular sizing in transcatheter aortic valve replacement: Influence on postprocedural paravalvular aortic regurgitation. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 82, 977-86	5.1	3
119	Moderate overweight is beneficial and severe obesity detrimental for patients with documented atherosclerotic heart disease. <i>Heart</i> , 2013 , 99, 655-60	2.7	34
118	Outcomes after primary percutaneous coronary intervention in octogenarians and nonagenarians with ST-segment elevation myocardial infarction: from the Western Denmark heart registry. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 81, 912-9	5.1	51
117	Incidence of definite stent thrombosis or in-stent restenosis after drug-eluting stent implantation for treatment of coronary in-stent restenosis: from Western Denmark Heart Registry. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 81, 260-5	2.7	48
116	Event detection using population-based health care databases in randomized clinical trials: a novel research tool in interventional cardiology. <i>Clinical Epidemiology</i> , 2013 , 5, 357-61	2.7	11
115	Consensus from the 7th European Bifurcation Club meeting. <i>EuroIntervention</i> , 2013 , 9, 36-45	5.9	21
114	Zotarolimus-eluting vs. sirolimus-eluting coronary stents in patients with and without acute coronary syndromes: a SORT OUT III substudy. <i>European Journal of Clinical Investigation</i> , 2012 , 42, 1047-54	3.1	51
113	Clinical outcomes after treatment of multiple lesions with zotarolimus-eluting versus sirolimus-eluting coronary stents (a SORT OUT III substudy). <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 18	4.6	9
112	Comparison of zotarolimus-eluting and sirolimus-eluting coronary stents: a study from the Western Denmark Heart Registry. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 84	2.3	0
111	Earlier reperfusion in patients with ST-elevation myocardial infarction by use of helicopter. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2012 , 20, 70	2.3	2
110		3.6	18

109	2-year patient-related versus stent-related outcomes: the SORT OUT IV (Scandinavian Organization for Randomized Trials With Clinical Outcome IV) Trial. <i>Journal of the American College of Cardiology</i> , 2012 , 60, 1140-7	15.1	41
108	3-Year clinical outcomes in the randomized SORT OUT III superiority trial comparing zotarolimus- and sirolimus-eluting coronary stents. <i>JACC: Cardiovascular Interventions</i> , 2012 , 5, 812-8	5	42
107	Comparison of outcomes in patients with versus without diabetes mellitus after revascularization with everolimus- and sirolimus-eluting stents (from the SORT OUT IV trial). <i>American Journal of Cardiology</i> , 2012 , 110, 1585-91	3	44
106	Influence of diabetes mellitus on clinical outcomes following primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. <i>American Journal of Cardiology</i> , 2012 , 109, 629-35	3	43
105	Dimensions of socioeconomic status and clinical outcome after primary percutaneous coronary intervention. <i>Circulation: Cardiovascular Interventions</i> , 2012 , 5, 641-8	6	30
104	Randomized comparison of everolimus-eluting and sirolimus-eluting stents in patients treated with percutaneous coronary intervention: the Scandinavian Organization for Randomized Trials with Clinical Outcome IV (SORT OUT IV). <i>Circulation</i> , 2012 , 125, 1246-55	16.7	131
103	Side branch fractional flow reserve measurements after main vessel stenting: a Nordic-Baltic Bifurcation Study III substudy. <i>EuroIntervention</i> , 2012 , 7, 1155-61	3.1	41
102	Culprit only or multivessel percutaneous coronary interventions in patients with ST-segment elevation myocardial infarction and multivessel disease. <i>EuroIntervention</i> , 2012 , 8, 456-64	3.1	32
101	The risk and prognostic impact of definite stent thrombosis or in-stent restenosis after coronary stent implantation. <i>EuroIntervention</i> , 2012 , 8, 591-8	3.1	17
100	Sex- and age-related differences in clinical outcome after primary percutaneous coronary intervention. <i>EuroIntervention</i> , 2012 , 8, 904-11	3.1	27
99	Simple or complex stenting for bifurcation coronary lesions: a patient-level pooled-analysis of the Nordic Bifurcation Study and the British Bifurcation Coronary Study. <i>Circulation: Cardiovascular Interventions</i> , 2011 , 4, 57-64	6	118
98	Left ventricular contractile function after distal protection in primary percutaneous coronary intervention: results from the Drug Elution and Distal Protection in ST-Elevation Myocardial Infarction trial. <i>International Journal of Cardiology</i> , 2011 , 146, 395-8	3.2	4
97	Long-term safety and efficacy of drug-eluting stents. <i>Lancet, The</i> , 2011 , 377, 1213-4	40	2
96	Health care system delay and heart failure in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention: follow-up of population-based medical registry data. <i>Annals of Internal Medicine</i> , 2011 , 155, 361-7	8	69
95	Nonsteroidal antiinflammatory drug use and cardiovascular risks after coronary stent implantation. <i>Pharmacotherapy</i> , 2011 , 31, 458-68	5.8	8
94	Prehospital troponin T testing in the diagnosis and triage of patients with suspected acute myocardial infarction. <i>American Journal of Cardiology</i> , 2011 , 107, 1436-40	3	46
93	Outcome of sirolimus-eluting versus zotarolimus-eluting coronary stent implantation in patients with and without diabetes mellitus (a SORT OUT III Substudy). <i>American Journal of Cardiology</i> , 2011 , 108, 1232-7	3	35
92	Comparison of outcomes of patients \geq 60 years of age having percutaneous coronary intervention according to presentation (stable vs unstable angina pectoris/non-ST-segment elevation myocardial infarction vs ST-segment elevation myocardial infarction). <i>American Journal of Cardiology</i> , 2011 , 108, 1385-100	3	14

91	Quantitative coronary analysis in the Nordic Bifurcation studies. <i>International Journal of Cardiovascular Imaging</i> , 2011 , 27, 175-80	2.5	14
90	Quality of cardiopulmonary resuscitation in out-of-hospital cardiac arrest is hampered by interruptions in chest compressions—a nationwide prospective feasibility study. <i>Resuscitation</i> , 2011 , 82, 263-9	4	51
89	Randomized comparison of final kissing balloon dilatation versus no final kissing balloon dilatation in patients with coronary bifurcation lesions treated with main vessel stenting: the Nordic-Baltic Bifurcation Study III. <i>Circulation</i> , 2011 , 123, 79-86	16.7	219
88	Primary percutaneous coronary intervention as a national reperfusion strategy in patients with ST-segment elevation myocardial infarction. <i>Circulation: Cardiovascular Interventions</i> , 2011 , 4, 570-6	6	31
87	Urban and rural implementation of pre-hospital diagnosis and direct referral for primary percutaneous coronary intervention in patients with acute ST-elevation myocardial infarction. <i>European Heart Journal</i> , 2011 , 32, 430-6	9.5	132
86	Late lumen loss and intima hyperplasia after sirolimus-eluting and zotarolimus-eluting stent implantation in diabetic patients: the diabetes and drug-eluting stent (DiabeDES III) angiography and intravascular ultrasound trial. <i>EuroIntervention</i> , 2011 , 7, 323-31	3.1	19
85	Use of three-dimensional optical coherence tomography to verify correct wire position in a jailed side branch after main vessel stent implantation. <i>EuroIntervention</i> , 2011 , 7, 528-9	3.1	9
84	Rationale and design of a randomized clinical comparison of everolimus-eluting (Xience v/Promus) and sirolimus-eluting (cypher select+) coronary stents in unselected patients with coronary heart disease. <i>Cardiology</i> , 2010 , 116, 73-8	1.6	17
83	System delay and mortality among patients with STEMI treated with primary percutaneous coronary intervention. <i>JAMA - Journal of the American Medical Association</i> , 2010 , 304, 763-71	27.4	431
82	Increased rate of stent thrombosis and target lesion revascularization after filter protection in primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: 15-month follow-up of the DEDICATION (Drug Elution and Distal Protection in ST Elevation Myocardial Infarction) Trial. <i>Journal of the American College of Cardiology</i> , 2010 , 55, 267-71	15.1	30
81	Long-term outcome after drug-eluting versus bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: 3-year follow-up of the randomized DEDICATION (Drug Elution and Distal Protection in Acute Myocardial Infarction) Trial. <i>Journal of the American College of Cardiology</i> , 2010 , 56, 641-5	15.1	63
80	Sirolimus-eluting versus bare-metal stent implantation in patients with ostial lesions. <i>International Journal of Cardiology</i> , 2010 , 145, 162-3	3.2	1
79	Remote ischaemic conditioning before hospital admission, as a complement to angioplasty, and effect on myocardial salvage in patients with acute myocardial infarction: a randomised trial. <i>Lancet, The</i> , 2010 , 375, 727-34	40	773
78	Efficacy and safety of zotarolimus-eluting and sirolimus-eluting coronary stents in routine clinical care (SORT OUT III): a randomised controlled superiority trial. <i>Lancet, The</i> , 2010 , 375, 1090-9	40	174
77	Outcome in high risk patients with unprotected left main coronary artery stenosis treated with percutaneous coronary intervention. <i>Catheterization and Cardiovascular Interventions</i> , 2010 , 75, 101-8	2.7	18
76	Zotarolimus-eluting versus sirolimus-eluting coronary stent implantation. <i>Interventional Cardiology</i> , 2010 , 2, 807-812	3	1
75	Long-term outcomes after percutaneous coronary intervention in patients with and without diabetes mellitus in Western Denmark. <i>American Journal of Cardiology</i> , 2010 , 105, 1513-9	3	35
74	Cardiac arrest in the catheterisation laboratory: a 5-year experience of using mechanical chest compressions to facilitate PCI during prolonged resuscitation efforts. <i>Resuscitation</i> , 2010 , 81, 383-7	4	121

73	Paclitaxel and sirolimus eluting stents versus bare metal stents: long-term risk of stent thrombosis and other outcomes. From the Western Denmark Heart Registry. <i>EuroIntervention</i> , 2010 , 5, 898-905	3.1	39
72	Consensus from the 5th European Bifurcation Club meeting. <i>EuroIntervention</i> , 2010 , 6, 34-38	3.1	117
71	One or two stents for coronary bifurcation lesions?. <i>EuroIntervention</i> , 2010 , 6, J61-J64	3.1	4
70	Infarct size and myocardial salvage after primary angioplasty in patients presenting with symptoms for . <i>European Heart Journal</i> , 2009 , 30, 1322-30	9.5	65
69	Randomized comparison of coronary bifurcation stenting with the crush versus the culotte technique using sirolimus eluting stents: the Nordic stent technique study. <i>Circulation: Cardiovascular Interventions</i> , 2009 , 2, 27-34	6	133
68	ST changes before and during primary percutaneous coronary intervention predict final infarct size in patients with ST elevation myocardial infarction. <i>Journal of Electrocardiology</i> , 2009 , 42, 64-72	1.4	22
67	Temporal course of pregnancy-associated plasma protein-A in angioplasty-treated ST-elevation myocardial infarction patients and potential significance of concomitant heparin administration. <i>American Journal of Cardiology</i> , 2009 , 103, 29-35	3	40
66	Comparison of the sirolimus-eluting versus paclitaxel-eluting coronary stent in patients with diabetes mellitus: the diabetes and drug-eluting stent (DiabeDES) randomized angiography trial. <i>American Journal of Cardiology</i> , 2009 , 103, 345-9	3	48
65	Serial intravascular ultrasound analysis of peri-stent remodeling and proximal and distal edge effects after sirolimus-eluting or paclitaxel-eluting stent implantation in patients with diabetes mellitus. <i>American Journal of Cardiology</i> , 2009 , 103, 1083-8	3	16
64	Prevalence and significance of accelerated idioventricular rhythm in patients with ST-elevation myocardial infarction treated with primary percutaneous coronary intervention. <i>American Journal of Cardiology</i> , 2009 , 104, 1641-6	3	39
63	Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 784-91	2.1	10
62	2-year clinical outcomes after implantation of sirolimus-eluting, paclitaxel-eluting, and bare-metal coronary stents: results from the WDHR (Western Denmark Heart Registry). <i>Journal of the American College of Cardiology</i> , 2009 , 53, 658-64	15.1	268
61	Primary PCI as the preferred reperfusion therapy in STEMI: it is a matter of time. <i>Heart</i> , 2009 , 95, 362-9	5.1	51
60	Percutaneous coronary intervention for bifurcation lesions: 2008 consensus document from the fourth meeting of the European Bifurcation Club. <i>EuroIntervention</i> , 2009 , 5, 39-49	3.1	92
59	Intravascular ultrasound assessment of expansion of the sirolimus-eluting (cypher select) and paclitaxel-eluting (Taxus Express-2) stent in patients with diabetes mellitus. <i>American Journal of Cardiology</i> , 2008 , 102, 19-26	3	13
58	Comparison of stent thrombosis, myocardial infarction, and mortality following drug-eluting versus bare-metal stent coronary intervention in patients with diabetes mellitus. <i>American Journal of Cardiology</i> , 2008 , 102, 165-72	3	26
57	Randomized comparison of distal protection versus conventional treatment in primary percutaneous coronary intervention: the drug elution and distal protection in ST-elevation myocardial infarction (DEDICATION) trial. <i>Journal of the American College of Cardiology</i> , 2008 , 51, 899-905	15.1	117
56	Long-term outcome in patients treated with sirolimus-eluting stents in complex coronary artery lesions: 3-year results of the SCANDSTENT (Stenting Coronary Arteries in Non-Stress/Benestent Disease) trial. <i>Journal of the American College of Cardiology</i> , 2008 , 51, 2011-6	15.1	44

55	Neointimal hyperplasia after sirolimus-eluting and paclitaxel-eluting stent implantation in diabetic patients: the Randomized Diabetes and Drug-Eluting Stent (DiabeDES) Intravascular Ultrasound Trial. <i>European Heart Journal</i> , 2008 , 29, 2733-41	9.5	32
54	Two stent or not two stent--that is the question. <i>European Heart Journal</i> , 2008 , 29, 2829-30	9.5	1
53	Comparison of paclitaxel- and sirolimus-eluting stents in everyday clinical practice: the SORT OUT II randomized trial. <i>JAMA - Journal of the American Medical Association</i> , 2008 , 299, 409-16	27.4	116
52	Clinical outcome after primary percutaneous coronary intervention with drug-eluting and bare metal stents in patients with ST-segment elevation myocardial infarction. <i>Circulation: Cardiovascular Interventions</i> , 2008 , 1, 176-84	6	26
51	Drug-eluting versus bare metal stents in patients with st-segment-elevation myocardial infarction: eight-month follow-up in the Drug Elution and Distal Protection in Acute Myocardial Infarction (DEDICATION) trial. <i>Circulation</i> , 2008 , 118, 1155-62	16.7	58
50	Classification of coronary artery bifurcation lesions and treatments: time for a consensus!. <i>Catheterization and Cardiovascular Interventions</i> , 2008 , 71, 175-83	2.7	210
49	Safety in simple versus complex stenting of coronary artery bifurcation lesions. The nordic bifurcation study 14-month follow-up results. <i>EuroIntervention</i> , 2008 , 4, 229-33	3.1	53
48	Does postsystolic motion or shortening predict recovery of myocardial function after primary percutaneous coronary intervention?. <i>Journal of the American Society of Echocardiography</i> , 2007 , 20, 505-11	5.8	19
47	Stent thrombosis, myocardial infarction, and death after drug-eluting and bare-metal stent coronary interventions. <i>Journal of the American College of Cardiology</i> , 2007 , 50, 463-70	15.1	195
46	Influence of a pressure gradient distal to implanted bare-metal stent on in-stent restenosis after percutaneous coronary intervention. <i>Circulation</i> , 2007 , 116, 2802-8	16.7	17
45	Recruitable collateral blood flow index predicts coronary in-stent restenosis after percutaneous coronary intervention. <i>European Heart Journal</i> , 2007 , 28, 1820-6	9.5	14
44	Target lesion revascularisation in patients treated with a sirolimus-eluting or paclitaxel-eluting stent. <i>Heart</i> , 2007 , 93, 694-7	5.1	10
43	Percutaneous coronary intervention of bifurcation lesions: state-of-the-art. Insights from the second meeting of the European Bifurcation Club. <i>EuroIntervention</i> , 2007 , 3, 44-9	3.1	22
42	Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). <i>European Heart Journal</i> , 2006 , 27, 267-75	9.5	57
41	Percutaneous treatment of coronary bifurcation lesions: a novel "extended Y" technique with complete lesion stent coverage. <i>Heart</i> , 2006 , 92, 981-2	5.1	11
40	Randomized study on simple versus complex stenting of coronary artery bifurcation lesions: the Nordic bifurcation study. <i>Circulation</i> , 2006 , 114, 1955-61	16.7	574
39	Routine thrombectomy in percutaneous coronary intervention for acute ST-segment-elevation myocardial infarction: a randomized, controlled trial. <i>Circulation</i> , 2006 , 114, 40-7	16.7	216
38	The Stenting Coronary Arteries in Non-stress/benestent Disease (SCANDSTENT) trial. <i>Journal of the American College of Cardiology</i> , 2006 , 47, 449-55	15.1	95

37	Comparison of sirolimus-eluting and bare metal stents in coronary bifurcation lesions: subgroup analysis of the Stenting Coronary Arteries in Non-Stress/Benestent Disease Trial (SCANDSTENT). <i>American Heart Journal</i> , 2006 , 152, 1140-5	4.9	67
36	Gains and losses of warfarin therapy as performed in an anticoagulation clinic. <i>Journal of Internal Medicine</i> , 2006 , 259, 296-304	10.8	24
35	Outcome of unprotected left main percutaneous coronary intervention in surgical low-risk, surgical high-risk, and acute myocardial infarction patients. <i>EuroIntervention</i> , 2006 , 1, 403-8	3.1	10
34	Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. <i>European Heart Journal</i> , 2005 , 26, 18-26	9.5	213
33	Reduction of treatment delay in patients with ST-elevation myocardial infarction: impact of pre-hospital diagnosis and direct referral to primary percutaneous intervention. <i>European Heart Journal</i> , 2005 , 26, 1343; author reply 1343-4	9.5	2
32	Prehospital evaluation in ST-elevation myocardial infarction patients treated with primary percutaneous coronary intervention. <i>Journal of Electrocardiology</i> , 2005 , 38, 187-92	1.4	34
31	When analyses are invalidated by erroneous assumptions. <i>International Journal of Technology Assessment in Health Care</i> , 2005 , 21, 414-5; author reply 416-20	1.8	
30	Reduction of treatment delay in patients with ST-elevation myocardial infarction: impact of pre-hospital diagnosis and direct referral to primary percutaneous coronary intervention. <i>European Heart Journal</i> , 2005 , 26, 770-7	9.5	195
29	Myocardial infarction centres: the way forward. <i>Heart</i> , 2005 , 91 Suppl 3, iii12-5	5.1	5
28	Should patients with acute ST elevation MI be transferred for primary PCI?. <i>Heart</i> , 2004 , 90, 1358-63	5.1	6
27	Quality of lipid-lowering therapy in patients with ischaemic heart disease: a register-based study in 3477 patients. <i>Journal of Internal Medicine</i> , 2004 , 255, 367-72	10.8	13
26	The platelet polymorphism PLA2 is a genetic risk factor for myocardial infarction. <i>Journal of Internal Medicine</i> , 2004 , 255, 637-44	10.8	40
25	Haemostatic activity in patients with atrial fibrillation treated with low-molecular-weight heparin before and after electrical cardioversion. <i>Journal of Thrombosis and Thrombolysis</i> , 2004 , 17, 185-9	5.1	5
24	Is there a hypercoagulable phase during initiation of antithrombotic therapy with oral anticoagulants in patients with atrial fibrillation?. <i>Thrombosis Research</i> , 2003 , 109, 241-6	8.2	18
23	Are we underestimating the full potential of early thrombolytic treatment in patients with acute myocardial infarction?. <i>British Heart Journal</i> , 2003 , 89, 483-4		8
22	A randomized controlled trial of shared care versus routine care for patients receiving oral anticoagulant therapy. <i>Journal of Internal Medicine</i> , 2002 , 252, 322-31	10.8	18
21	Telemedicine used for remote prehospital diagnosing in patients suspected of acute myocardial infarction. <i>Journal of Internal Medicine</i> , 2002 , 252, 412-20	10.8	73
20	Cholesterol reduction following health screening in general practice. <i>Scandinavian Journal of Primary Health Care</i> , 2002 , 20, 219-23	2.7	11

19	Prospective evaluation of the quality of oral anticoagulation management in an outpatient clinic and in general practices. <i>Thrombosis Research</i> , 2002 , 105, 103-8	8.2	9
18	Electromechanical mapping for detection of myocardial viability in patients with ischemic cardiomyopathy. <i>Circulation</i> , 2001 , 103, 1631-7	16.7	68
17	HbA1c in an unselected population of 4438 people with type 2 diabetes in a Danish county. <i>Scandinavian Journal of Primary Health Care</i> , 2001 , 19, 241-6	2.7	14
16	Intravenous magnesium does not influence the activity of the coagulation cascade. <i>Blood Coagulation and Fibrinolysis</i> , 2001 , 12, 223-8	1	16
15	Evaluation of a simple dosage scheme for transition from phenprocoumon to warfarin in oral anticoagulation. <i>Thrombosis Research</i> , 2000 , 98, 157-63	8.2	9
14	Near-Patient Test for C-Reactive Protein in General Practice: Assessment of Clinical, Organizational, and Economic Outcomes. <i>Clinical Chemistry</i> , 1999 , 45, 478-485	5.5	50
13	From science to everyday clinical practice. Need for systematic evaluation of research findings. <i>Scandinavian Journal of Primary Health Care</i> , 1999 , 17, 6-10	2.7	3
12	The Arg506Gln mutation (FV Leiden) among a cohort of 4188 unselected Danish newborns. <i>Thrombosis Research</i> , 1998 , 89, 211-5	8.2	39
11	Effect of anticoagulant therapy on the hypercoagulable state in patients carrying the factor V Arg506Gln mutation. <i>Thrombosis Research</i> , 1998 , 92, 157-62	8.2	4
10	Markers of coagulation and fibrinolysis as measures of disease activity in inflammatory bowel disease. <i>Scandinavian Journal of Gastroenterology</i> , 1998 , 33, 637-43	2.4	41
9	C-reactive protein in general practice--how commonly is it used and why?. <i>Scandinavian Journal of Primary Health Care</i> , 1997 , 15, 35-8	2.7	10
8	Evaluation of a near-patient test for C-reactive protein used in daily routine in primary healthcare by use of difference plots. <i>Clinical Chemistry</i> , 1997 , 43, 2064-2075	5.5	33
7	Biological variation of International Normalized Ratio for prothrombin times, and consequences in monitoring oral anticoagulant therapy: computer simulation of serial measurements with goal-setting for analytical quality. <i>Clinical Chemistry</i> , 1997 , 43, 2175-2182	5.5	21
6	Graphical interpretation of analytical data from comparison of a field method with a Reference Method by use of difference plots. <i>Clinical Chemistry</i> , 1997 , 43, 2039-2046	5.5	91
5	Research, technology assessment, and quality assurance. <i>European Journal of General Practice</i> , 1996 , 2, 162-165	2.8	5
4	Interpretation of serial measurements of international normalized ratio for prothrombin times in monitoring oral anticoagulant therapy. <i>Clinical Chemistry</i> , 1995 , 41, 1171-1176	5.5	40
3	International normalized ratio for prothrombin times in patients taking oral anticoagulants: critical difference and probability of significant change in consecutive measurements. <i>Clinical Chemistry</i> , 1995 , 41, 444-447	5.5	60
2	Hallucinogenic mushroom use by Danish students: pattern of consumption. <i>Journal of Internal Medicine</i> , 1993 , 233, 111-2	10.8	4

1 The Nordic Experience 21-33