

Richard K Johnson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1427335/richard-k-johnson-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

127 papers	5,672 citations	36 h-index	73 g-index
129 ext. papers	6,228 ext. citations	4.2 avg, IF	5.61 L-index

#	Paper	IF	Citations
127	Species distribution models as a tool for early detection of the invasive <i>Raphidiopsis raciborskii</i> in European lakes.. <i>Harmful Algae</i> , 2022 , 113, 102202	5.3	1
126	Responses of multiple structural and functional indicators along three contrasting disturbance gradients. <i>Ecological Indicators</i> , 2022 , 135, 108514	5.8	0
125	Panarchy and management of lake ecosystems. <i>Ecology and Society</i> , 2021 , 26, 1-7	4.1	2
124	A Bayesian Belief Network learning tool integrates multi-scale effects of riparian buffers on stream invertebrates. <i>Science of the Total Environment</i> , 2021 , 810, 152146	10.2	2
123	A global agenda for advancing freshwater biodiversity research. <i>Ecology Letters</i> , 2021 ,	10	6
122	Contrasting responses of terrestrial and aquatic consumers in riparian stream networks to local and landscape level drivers of environmental change. <i>Basic and Applied Ecology</i> , 2021 ,	3.2	1
121	Forested Riparian Buffers Change the Taxonomic and Functional Composition of Stream Invertebrate Communities in Agricultural Catchments. <i>Water (Switzerland)</i> , 2021 , 13, 1028	3	3
120	Local habitat is a strong determinant of spatial and temporal patterns of macrophyte diversity and composition in boreal lakes. <i>Freshwater Biology</i> , 2021 , 66, 1490-1501	3.1	0
119	Assessing the Ecological Status of European Rivers and Lakes Using Benthic Invertebrate Communities: A Practical Catalogue of Metrics and Methods. <i>Water (Switzerland)</i> , 2021 , 13, 346	3	4
118	Stochastic processes and ecological connectivity drive stream invertebrate community responses to short-term drought. <i>Journal of Animal Ecology</i> , 2021 , 90, 886-898	4.7	4
117	The Structure of Riparian Vegetation in Agricultural Landscapes Influences Spider Communities and Aquatic-Terrestrial Linkages. <i>Water (Switzerland)</i> , 2020 , 12, 2855	3	9
116	Habitat patchiness, ecological connectivity and the uneven recovery of boreal stream ecosystems from an experimental drought. <i>Global Change Biology</i> , 2020 , 26, 3455-3472	11.4	9
115	Assessing the Benefits of Forested Riparian Zones: A Qualitative Index of Riparian Integrity Is Positively Associated with Ecological Status in European Streams. <i>Water (Switzerland)</i> , 2020 , 12, 1178	3	32
114	Phytoplankton size- and abundance-based resilience assessments reveal nutrient rather than water level effects. <i>Science of the Total Environment</i> , 2020 , 746, 141110	10.2	1
113	Partitioning spatial, environmental, and community drivers of ecosystem functioning. <i>Landscape Ecology</i> , 2019 , 34, 2371-2384	4.3	11
112	Policy-driven monitoring and evaluation: Does it support adaptive management of socio-ecological systems?. <i>Science of the Total Environment</i> , 2019 , 662, 373-384	10.2	30
111	Use of taxon-specific models of phytoplankton assemblage composition and biomass for detecting impact. <i>Ecological Indicators</i> , 2019 , 97, 447-456	5.8	3

110	Modelling outperforms typologies for establishing reference conditions of boreal lake and stream invertebrate assemblages. <i>Ecological Indicators</i> , 2018 , 93, 864-873	5.8	6
109	Disentangling the response of lake littoral invertebrate assemblages to multiple pressures. <i>Ecological Indicators</i> , 2018 , 85, 1149-1157	5.8	7
108	Decomposing multiple pressure effects on invertebrate assemblages of boreal streams. <i>Ecological Indicators</i> , 2017 , 77, 293-303	5.8	9
107	Using streamflow observations to estimate the impact of hydrological regimes and anthropogenic water use on European stream macroinvertebrate occurrences. <i>Ecohydrology</i> , 2017 , 10, e1895	2.5	15
106	Approaches for integrated assessment of ecological and eutrophication status of surface waters in Nordic Countries. <i>Ambio</i> , 2016 , 45, 681-91	6.5	12
105	Strong land-use effects on the dispersal patterns of adult stream insects: implications for transfers of aquatic subsidies to terrestrial consumers. <i>Freshwater Biology</i> , 2016 , 61, 848-861	3.1	33
104	Benthic macroinvertebrates in lake ecological assessment: A review of methods, intercalibration and practical recommendations. <i>Science of the Total Environment</i> , 2016 , 543, 123-134	10.2	55
103	Algal blooms increase heterotrophy at the base of boreal lake food webs-evidence from fatty acid biomarkers. <i>Limnology and Oceanography</i> , 2016 , 61, 1563-1573	4.8	9
102	Adapting boreal streams to climate change: effects of riparian vegetation on water temperature and biological assemblages. <i>Freshwater Science</i> , 2016 , 35, 984-997	2	25
101	Linking degradation status with ecosystem vulnerability to environmental change. <i>Oecologia</i> , 2015 , 178, 899-913	2.9	11
100	Assessing temporal scales and patterns in time series: Comparing methods based on redundancy analysis. <i>Ecological Complexity</i> , 2015 , 22, 162-168	2.6	16
99	Linking Biodiversity, Ecosystem Functioning and Services, and Ecological Resilience. <i>Advances in Ecological Research</i> , 2015 , 53, 55-96	4.6	44
98	Spatial Patterns and Functional Redundancies in a Changing Boreal Lake Landscape. <i>Ecosystems</i> , 2015 , 18, 889-902	3.9	12
97	Effects of agricultural land use on stream assemblages: Taxon-specific responses of alpha and beta diversity. <i>Ecological Indicators</i> , 2014 , 45, 386-393	5.8	41
96	Similar resilience attributes in lakes with different management practices. <i>PLoS ONE</i> , 2014 , 9, e91881	3.7	20
95	The Swedish monitoring of surface waters: 50 years of adaptive monitoring. <i>Ambio</i> , 2014 , 43 Suppl 1, 3-18	6.5	94
94	Assessing and managing freshwater ecosystems vulnerable to environmental change. <i>Ambio</i> , 2014 , 43 Suppl 1, 113-25	6.5	49
93	Cross-taxon responses to elevated nutrients in European streams and lakes. <i>Aquatic Sciences</i> , 2014 , 76, 51-60	2.5	8

92	The impact of climate on the geographical distribution of phytoplankton species in boreal lakes. <i>Oecologia</i> , 2013 , 173, 1625-38	2.9	12
91	Optimizing stream bioassessment: habitat, season, and the impacts of land use on benthic macroinvertebrates. <i>Hydrobiologia</i> , 2013 , 704, 363-373	2.4	26
90	Community structure in boreal lakes with recurring blooms of the nuisance flagellate <i>Gonyostomum semen</i> . <i>Aquatic Sciences</i> , 2013 , 75, 447-455	2.5	16
89	Factors affecting occurrence and bloom formation of the nuisance flagellate <i>Gonyostomum semen</i> in boreal lakes. <i>Harmful Algae</i> , 2013 , 27, 60-67	5.3	26
88	Algal invasions, blooms and biodiversity in lakes: Accounting for habitat-specific responses. <i>Harmful Algae</i> , 2013 , 23, 60-69	5.3	13
87	Measuring the relative resilience of subarctic lakes to global change: redundancies of functions within and across temporal scales. <i>Journal of Applied Ecology</i> , 2013 , 50, 572-584	5.8	37
86	Effects of Dispersal-Related Factors on Species Distribution Model Accuracy for Boreal Lake Ecosystems. <i>Diversity</i> , 2013 , 5, 393-408	2.5	6
85	Zooplankton feeding on the nuisance flagellate <i>Gonyostomum semen</i> . <i>PLoS ONE</i> , 2013 , 8, e62557	3.7	20
84	Hierarchical dynamics of ecological communities: do scales of space and time match?. <i>PLoS ONE</i> , 2013 , 8, e69174	3.7	14
83	Comparison of classification-then-modelling and species-by-species modelling for predicting lake phytoplankton assemblages. <i>Ecological Modelling</i> , 2012 , 231, 11-19	3	11
82	Correcting a fundamental error in greenhouse gas accounting related to bioenergy. <i>Energy Policy</i> , 2012 , 45-222, 18-23	7.2	148
81	Patterns of temporal community turnover are spatially synchronous across boreal lakes. <i>Freshwater Biology</i> , 2012 , 57, 1782-1793	3.1	13
80	Insight on Invasions and Resilience Derived from Spatiotemporal Discontinuities of Biomass at Local and Regional Scales. <i>Ecology and Society</i> , 2012 , 17,	4.1	23
79	Temporal scales and patterns of invertebrate biodiversity dynamics in boreal lakes recovering from acidification 2012 , 22, 1172-86		32
78	Revealing the Organization of Complex Adaptive Systems through Multivariate Time Series Modeling. <i>Ecology and Society</i> , 2011 , 16,	4.1	31
77	Changes in phytoplankton, benthic invertebrate and fish assemblages of boreal lakes following invasion by <i>Gonyostomum semen</i> . <i>Freshwater Biology</i> , 2011 , 56, 1937-1948	3.1	19
76	Spatial congruency of benthic diatom, invertebrate, macrophyte, and fish assemblages in European streams 2010 , 20, 978-92		44
75	Climate Change: Defining Reference Conditions and Restoring Freshwater Ecosystems 2010 , 203-235		3

74	Tracing recovery under changing climate: response of phytoplankton and invertebrate assemblages to decreased acidification. <i>Journal of the North American Benthological Society</i> , 2010 , 29, 1472-1490		29
73	Identifying resilience mechanisms to recurrent ecosystem perturbations. <i>Oecologia</i> , 2010 , 164, 231-41	2.9	24
72	The European Water Framework Directive at the age of 10: a critical review of the achievements with recommendations for the future. <i>Science of the Total Environment</i> , 2010 , 408, 4007-19	10.2	631
71	Nature as the "natural" goal for water management: a conversation. <i>Ambio</i> , 2009 , 38, 209-14	6.5	21
70	Effects of nutrient enrichment on C and N stable isotope ratios of invertebrates, fish and their food resources in boreal streams. <i>Hydrobiologia</i> , 2009 , 628, 67-79	2.4	27
69	Response of taxonomic groups in streams to gradients in resource and habitat characteristics. <i>Journal of Applied Ecology</i> , 2009 , 46, 175-186	5.8	95
68	Detection of ecological change in stream macroinvertebrate assemblages using single metric, multimetric or multivariate approaches. <i>Ecological Indicators</i> , 2009 , 9, 659-669	5.8	33
67	Climate Change and the Future of Freshwater Biodiversity in Europe: A Primer for Policy-Makers. <i>Freshwater Reviews: A Journal of the Freshwater Biological Association</i> , 2009 , 2, 103-130		62
66	Tracking recovery trends of boreal lakes: use of multiple indicators and habitats. <i>Journal of the North American Benthological Society</i> , 2008 , 27, 529-540		28
65	Habitat-specific stability and persistence of benthic invertebrate communities in boreal lakes. <i>Fundamental and Applied Limnology</i> , 2008 , 171, 311-322	1.9	10
64	Ecological relationships between stream communities and spatial scale: implications for designing catchment-level monitoring programmes. <i>Freshwater Biology</i> , 2007 , 52, 939-958	3.1	124
63	Effects of nutrient enrichment on boreal streams: invertebrates, fungi and leaf-litter breakdown. <i>Freshwater Biology</i> , 2007 , 52, 1618-1633	3.1	37
62	Special Section: Environmental Assessment meets Landscape Ecology meets Land use Planning. <i>Freshwater Biology</i> , 2007 , 52, 907-907	3.1	4
61	Relationships Between Macroinvertebrate Assemblages of Stony Littoral Habitats and Water Chemistry Variables Indicative of Acid-stress. <i>Water, Air and Soil Pollution</i> , 2007 , 7, 323-330		10
60	A Novel Environmental Quality Criterion for Acidification in Swedish Lakes [An Application of Studies on the Relationship Between Biota and Water Chemistry. <i>Water, Air and Soil Pollution</i> , 2007 , 7, 331-338		29
59	Assessing the ecological integrity of boreal streams: a comparison of functional and structural responses. <i>Fundamental and Applied Limnology</i> , 2007 , 168, 113-125	1.9	27
58	Relationships Between Macroinvertebrate Assemblages of Stony Littoral Habitats and Water Chemistry Variables Indicative of Acid-stress 2007 , 323-330		0
57	Multiscale drivers of water chemistry of boreal lakes and streams. <i>Environmental Management</i> , 2006 , 38, 760-70	3.1	20

56	Assessing the acidity of Swedish streams using benthic macroinvertebrates and weighted averaging (WA) regression and calibration. <i>Archiv Für Hydrobiologie</i> , 2006 , 166, 343-362		3
55	A comparison of the European Water Framework Directive physical typology and RIVPACS-type models as alternative methods of establishing reference conditions for benthic macroinvertebrates 2006 , 91-105		1
54	Detection of ecological change using multiple organism groups: metrics and uncertainty 2006 , 115-137		
53	Indicators of ecological change: comparison of the early response of four organism groups to stress gradients 2006 , 139-152		2
52	Setting expectations for the ecological condition of streams: the concept of reference condition 2006 , 16, 1267-76		716
51	The effects of organic enrichment on leaf litter breakdown in three boreal streams. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2006 , 29, 1362-1366		
50	Assessment of European streams with diatoms, macrophytes, macroinvertebrates and fish: a comparative metric-based analysis of organism response to stress. <i>Freshwater Biology</i> , 2006 , 51, 1757-1785	3.1	403
49	The STAR project: context, objectives and approaches. <i>Hydrobiologia</i> , 2006 , 566, 3-29	2.4	165
48	A comparison of the European Water Framework Directive physical typology and RIVPACS-type models as alternative methods of establishing reference conditions for benthic macroinvertebrates. <i>Hydrobiologia</i> , 2006 , 566, 91-105	2.4	42
47	Estimates and comparisons of the effects of sampling variation using Bational macroinvertebrate sampling protocols on the precision of metrics used to assess ecological status. <i>Hydrobiologia</i> , 2006 , 566, 477-503	2.4	31
46	Linking organism groups major results and conclusions from the STAR project. <i>Hydrobiologia</i> , 2006 , 566, 109-113	2.4	40
45	Indicators of ecological change: comparison of the early response of four organism groups to stress gradients. <i>Hydrobiologia</i> , 2006 , 566, 139-152	2.4	50
44	Detection of ecological change using multiple organism groups: metrics and uncertainty. <i>Hydrobiologia</i> , 2006 , 566, 115-137	2.4	115
43	Estimates and comparisons of the effects of sampling variation using Bational macroinvertebrate sampling protocols on the precision of metrics used to assess ecological status 2006 , 477-503		
42	Linking organism groups major results and conclusions from the STAR project 2006 , 109-113		1
41	The STAR project: context, objectives and approaches 2006 , 3-29		2
40	Identifying, managing and monitoring conflicts between forest biodiversity conservation and other human interests in Europe. <i>Forest Policy and Economics</i> , 2005 , 7, 877-890	3.6	90
39	Additive partitioning of aquatic invertebrate species diversity across multiple spatial scales. <i>Freshwater Biology</i> , 2005 , 50, 1360-1375	3.1	88

38	Towards sustainable land use: identifying and managing the conflicts between human activities and biodiversity conservation in Europe. <i>Biodiversity and Conservation</i> , 2005 , 14, 1641-1661	3.4	189
37	A multimetric macroinvertebrate index for detecting organic pollution of streams in southern Sweden. <i>Archiv Für Hydrobiologie</i> , 2004 , 160, 487-513		22
36	Assessing Acid Stress in Swedish Boreal and Alpine Streams Using Benthic Macroinvertebrates 2004 , 129-148		1
35	Spatial scale and ecological relationships between the macroinvertebrate communities of stony habitats of streams and lakes. <i>Freshwater Biology</i> , 2004 , 49, 1179-1194	3.1	115
34	Representativity of a mid-lake surface water chemistry sample. <i>Environmental Monitoring and Assessment</i> , 2004 , 95, 221-38	3.1	11
33	Assessing acid stress in Swedish boreal and alpine streams using benthic macroinvertebrates. <i>Hydrobiologia</i> , 2004 , 516, 129-148	2.4	33
32	Detection of organic pollution of streams in southern Sweden using benthic macroinvertebrates. <i>Hydrobiologia</i> , 2004 , 516, 161-172	2.4	42
31	Local, landscape and regional factors structuring benthic macroinvertebrate assemblages in Swedish streams. <i>Landscape Ecology</i> , 2004 , 19, 501-515	4.3	105
30	Detection of Organic Pollution of Streams in Southern Sweden Using Benthic Macroinvertebrates 2004 , 161-172		1
29	Development of a prediction system for lake stony-bottom littoral macroinvertebrate communities. <i>Archiv Für Hydrobiologie</i> , 2003 , 158, 517-540		21
28	The Development of a System to Assess the Ecological Quality of Streams Based on Macroinvertebrates Design of the Sampling Programme within the AQEM Project. <i>International Review of Hydrobiology</i> , 2003 , 88, 345-361	2.3	158
27	The statistical power of selected indicator metrics using macroinvertebrates for assessing acidification and eutrophication of running waters. <i>Hydrobiologia</i> , 2000 , 422/423, 233-243	2.4	55
26	The use of biogeographical regions for partitioning variance of littoral macroinvertebrate communities. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2000 , 27, 333-339		
25	The importance of sampling effort for the assessment of ecological quality using macroinvertebrates. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2000 , 27, 326-332		1
24	Spatial scale of benthic macroinvertebrate communities in Swedish streams: variation partitioning using partial Canonical Correspondence Analysis. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2000 , 27, 382-383		
23	Ecoregions and benthic macroinvertebrate assemblages of Swedish streams. <i>Journal of the North American Benthological Society</i> , 2000 , 19, 462-474		83
22	Evaluation of the use of landscape classifications for the prediction of freshwater biota: synthesis and recommendations. <i>Journal of the North American Benthological Society</i> , 2000 , 19, 541-556		208
21	Spatial congruence between ecoregions and littoral macroinvertebrate assemblages. <i>Journal of the North American Benthological Society</i> , 2000 , 19, 475-486		34

20	The statistical power of selected indicator metrics using macroinvertebrates for assessing acidification and eutrophication of running waters 2000 , 233-243		2
19	Regional Representativeness of Swedish Reference Lakes. <i>Environmental Management</i> , 1999 , 23, 115-124	4.1	24
18	SPATIOTEMPORAL VARIABILITY OF TEMPERATE LAKE MACROINVERTEBRATE COMMUNITIES: DETECTION OF IMPACT 1998 , 8, 61-70		52
17	The fate of diatom carbon within a freshwater benthic community-a microcosm study. <i>Limnology and Oceanography</i> , 1997 , 42, 452-460	4.8	16
16	Pelagic-benthic coupling: Profundal benthic community response to spring diatom deposition in mesotrophic Lake Erken. <i>Limnology and Oceanography</i> , 1996 , 41, 636-647	4.8	70
15	Exploitation of sediment bacterial carbon by juveniles of the amphipod <i>Monoporeia affinis</i> . <i>Freshwater Biology</i> , 1994 , 32, 553-563	3.1	18
14	Effects of Deposit-Feeder Activity on Bacterial Production and Abundance in Profundal Lake Sediment. <i>Journal of the North American Benthological Society</i> , 1994 , 13, 532-539		61
13	Classification of littoral macroinvertebrate communities of Swedish reference lakes. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 1993 , 25, 512-517		3
12	Pelagic-benthic coupling-The importance of diatom interannual variability for population oscillations of <i>Monoporeia affinis</i> . <i>Limnology and Oceanography</i> , 1992 , 37, 1596-1607	4.8	50
11	Ordination of profundal zoobenthos along a trace metal pollution gradient in Northern Sweden. <i>Water, Air, and Soil Pollution</i> , 1992 , 65, 339-351	2.6	15
10	Modelling the importance of sediment bacterial carbon for profundal macroinvertebrates along a lake nutrient gradient. <i>Netherlands Journal of Aquatic Ecology</i> , 1992 , 26, 477-483		16
9	Interactions of <i>Monoporeia affinis</i> (Lindström) (Amphipoda) with sedentary Chironomidae. <i>Netherlands Journal of Aquatic Ecology</i> , 1992 , 26, 491-497		5
8	The influence of season on the classification and ordination of profundal communities of nutrient poor, oligo-mesohumic Swedish lakes using environmental data. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 1990 , 24, 646-652		
7	Long-term growth oscillations of <i>Pontoporeia affinis</i> Lindström (Crustacea: Amphipoda) in Lake Mälaren. <i>Hydrobiologia</i> , 1989 , 175, 183-194	2.4	15
6	Classification and ordination of profundal macroinvertebrate communities in nutrient poor, oligo-mesohumic lakes in relation to environmental data. <i>Freshwater Biology</i> , 1989 , 21, 375-386	3.1	63
5	Interactions between <i>Chironomus plumosus</i> (L.) and the microbial community in surficial sediments of a shallow, eutrophic lake. <i>Limnology and Oceanography</i> , 1989 , 34, 992-1003	4.8	63
4	The life history, production and food habits of <i>Pontoporeia affinis</i> Lindström (Crustacea: Amphipoda) in mesotrophic Lake Erken. <i>Hydrobiologia</i> , 1987 , 144, 277-283	2.4	17
3	Seasonal variation in diet of <i>Chironomus plumosus</i> (L.) and <i>C. anthracinus</i> Zett. (Diptera: Chironomidae) in mesotrophic Lake Erken. <i>Freshwater Biology</i> , 1987 , 17, 525-532	3.1	34

2	Feeding efficiencies of <i>Chironomus plumosus</i> (L.) and <i>C. anthracinus</i> Zett. (Diptera: Chironomidae) in mesotrophic Lake Erken. <i>Freshwater Biology</i> , 1985 , 15, 605-612	3.1	29
1	Freshwater Ecosystem Responses to Climate Change: The Euro-Limpacs Project. <i>Water Quality Measurements Series</i> , 313-354		2