
MarÃ-a de Lourdes Mora-GarcÃ-a

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1431784/publications.pdf Version: 2024-02-01



MarÃa de Lourdes

#	Article	IF	CITATIONS
1	Up-regulation of HLA class-I antigen expression and antigen-specific CTL response in cervical cancer cells by the demethylating agent hydralazine and the histone deacetylase inhibitor valproic acid. Journal of Translational Medicine, 2006, 4, 55.	4.4	63
2	An HPV 16 L1-based chimeric human papilloma virus-like particles containing a string of epitopes produced in plants is able to elicit humoral and cytotoxic T-cell activity in mice. Virology Journal, 2009, 6, 2.	3.4	62
3	Mesenchymal stromal cells derived from cervical cancer produce high amounts of adenosine to suppress cytotoxic T lymphocyte functions. Journal of Translational Medicine, 2016, 14, 302.	4.4	58
4	Immunization with an HPV-16 L1-based chimeric virus-like particle containing HPV-16 E6 and E7 epitopes elicits long-lasting prophylactic and therapeutic efficacy in an HPV-16 tumor mice model. Archives of Virology, 2014, 159, 291-305.	2.1	31
5	Cervical cancer cells produce TGF-β1 through the CD73-adenosine pathway and maintain CD73 expression through the autocrine activity of TGF-β1. Cytokine, 2019, 118, 71-79.	3.2	22
6	Detection of antibodies against a human papillomavirus (HPV) type 16 peptide that differentiate high-risk from low-risk HPV-associated low-grade squamous intraepithelial lesions. Journal of General Virology, 2004, 85, 2643-2650.	2.9	21
7	HPV-16 Infection Is Associated with a High Content of CD39 and CD73 Ectonucleotidases in Cervical Samples from Patients with CIN-1. Mediators of Inflammation, 2019, 2019, 1-13.	3.0	13
8	A novel HPV 16 L1-based chimeric virus-like particle containing E6 and E7 seroreactive epitopes permits highly specific detection of antibodies in patients with CIN 1 and HPV-16 infection. Virology Journal, 2011, 8, 59.	3.4	10
9	Peripheral blood lymphocytes from low-grade squamous intraepithelial lesions patients recognize vaccine antigens in the presence of activated dendritic cells, and produced high levels of CD8 + IFNγâ€% cells and low levels of IL-2 when induced to proliferate. Infectious Agents and Cancer, 2012, 7, 12.	‰ ₽.â€ ‰T	6
10	Detection of CD39 and a Highly Glycosylated Isoform of Soluble CD73 in the Plasma of Patients with Cervical Cancer: Correlation with Disease Progression. Mediators of Inflammation, 2020, 2020, 1-14.	3.0	6
11	Sodium Caseinate (CasNa) Induces Mobilization of Hematopoietic Stem Cells in a BALB/c Mouse Model. Medical Science Monitor Basic Research, 2015, 21, 206-212.	2.6	4
12	GLMEEMSAL epitope common in different isoforms of hMena elicits in vitro activation of cytotoxic T cells and stimulates specific antitumor immunity in BALB/c mice. International Immunopharmacology, 2018, 56, 291-300.	3.8	0