

MarÃ-a de Lourdes Mora-GarcÃ-a

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1431784/publications.pdf>

Version: 2024-02-01

12
papers

296
citations

1163117

8
h-index

1281871

11
g-index

12
all docs

12
docs citations

12
times ranked

559
citing authors

#	ARTICLE	IF	CITATIONS
1	Detection of CD39 and a Highly Glycosylated Isoform of Soluble CD73 in the Plasma of Patients with Cervical Cancer: Correlation with Disease Progression. <i>Mediators of Inflammation</i> , 2020, 2020, 1-14.	3.0	6
2	HPV-16 Infection Is Associated with a High Content of CD39 and CD73 Ectonucleotidases in Cervical Samples from Patients with CIN-1. <i>Mediators of Inflammation</i> , 2019, 2019, 1-13.	3.0	13
3	Cervical cancer cells produce TGF- β 1 through the CD73-adenosine pathway and maintain CD73 expression through the autocrine activity of TGF- β 1. <i>Cytokine</i> , 2019, 118, 71-79.	3.2	22
4	GLMEEMSAL epitope common in different isoforms of hMena elicits in vitro activation of cytotoxic T cells and stimulates specific antitumor immunity in BALB/c mice. <i>International Immunopharmacology</i> , 2018, 56, 291-300.	3.8	0
5	Mesenchymal stromal cells derived from cervical cancer produce high amounts of adenosine to suppress cytotoxic T lymphocyte functions. <i>Journal of Translational Medicine</i> , 2016, 14, 302.	4.4	58
6	Sodium Caseinate (CasNa) Induces Mobilization of Hematopoietic Stem Cells in a BALB/c Mouse Model. <i>Medical Science Monitor Basic Research</i> , 2015, 21, 206-212.	2.6	4
7	Immunization with an HPV-16 L1-based chimeric virus-like particle containing HPV-16 E6 and E7 epitopes elicits long-lasting prophylactic and therapeutic efficacy in an HPV-16 tumor mice model. <i>Archives of Virology</i> , 2014, 159, 291-305.	2.1	31
8	Peripheral blood lymphocytes from low-grade squamous intraepithelial lesions patients recognize vaccine antigens in the presence of activated dendritic cells, and produced high levels of CD8 α ⁺ IFN γ ⁺ T cells and low levels of IL-2 when induced to proliferate. <i>Infectious Agents and Cancer</i> , 2012, 7, 12.	2.3	6
9	A novel HPV 16 L1-based chimeric virus-like particle containing E6 and E7 seroreactive epitopes permits highly specific detection of antibodies in patients with CIN 1 and HPV-16 infection. <i>Virology Journal</i> , 2011, 8, 59.	3.4	10
10	An HPV 16 L1-based chimeric human papilloma virus-like particles containing a string of epitopes produced in plants is able to elicit humoral and cytotoxic T-cell activity in mice. <i>Virology Journal</i> , 2009, 6, 2.	3.4	62
11	Up-regulation of HLA class-I antigen expression and antigen-specific CTL response in cervical cancer cells by the demethylating agent hydralazine and the histone deacetylase inhibitor valproic acid. <i>Journal of Translational Medicine</i> , 2006, 4, 55.	4.4	63
12	Detection of antibodies against a human papillomavirus (HPV) type 16 peptide that differentiate high-risk from low-risk HPV-associated low-grade squamous intraepithelial lesions. <i>Journal of General Virology</i> , 2004, 85, 2643-2650.	2.9	21