

# Kristian Stengaard-Pedersen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1432873/publications.pdf>

Version: 2024-02-01

75  
papers

3,164  
citations

201674

27  
h-index

161849

54  
g-index

76  
all docs

76  
docs citations

76  
times ranked

4449  
citing authors

#	ARTICLE	IF	CITATIONS
1	Treatment of Lateral Epicondylitis With Platelet-Rich Plasma, Glucocorticoid, or Saline. <i>American Journal of Sports Medicine</i> , 2013, 41, 625-635.	4.2	305
2	Inherited Deficiency of Mannan-Binding Lectin-Associated Serine Protease 2. <i>New England Journal of Medicine</i> , 2003, 349, 554-560.	27.0	239
3	Comparative Effectiveness of Injection Therapies in Lateral Epicondylitis. <i>American Journal of Sports Medicine</i> , 2013, 41, 1435-1446.	4.2	213
4	Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CIMESTR trial. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1789-1795.	0.9	168
5	Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: An investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> . 2006. 54. 1401-1409.	6.7	160
6	Telehealth Followup Strategy for Tight Control of Disease Activity in Rheumatoid Arthritis: Results of a Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2018, 70, 353-360.	3.4	124
7	Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> . 2014. 73. 654-661.	0.9	108
8	Plasma Adiponectin in Patients with Active, Early, and Chronic Rheumatoid Arthritis Who Are Steroid- and Disease-Modifying Antirheumatic Drug-Naive Compared with Patients with Osteoarthritis and Controls. <i>Journal of Rheumatology</i> , 2009, 36, 1885-1891.	2.0	106
9	Increased Interleukin 21 (IL-21) and IL-23 Are Associated with Increased Disease Activity and with Radiographic Status in Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2010, 37, 2014-2020.	2.0	94
10	Compliance with methotrexate treatment in patients with rheumatoid arthritis: influence of patients' beliefs about the medicine. A prospective cohort study. <i>Rheumatology International</i> , 2010, 30, 1441-1448.	3.0	72
11	Intra-articular hyaluronan is without clinical effect in knee osteoarthritis: a multicentre, randomised, placebo-controlled, double-blind study of 337 patients followed for 1 year. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1097.2-1102.	0.9	71
12	Short- and long-term efficacy of intra-articular injections with betamethasone as part of a treat-to-target strategy in early rheumatoid arthritis: impact of joint area, repeated injections, MRI findings, anti-CCP, IgM-RF and CRP. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 851-856.	0.9	70
13	Effect of Magnetic Resonance Imaging vs Conventional Treat-to-Target Strategies on Disease Activity Remission and Radiographic Progression in Rheumatoid Arthritis. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 461.	7.4	70
14	Type 1 Modic changes was a significant risk factor for 1-year outcome in sick-listed low back pain patients: a nested cohort study using magnetic resonance imaging of the lumbar spine. <i>Spine Journal</i> , 2014, 14, 2568-2581.	1.3	63
15	CXCL13 predicts disease activity in early rheumatoid arthritis and could be an indicator of the therapeutic 'window of opportunity'. <i>Arthritis Research and Therapy</i> , 2014, 16, 434.	3.5	61
16	A 6-months, randomised, placebo-controlled evaluation of efficacy and tolerability of a low-dose 7-day buprenorphine transdermal patch in osteoarthritis patients naïve to potent opioids. <i>Scandinavian Journal of Pain</i> , 2010, 1, 122-141.	1.3	55
17	Inhibition of Enkephalin Binding to Opiate Receptors by Zinc Ions: Possible Physiological Importance in the Brain. <i>Acta Pharmacologica Et Toxicologica</i> , 1982, 50, 213-220.	0.0	54
18	Overexpression of microRNA-155 increases IL-21 mediated STAT3 signaling and IL-21 production in systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , 2015, 17, 154.	3.5	52

#	ARTICLE	IF	CITATIONS
19	One-year prognosis in sick-listed low back pain patients with and without radiculopathy. Prognostic factors influencing pain and disability. <i>Spine Journal</i> , 2010, 10, 659-675.	1.3	50
20	Polymorphisms within Toll-like receptors are associated with systemic lupus erythematosus in a cohort of Danish females. <i>Rheumatology</i> , 2014, 53, 48-55.	1.9	50
21	A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 867-875.	0.9	49
22	Body Fat Percentage, Waist Circumference, and Obesity As Risk Factors for Rheumatoid Arthritis: A Danish Cohort Study. <i>Arthritis Care and Research</i> , 2019, 71, 777-786.	3.4	40
23	Prediction model for unsuccessful return to work after hospital-based intervention in low back pain patients. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 140.	1.9	36
24	The Lectin Pathway of Complement Activation in Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2018, 45, 1136-1144.	2.0	36
25	Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , 2018, 45, 53-61.	2.0	35
26	The C3dg Fragment of Complement Is Superior to Conventional C3 as a Diagnostic Biomarker in Systemic Lupus Erythematosus. <i>Frontiers in Immunology</i> , 2018, 9, 581.	4.8	32
27	Increased plasma levels of IL-21 and IL-23 in spondyloarthritis are not associated with clinical and MRI findings. <i>Rheumatology International</i> , 2012, 32, 387-393.	3.0	30
28	Soluble OX40L is associated with presence of autoantibodies in early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014, 16, 474.	3.5	28
29	Ficolin-3 Deficiency Is Associated with Disease and an Increased Risk of Systemic Lupus Erythematosus. <i>Journal of Clinical Immunology</i> , 2019, 39, 421-429.	3.8	28
30	Postural balance in low back pain patients: criterion-related validity of centre of pressure assessed on a portable force platform. <i>European Spine Journal</i> , 2012, 21, 425-431.	2.2	27
31	Non-Synonymous Polymorphisms in the FCN1 Gene Determine Ligand-Binding Ability and Serum Levels of M-Ficolin. <i>PLoS ONE</i> , 2012, 7, e50585.	2.5	27
32	The interleukin-20 receptor axis in early rheumatoid arthritis: novel links between disease-associated autoantibodies and radiographic progression. <i>Arthritis Research and Therapy</i> , 2016, 18, 61.	3.5	26
33	Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1707-1715.	0.9	26
34	Polymorphisms in the MASP1 Gene Are Associated with Serum Levels of MASP-1, MASP-3, and MAP44. <i>PLoS ONE</i> , 2013, 8, e73317.	2.5	26
35	Positive predictive value of first-time rheumatoid arthritis diagnoses and their serological subtypes in the Danish National Patient Registry. <i>Clinical Epidemiology</i> , 2018, Volume 10, 1709-1720.	3.0	25
36	Human Endogenous Retroviral Genetic Element With Immunosuppressive Activity in Both Human Autoimmune Diseases and Experimental Arthritis. <i>Arthritis and Rheumatology</i> , 2017, 69, 398-409.	5.6	23

#	ARTICLE	IF	CITATIONS
37	Brief Report: Mâ€Ficolin Levels Reflect Disease Activity and Predict Remission in Early Rheumatoid Arthritis. <i>Arthritis and Rheumatism</i> , 2013, 65, 3045-3050.	6.7	22
38	Expression of soluble CD83 in plasma from early-stage rheumatoid arthritis patients is not modified by anti-TNF-Î± therapy. <i>Cytokine</i> , 2017, 96, 1-7.	3.2	22
39	Low back pain may be caused by disturbed pain regulation. <i>European Journal of Pain</i> , 2010, 14, 514-522.	2.8	21
40	Postural balance in low back pain patients: Intra-session reliability of center of pressure on a portable force platform and of the one leg stand test. <i>Gait and Posture</i> , 2011, 34, 213-217.	1.4	20
41	Calprotectin in patients with chronic rheumatoid arthritis correlates with disease activity and responsiveness to methotrexate. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 62-67.	1.2	20
42	Mannose-Binding Lectin Gene Polymorphisms Are Associated with Disease Activity and Physical Disability in Untreated, Anti-Cyclic Citrullinated Peptide-Positive Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2009, 36, 731-735.	2.0	19
43	Circulating surfactant protein -D is low and correlates negatively with systemic inflammation in early, untreated rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2010, 12, R39.	3.5	19
44	Levels in Plasma of the Serine Proteases and Associated Proteins of the Lectin Pathway Are Altered in Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2015, 42, 948-951.	2.0	19
45	Patient self-assessment of flare in rheumatoid arthritis: criterion and concurrent validity of the Flare instrument. <i>Clinical Rheumatology</i> , 2016, 35, 467-471.	2.2	19
46	Effects of maternal health anxiety on childrenâ€™s health complaints, emotional symptoms, and quality of life. <i>European Child and Adolescent Psychiatry</i> , 2017, 26, 591-601.	4.7	19
47	SKG arthritis as a model for evaluating therapies in rheumatoid arthritis with special focus on bone changes. <i>Rheumatology International</i> , 2013, 33, 1127-1133.	3.0	18
48	A non-synonymous single-nucleotide polymorphism in the gene encoding Toll-like Receptor 3 (TLR3) is associated with sero-negative Rheumatoid Arthritis (RA) in a Danish population. <i>BMC Research Notes</i> , 2014, 7, 716.	1.4	18
49	Back pain was less explained than leg pain: a cross-sectional study using magnetic resonance imaging in low back pain patients with and without radiculopathy. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 374.	1.9	17
50	A disintegrin and metalloprotease-17 and galectin-9 are important regulators of local 4-1BB activity and disease outcome in rheumatoid arthritis. <i>Rheumatology</i> , 2016, 55, 1871-1879.	1.9	17
51	Patterns of Magnetic Resonance Imaging Bone Erosion in Rheumatoid Arthritis â€” Which Bones Are Most Frequently Involved and Show the Most Change?: Figure 1.. <i>Journal of Rheumatology</i> , 2011, 38, 2014-2017.	2.0	15
52	Macrophage activity assessed by soluble CD163 in early rheumatoid arthritis: association with disease activity but different response patterns to synthetic and biologic DMARDs. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, 498-502.	0.8	15
53	Cartilage Oligomeric Matrix Protein Associates Differentially with Erosions and Synovitis and Has a Different Temporal Course in Cyclic Citrullinated Peptide Antibody (Anti-CCP)-positive versus Anti-CCP-negative Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2011, 38, 1563-1568.	2.0	14
54	The Dâ€vitamin metabolite 1,25( <sc&gt;oh&lt; <i="" active="" activity="" and="" antibodies="" antiâ€citrullinated="" arthritis="" associated="" disease="" early="" in="" is="" naãˆve,="" patients.="" protein="" rheumatoid="" sc&gt;)&lt;sub&gt;2&lt;="" serum="" sub&gt;d="" treatment="" with="">Scandinavian Journal of Immunology, 2018, 88, e12704.</sc&gt;oh&lt;>	2.7	14

#	ARTICLE	IF	CITATIONS
55	Uncoupling of Collagen II Metabolism in Newly Diagnosed, Untreated Rheumatoid Arthritis Is Linked to Inflammation and Antibodies Against Cyclic Citrullinated Peptides. <i>Journal of Rheumatology</i> , 2010, 37, 1113-1120.	2.0	13
56	Patient-self assessment of flare in rheumatoid arthritis: translation and reliability of the Flare instrument. <i>Clinical Rheumatology</i> , 2016, 35, 1053-1058.	2.2	13
57	CRP genotype and haplotype associations with serum C-reactive protein level and DAS28 in untreated early rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2014, 16, 475.	3.5	12
58	Changes in Soluble CD18 in Murine Autoimmune Arthritis and Rheumatoid Arthritis Reflect Disease Establishment and Treatment Response. <i>PLoS ONE</i> , 2016, 11, e0148486.	2.5	12
59	Bone Formation and Resorption Are Both Increased in Experimental Autoimmune Arthritis. <i>PLoS ONE</i> , 2012, 7, e53034.	2.5	12
60	Impact of a magnetic resonance imaging-guided treat-to-target strategy on disease activity and progression in patients with rheumatoid arthritis (the IMAGINE-RA trial): study protocol for a randomized controlled trial. <i>Trials</i> , 2015, 16, 178.	1.6	11
61	Soluble CD206 plasma levels in rheumatoid arthritis reflect decrease in disease activity. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2017, 77, 385-389.	1.2	11
62	Programmed death ligand 2 – A link between inflammation and bone loss in rheumatoid arthritis. <i>Journal of Translational Autoimmunity</i> , 2020, 3, 100028.	4.0	10
63	The Programmed Death-1 Pathway Counter-Regulates Inflammation-Induced Osteoclast Activity in Clinical and Experimental Settings. <i>Frontiers in Immunology</i> , 2022, 13, 773946.	4.8	9
64	Insulin-like growth factor I receptor density on CD4+T-lymphocytes from active early steroid- and DMARD-naïve rheumatoid arthritis patients is up-regulated and not influenced by 1 year of clinically effective treatment. <i>Rheumatology International</i> , 2012, 32, 501-504.	3.0	8
65	Systemic but No Local Effects of Combined Zoledronate and Parathyroid Hormone Treatment in Experimental Autoimmune Arthritis. <i>PLoS ONE</i> , 2014, 9, e92359.	2.5	8
66	Characterization of DNA-protein complexes by nanoparticle tracking analysis and their association with systemic lupus erythematosus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	7
67	One-year progression of erosive disease in patients with anti-citrullinated peptide antibodies and arthralgia. <i>Joint Bone Spine</i> , 2020, 87, 181-183.	1.6	6
68	Predictors of joint damage progression and stringent remission in patients with established rheumatoid arthritis in clinical remission. <i>Rheumatology</i> , 2021, 60, 380-391.	1.9	6
69	Diffuse central sensitization in low back patients. <i>Medicine (United States)</i> , 2020, 99, e22198.	1.0	4
70	Calcified cartilage in patients with osteoarthritis of the hip compared to that of healthy subjects. A design-based histological study. <i>Bone</i> , 2021, 143, 115660.	2.9	4
71	Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , 2021, 41, 543-549.	3.0	4
72	Methotrexate utilization in Rheumatoid arthritis. A register-based cohort-study of treatment re-starts after gabs of at least 90 days. <i>SpringerPlus</i> , 2015, 4, 227.	1.2	3

#	ARTICLE	IF	CITATIONS
73	A positive <i>Helicobacter pylori</i> test is associated with low spondylarthritis incidence in a Danish historical cohort study. <i>Rheumatology International</i> , 2020, 40, 359-366.	3.0	2
74	Methotrexate pharmacokinetic is influenced by co-administration of cyclosporin in rheumatoid arthritis patients. Results from a randomized clinical trial. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 185-190.	1.2	1
75	SAT0092â€¦LOW SERUM LEVEL OF VITAMIN D AT TIME OF DIAGNOSIS IS ASSOCIATED WITH HIGHER ONE-YEAR REMISSION RATE IN PATIENTS WITH NEWLY DIAGNOSED RA, TREATED AGGRESSIVELY DURING FOLLOW-UP: POST-HOC ANALYSES OF THE CIMESTRA TRIAL. , 2019, , .		0