

Kristian Stengaard-Pedersen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1432873/publications.pdf>

Version: 2024-02-01

75
papers

3,164
citations

201674

27
h-index

161849

54
g-index

76
all docs

76
docs citations

76
times ranked

4449
citing authors

#	ARTICLE	IF	CITATIONS
1	The Programmed Death-1 Pathway Counter-Regulates Inflammation-Induced Osteoclast Activity in Clinical and Experimental Settings. <i>Frontiers in Immunology</i> , 2022, 13, 773946.	4.8	9
2	Calcified cartilage in patients with osteoarthritis of the hip compared to that of healthy subjects. A design-based histological study. <i>Bone</i> , 2021, 143, 115660.	2.9	4
3	Predictors of joint damage progression and stringent remission in patients with established rheumatoid arthritis in clinical remission. <i>Rheumatology</i> , 2021, 60, 380-391.	1.9	6
4	Characterization of DNA-protein complexes by nanoparticle tracking analysis and their association with systemic lupus erythematosus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	7
5	Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , 2021, 41, 543-549.	3.0	4
6	One-year progression of erosive disease in patients with anti-citrullinated peptide antibodies and arthralgia. <i>Joint Bone Spine</i> , 2020, 87, 181-183.	1.6	6
7	Methotrexate pharmacokinetic is influenced by co-administration of cyclosporin in rheumatoid arthritis patients. Results from a randomized clinical trial. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 185-190.	1.2	1
8	Programmed death ligand 2 - A link between inflammation and bone loss in rheumatoid arthritis. <i>Journal of Translational Autoimmunity</i> , 2020, 3, 100028.	4.0	10
9	A positive <i>Helicobacter pylori</i> test is associated with low spondylarthritis incidence in a Danish historical cohort study. <i>Rheumatology International</i> , 2020, 40, 359-366.	3.0	2
10	Diffuse central sensitization in low back patients. <i>Medicine (United States)</i> , 2020, 99, e22198.	1.0	4
11	Body Fat Percentage, Waist Circumference, and Obesity As Risk Factors for Rheumatoid Arthritis: A Danish Cohort Study. <i>Arthritis Care and Research</i> , 2019, 71, 777-786.	3.4	40
12	Effect of Magnetic Resonance Imaging vs Conventional Treat-to-Target Strategies on Disease Activity Remission and Radiographic Progression in Rheumatoid Arthritis. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 461.	7.4	70
13	Ficolin-3 Deficiency Is Associated with Disease and an Increased Risk of Systemic Lupus Erythematosus. <i>Journal of Clinical Immunology</i> , 2019, 39, 421-429.	3.8	28
14	SAT0092...LOW SERUM LEVEL OF VITAMIN D AT TIME OF DIAGNOSIS IS ASSOCIATED WITH HIGHER ONE-YEAR REMISSION RATE IN PATIENTS WITH NEWLY DIAGNOSED RA, TREATED AGGRESSIVELY DURING FOLLOW-UP: POST-HOC ANALYSES OF THE CIMESTRA TRIAL. , 2019, , .		0
15	Tele-Health Followup Strategy for Tight Control of Disease Activity in Rheumatoid Arthritis: Results of a Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2018, 70, 353-360.	3.4	124
16	Calprotectin in patients with chronic rheumatoid arthritis correlates with disease activity and responsiveness to methotrexate. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 62-67.	1.2	20
17	Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , 2018, 45, 53-61.	2.0	35
18	Positive predictive value of first-time rheumatoid arthritis diagnoses and their serological subtypes in the Danish National Patient Registry. <i>Clinical Epidemiology</i> , 2018, Volume 10, 1709-1720.	3.0	25

#	ARTICLE	IF	CITATIONS
19	The Dâ€vitamin metabolite 1,25(<sc>OH</sc>)₂D in serum is associated with disease activity and Antiâ€CitruLLinated Protein Antibodies in active and treatment naÃve, early Rheumatoid Arthritis Patients. Scandinavian Journal of Immunology, 2018, 88, e12704.	2.7	14
20	The C3dg Fragment of Complement Is Superior to Conventional C3 as a Diagnostic Biomarker in Systemic Lupus Erythematosus. Frontiers in Immunology, 2018, 9, 581.	4.8	32
21	The Lectin Pathway of Complement Activation in Patients with Systemic Lupus Erythematosus. Journal of Rheumatology, 2018, 45, 1136-1144.	2.0	36
22	Soluble CD206 plasma levels in rheumatoid arthritis reflect decrease in disease activity. Scandinavian Journal of Clinical and Laboratory Investigation, 2017, 77, 385-389.	1.2	11
23	Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. Annals of the Rheumatic Diseases, 2017, 76, 1707-1715.	0.9	26
24	Expression of soluble CD83 in plasma from early-stage rheumatoid arthritis patients is not modified by anti-TNF-Î± therapy. Cytokine, 2017, 96, 1-7.	3.2	22
25	Effects of maternal health anxiety on childrenâ€™s health complaints, emotional symptoms, and quality of life. European Child and Adolescent Psychiatry, 2017, 26, 591-601.	4.7	19
26	Human Endogenous Retroviral Genetic Element With Immunosuppressive Activity in Both Human Autoimmune Diseases and Experimental Arthritis. Arthritis and Rheumatology, 2017, 69, 398-409.	5.6	23
27	Changes in Soluble CD18 in Murine Autoimmune Arthritis and Rheumatoid Arthritis Reflect Disease Establishment and Treatment Response. PLoS ONE, 2016, 11, e0148486.	2.5	12
28	A disintegrin and metalloprotease-17 and galectin-9 are important regulators of local 4-1BB activity and disease outcome in rheumatoid arthritis. Rheumatology, 2016, 55, 1871-1879.	1.9	17
29	The interleukin-20 receptor axis in early rheumatoid arthritis: novel links between disease-associated autoantibodies and radiographic progression. Arthritis Research and Therapy, 2016, 18, 61.	3.5	26
30	Patient self-assessment of flare in rheumatoid arthritis: criterion and concurrent validity of the Flare instrument. Clinical Rheumatology, 2016, 35, 467-471.	2.2	19
31	Patient-self assessment of flare in rheumatoid arthritis: translation and reliability of the Flare instrument. Clinical Rheumatology, 2016, 35, 1053-1058.	2.2	13
32	Overexpression of microRNA-155 increases IL-21 mediated STAT3 signaling and IL-21 production in systemic lupus erythematosus. Arthritis Research and Therapy, 2015, 17, 154.	3.5	52
33	Back pain was less explained than leg pain: a cross-sectional study using magnetic resonance imaging in low back pain patients with and without radiculopathy. BMC Musculoskeletal Disorders, 2015, 16, 374.	1.9	17
34	Methotrexate utilization in Rheumatoid arthritis. A register-based cohort-study of treatment re-starts after gabs of at least 90Ãdays. SpringerPlus, 2015, 4, 227.	1.2	3
35	Impact of a magnetic resonance imaging-guided treat-to-target strategy on disease activity and progression in patients with rheumatoid arthritis (the IMAGINE-RA trial): study protocol for a randomized controlled trial. Trials, 2015, 16, 178.	1.6	11
36	A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. Annals of the Rheumatic Diseases, 2015, 74, 867-875.	0.9	49

#	ARTICLE	IF	CITATIONS
37	Levels in Plasma of the Serine Proteases and Associated Proteins of the Lectin Pathway Are Altered in Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2015, 42, 948-951.	2.0	19
38	Macrophage activity assessed by soluble CD163 in early rheumatoid arthritis: association with disease activity but different response patterns to synthetic and biologic DMARDs. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, 498-502.	0.8	15
39	Systemic but No Local Effects of Combined Zoledronate and Parathyroid Hormone Treatment in Experimental Autoimmune Arthritis. <i>PLoS ONE</i> , 2014, 9, e92359.	2.5	8
40	CXCL13 predicts disease activity in early rheumatoid arthritis and could be an indicator of the therapeutic 'window of opportunity'. <i>Arthritis Research and Therapy</i> , 2014, 16, 434.	3.5	61
41	Soluble OX40L is associated with presence of autoantibodies in early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014, 16, 474.	3.5	28
42	CRP genotype and haplotype associations with serum C-reactive protein level and DAS28 in untreated early rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2014, 16, 475.	3.5	12
43	Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 654-661.	0.9	108
44	A non-synonymous single-nucleotide polymorphism in the gene encoding Toll-like Receptor 3 (TLR3) is associated with sero-negative Rheumatoid Arthritis (RA) in a Danish population. <i>BMC Research Notes</i> , 2014, 7, 716.	1.4	18
45	Polymorphisms within Toll-like receptors are associated with systemic lupus erythematosus in a cohort of Danish females. <i>Rheumatology</i> , 2014, 53, 48-55.	1.9	50
46	Type 1 Modic changes was a significant risk factor for 1-year outcome in sick-listed low back pain patients: a nested cohort study using magnetic resonance imaging of the lumbar spine. <i>Spine Journal</i> , 2014, 14, 2568-2581.	1.3	63
47	Prediction model for unsuccessful return to work after hospital-based intervention in low back pain patients. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 140.	1.9	36
48	Treatment of Lateral Epicondylitis With Platelet-Rich Plasma, Glucocorticoid, or Saline. <i>American Journal of Sports Medicine</i> , 2013, 41, 625-635.	4.2	305
49	SKG arthritis as a model for evaluating therapies in rheumatoid arthritis with special focus on bone changes. <i>Rheumatology International</i> , 2013, 33, 1127-1133.	3.0	18
50	Comparative Effectiveness of Injection Therapies in Lateral Epicondylitis. <i>American Journal of Sports Medicine</i> , 2013, 41, 1435-1446.	4.2	213
51	Brief Report: Mâ€Ficolin Levels Reflect Disease Activity and Predict Remission in Early Rheumatoid Arthritis. <i>Arthritis and Rheumatism</i> , 2013, 65, 3045-3050.	6.7	22
52	Polymorphisms in the MASP1 Gene Are Associated with Serum Levels of MASP-1, MASP-3, and MAP44. <i>PLoS ONE</i> , 2013, 8, e73317.	2.5	26
53	Short- and long-term efficacy of intra-articular injections with betamethasone as part of a treat-to-target strategy in early rheumatoid arthritis: impact of joint area, repeated injections, MRI findings, anti-CCP, IgM-RF and CRP. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 851-856.	0.9	70
54	Postural balance in low back pain patients: criterion-related validity of centre of pressure assessed on a portable force platform. <i>European Spine Journal</i> , 2012, 21, 425-431.	2.2	27

#	ARTICLE	IF	CITATIONS
55	Increased plasma levels of IL-21 and IL-23 in spondyloarthritis are not associated with clinical and MRI findings. <i>Rheumatology International</i> , 2012, 32, 387-393.	3.0	30
56	Insulin-like growth factor I receptor density on CD4+T-lymphocytes from active early steroid- and DMARD-naïve rheumatoid arthritis patients is up-regulated and not influenced by 1 year of clinically effective treatment. <i>Rheumatology International</i> , 2012, 32, 501-504.	3.0	8
57	Non-Synonymous Polymorphisms in the FCN1 Gene Determine Ligand-Binding Ability and Serum Levels of M-Ficolin. <i>PLoS ONE</i> , 2012, 7, e50585.	2.5	27
58	Bone Formation and Resorption Are Both Increased in Experimental Autoimmune Arthritis. <i>PLoS ONE</i> , 2012, 7, e53034.	2.5	12
59	Postural balance in low back pain patients: Intra-session reliability of center of pressure on a portable force platform and of the one leg stand test. <i>Gait and Posture</i> , 2011, 34, 213-217.	1.4	20
60	Cartilage Oligomeric Matrix Protein Associates Differentially with Erosions and Synovitis and Has a Different Temporal Course in Cyclic Citrullinated Peptide Antibody (Anti-CCP)-positive versus Anti-CCP-negative Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2011, 38, 1563-1568.	2.0	14
61	Patterns of Magnetic Resonance Imaging Bone Erosion in Rheumatoid Arthritis – Which Bones Are Most Frequently Involved and Show the Most Change?: Figure 1.. <i>Journal of Rheumatology</i> , 2011, 38, 2014-2017.	2.0	15
62	Compliance with methotrexate treatment in patients with rheumatoid arthritis: influence of patients' beliefs about the medicine. A prospective cohort study. <i>Rheumatology International</i> , 2010, 30, 1441-1448.	3.0	72
63	Low back pain may be caused by disturbed pain regulation. <i>European Journal of Pain</i> , 2010, 14, 514-522.	2.8	21
64	Uncoupling of Collagen II Metabolism in Newly Diagnosed, Untreated Rheumatoid Arthritis Is Linked to Inflammation and Antibodies Against Cyclic Citrullinated Peptides. <i>Journal of Rheumatology</i> , 2010, 37, 1113-1120.	2.0	13
65	Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CESTRA trial. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1789-1795.	0.9	168
66	Intra-articular hyaluronan is without clinical effect in knee osteoarthritis: a multicentre, randomised, placebo-controlled, double-blind study of 337 patients followed for 1 year. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1097.2-1102.	0.9	71
67	A 6-months, randomised, placebo-controlled evaluation of efficacy and tolerability of a low-dose 7-day buprenorphine transdermal patch in osteoarthritis patients naïve to potent opioids. <i>Scandinavian Journal of Pain</i> , 2010, 1, 122-141.	1.3	55
68	One-year prognosis in sick-listed low back pain patients with and without radiculopathy. Prognostic factors influencing pain and disability. <i>Spine Journal</i> , 2010, 10, 659-675.	1.3	50
69	Increased Interleukin 21 (IL-21) and IL-23 Are Associated with Increased Disease Activity and with Radiographic Status in Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2010, 37, 2014-2020.	2.0	94
70	Circulating surfactant protein -D is low and correlates negatively with systemic inflammation in early, untreated rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2010, 12, R39.	3.5	19
71	Plasma Adiponectin in Patients with Active, Early, and Chronic Rheumatoid Arthritis Who Are Steroid- and Disease-Modifying Antirheumatic Drug-Naïve Compared with Patients with Osteoarthritis and Controls. <i>Journal of Rheumatology</i> , 2009, 36, 1885-1891.	2.0	106
72	Mannose-Binding Lectin Gene Polymorphisms Are Associated with Disease Activity and Physical Disability in Untreated, Anti-Cyclic Citrullinated Peptide-Positive Patients with Early Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2009, 36, 731-735.	2.0	19

#	ARTICLE	IF	CITATIONS
73	Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: An investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> , 2006, 54, 1401-1409.	6.7	160
74	Inherited Deficiency of Mannan-Binding Lectin-Associated Serine Protease 2. <i>New England Journal of Medicine</i> , 2003, 349, 554-560.	27.0	239
75	Inhibition of Enkephalin Binding to Opiate Receptors by Zinc Ions: Possible Physiological Importance in the Brain. <i>Acta Pharmacologica Et Toxicologica</i> , 1982, 50, 213-220.	0.0	54