

J Antoinette Killian

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1442085/j-antoinette-killian-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

134 papers	9,731 citations	59 h-index	96 g-index
139 ext. papers	10,458 ext. citations	4.2 avg, IF	5.97 L-index

#	Paper	IF	Citations
134	Synthesis and Evaluation of a Library of Alternating Amphipathic Copolymers to Solubilize and Study Membrane Proteins.. <i>Biomacromolecules</i> , 2022 ,	6.9	2
133	Mass Photometry of Membrane Proteins. <i>CheM</i> , 2021 , 7, 224-236	16.2	16
132	Factors influencing the solubilization of membrane proteins from Escherichia coli membranes by styrene-maleic acid copolymers. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2020 , 1862, 183125	3.8	16
131	Iterative RAFT-Mediated Copolymerization of Styrene and Maleic Anhydride toward Sequence- and Length-Controlled Copolymers and Their Applications for Solubilizing Lipid Membranes. <i>Biomacromolecules</i> , 2020 , 21, 3287-3300	6.9	15
130	A Broad-Spectrum Antiviral Peptide Blocks Infection of Viruses by Binding to Phosphatidylserine in the Viral Envelope. <i>Cells</i> , 2020 , 9,	7.9	6
129	The small molecule inhibitor anle145c thermodynamically traps human islet amyloid peptide in the form of non-cytotoxic oligomers. <i>Scientific Reports</i> , 2019 , 9, 19023	4.9	8
128	A simple and convenient method for the hydrolysis of styrene-maleic anhydride copolymers to styrene-maleic acid copolymers. <i>Chemistry and Physics of Lipids</i> , 2019 , 218, 85-90	3.7	12
127	A single mutation on the human amyloid polypeptide modulates fibril growth and affects the mechanism of amyloid-induced membrane damage. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2018 , 1860, 1783-1792	3.8	9
126	Bacillus subtilis MraY in detergent-free system of nanodiscs wrapped by styrene-maleic acid copolymers. <i>PLoS ONE</i> , 2018 , 13, e0206692	3.7	3
125	Membrane Solubilization by Styrene-Maleic Acid Copolymers: Delineating the Role of Polymer Length. <i>Biophysical Journal</i> , 2018 , 115, 129-138	2.9	19
124	Proton-Detected Solid-State NMR Spectroscopy of a Zinc Diffusion Facilitator Protein in Native Nanodiscs. <i>Angewandte Chemie - International Edition</i> , 2017 , 56, 2508-2512	16.4	63
123	Solubilization of lipids and lipid phases by the styrene-maleic acid copolymer. <i>European Biophysics Journal</i> , 2017 , 46, 91-101	1.9	57
122	Solubilization of human cells by the styrene-maleic acid copolymer: Insights from fluorescence microscopy. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2017 , 1859, 2155-2160	3.8	15
121	The effectiveness of styrene-maleic acid (SMA) copolymers for solubilisation of integral membrane proteins from SMA-accessible and SMA-resistant membranes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2017 , 1859, 2133-2143	3.8	52
120	Protonendetektierte Festkörper-NMR-Spektroskopie an einem Zinktransporter-Membranprotein in nativen Nanoscheiben. <i>Angewandte Chemie</i> , 2017 , 129, 2549-2553	3.6	5
119	Residue specific effects of human islet polypeptide amyloid on self-assembly and on cell toxicity. <i>Biochimie</i> , 2017 , 142, 22-30	4.6	20
118	Effect of Polymer Composition and pH on Membrane Solubilization by Styrene-Maleic Acid Copolymers. <i>Biophysical Journal</i> , 2016 , 111, 1974-1986	2.9	85

117	A Detergent-Free Approach to Membrane Protein Research: Polymer-Bounded Native Nanodiscs. <i>Biophysical Journal</i> , 2016 , 110, 580a	2.9	
116	Photophysics in single light-harvesting complexes II: from micelle to native nanodisks 2016 ,		3
115	The styrene-maleic acid copolymer: a versatile tool in membrane research. <i>European Biophysics Journal</i> , 2016 , 45, 3-21	1.9	241
114	Activation of the bacterial thermosensor DesK involves a serine zipper dimerization motif that is modulated by bilayer thickness. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 6353-8	11.5	37
113	E. coli MG1655 modulates its phospholipid composition through the cell cycle. <i>FEBS Letters</i> , 2015 , 589, 2726-30	3.8	22
112	Isolation of lipids from biological samples. <i>Molecular Membrane Biology</i> , 2015 , 32, 55-64	3.4	31
111	Molecular model for the solubilization of membranes into nanodisks by styrene maleic Acid copolymers. <i>Biophysical Journal</i> , 2015 , 108, 279-90	2.9	121
110	Bacterial Reaction Centers Purified with Styrene Maleic Acid Copolymer Retain Native Membrane Functional Properties and Display Enhanced Stability. <i>Angewandte Chemie</i> , 2014 , 126, 11997-12001	3.6	15
109	Bacterial reaction centers purified with styrene maleic acid copolymer retain native membrane functional properties and display enhanced stability. <i>Angewandte Chemie - International Edition</i> , 2014 , 53, 11803-7	16.4	107
108	Detergent-free isolation, characterization, and functional reconstitution of a tetrameric K ⁺ channel: the power of native nanodiscs. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 18607-12	11.5	232
107	Lipase activity in lipidomics - a hidden problem?. <i>Molecular Membrane Biology</i> , 2013 , 30, 347-9	3.4	4
106	Biophysical investigation of the membrane-disrupting mechanism of the antimicrobial and amyloid-like peptide dermaseptin S9. <i>PLoS ONE</i> , 2013 , 8, e75528	3.7	30
105	Thermodynamic measurements of bilayer insertion of a single transmembrane helix chaperoned by fluorinated surfactants. <i>Journal of Molecular Biology</i> , 2012 , 416, 328-34	6.5	16
104	How lipid headgroups sense the membrane environment: an application of ³¹ P NMR. <i>Biophysical Journal</i> , 2012 , 103, 1245-53	2.9	15
103	Modeling the membrane environment for membrane proteins. <i>Biophysical Journal</i> , 2011 , 100, 2073-4; author reply 2075	2.9	16
102	Sterols have higher affinity for sphingomyelin than for phosphatidylcholine bilayers even at equal acyl-chain order. <i>Biophysical Journal</i> , 2011 , 100, 2633-41	2.9	65
101	Probing the lipid-protein interface using model transmembrane peptides with a covalently linked acyl chain. <i>Biophysical Journal</i> , 2011 , 101, 1959-67	2.9	7
100	Low pH acts as inhibitor of membrane damage induced by human islet amyloid polypeptide. <i>Journal of the American Chemical Society</i> , 2011 , 133, 15598-604	16.4	61

99	Ruthenium-decorated lipid vesicles: light-induced release of [Ru(terpy)(bpy)(OH ₂)] ²⁺ and thermal back coordination. <i>Journal of the American Chemical Society</i> , 2011 , 133, 252-61	16.4	64
98	Lipid packing drives the segregation of transmembrane helices into disordered lipid domains in model membranes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 1343-8	11.5	193
97	Mechanism and kinetics of peptide partitioning into membranes from all-atom simulations of thermostable peptides. <i>Journal of the American Chemical Society</i> , 2010 , 132, 3452-60	16.4	70
96	Order parameters of a transmembrane helix in a fluid bilayer: case study of a WALP peptide. <i>Biophysical Journal</i> , 2010 , 98, 1864-72	2.9	43
95	Influence of hydrophobic mismatch and amino acid composition on the lateral diffusion of transmembrane peptides. <i>Biophysical Journal</i> , 2010 , 99, 1447-54	2.9	72
94	Self-reproduction of fatty acid vesicles: a combined experimental and simulation study. <i>Biophysical Journal</i> , 2010 , 99, 1520-8	2.9	44
93	The N-terminal fragment of human islet amyloid polypeptide is non-fibrillogenic in the presence of membranes and does not cause leakage of bilayers of physiologically relevant lipid composition. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2010 , 1798, 1805-11	3.8	23
92	Orientation and dynamics of transmembrane peptides: the power of simple models. <i>European Biophysics Journal</i> , 2010 , 39, 609-21	1.9	99
91	The role of the disulfide bond in the interaction of islet amyloid polypeptide with membranes. <i>European Biophysics Journal</i> , 2010 , 39, 1359-64	1.9	16
90	The influence of the acyl chain composition of cardiolipin on the stability of mitochondrial complexes; an unexpected effect of cardiolipin in alpha-ketoglutarate dehydrogenase and prohibitin complexes. <i>Journal of Proteomics</i> , 2010 , 73, 806-14	3.9	24
89	Aggregation of transmembrane peptides studied by spin-label EPR. <i>Journal of Physical Chemistry B</i> , 2009 , 113, 12257-64	3.4	18
88	Impaired processing of human pro-islet amyloid polypeptide is not a causative factor for fibril formation or membrane damage in vitro. <i>Biochemistry</i> , 2009 , 48, 10918-25	3.2	20
87	Lateral diffusion of membrane proteins. <i>Journal of the American Chemical Society</i> , 2009 , 131, 12650-6	16.4	250
86	Peptide Partitioning and Folding into Lipid Bilayers. <i>Journal of Chemical Theory and Computation</i> , 2009 , 5, 2202-5	6.4	16
85	Activation of phospholipase A2 by temporin B: formation of antimicrobial peptide-enzyme amyloid-type cofibrils. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2009 , 1788, 1064-72	3.8	18
84	Tilt and rotation angles of a transmembrane model peptide as studied by fluorescence spectroscopy. <i>Biophysical Journal</i> , 2009 , 97, 2258-66	2.9	39
83	Influence of trifluoroethanol on membrane interfacial anchoring interactions of transmembrane alpha-helical peptides. <i>Biophysical Journal</i> , 2008 , 94, 1315-25	2.9	20
82	Protein self-assembly and lipid binding in the folding of the potassium channel KcsA. <i>Biochemistry</i> , 2008 , 47, 2123-33	3.2	37

81	Is there a preferential interaction between cholesterol and tryptophan residues in membrane proteins?. <i>Biochemistry</i> , 2008 , 47, 2638-49	3.2	25
80	Membrane damage by human islet amyloid polypeptide through fibril growth at the membrane. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 6033-8	11.5	381
79	Recent insights in islet amyloid polypeptide-induced membrane disruption and its role in beta-cell death in type 2 diabetes mellitus. <i>Experimental Diabetes Research</i> , 2008 , 2008, 421287		97
78	On the orientation of a designed transmembrane peptide: toward the right tilt angle?. <i>Journal of the American Chemical Society</i> , 2007 , 129, 15174-81	16.4	92
77	How protein transmembrane segments sense the lipid environment. <i>Biochemistry</i> , 2007 , 46, 1457-65	3.2	134
76	2,2,2-Trifluoroethanol changes the transition kinetics and subunit interactions in the small bacterial mechanosensitive channel MscS. <i>Biophysical Journal</i> , 2007 , 92, 2771-84	2.9	26
75	Phosphatidic acid plays a special role in stabilizing and folding of the tetrameric potassium channel KcsA. <i>FEBS Letters</i> , 2007 , 581, 5715-22	3.8	41
74	Peptides in lipid bilayers: the power of simple models. <i>Current Opinion in Structural Biology</i> , 2006 , 16, 473-9	8.1	130
73	Striated domains: self-organizing ordered assemblies of transmembrane alpha-helical peptides and lipids in bilayers. <i>Biological Chemistry</i> , 2006 , 387, 235-41	4.5	10
72	Transmembrane peptides stabilize inverted cubic phases in a biphasic length-dependent manner: implications for protein-induced membrane fusion. <i>Biophysical Journal</i> , 2006 , 90, 200-11	2.9	39
71	Islet amyloid polypeptide inserts into phospholipid monolayers as monomer. <i>Journal of Molecular Biology</i> , 2006 , 356, 783-9	6.5	157
70	Molecular organization in striated domains induced by transmembrane alpha-helical peptides in dipalmitoyl phosphatidylcholine bilayers. <i>Biochemistry</i> , 2005 , 44, 2-10	3.2	19
69	A synergistic effect between cholesterol and tryptophan-flanked transmembrane helices modulates membrane curvature. <i>Biochemistry</i> , 2005 , 44, 4526-32	3.2	25
68	Influence of flanking residues on tilt and rotation angles of transmembrane peptides in lipid bilayers. A solid-state 2H NMR study. <i>Biochemistry</i> , 2005 , 44, 1004-12	3.2	91
67	A convenient solid phase synthesis of S-palmitoyl transmembrane peptides. <i>Tetrahedron Letters</i> , 2005 , 46, 3341-3345	2	21
66	Detection and identification of stable oligomeric protein complexes in Escherichia coli inner membranes: a proteomics approach. <i>Journal of Biological Chemistry</i> , 2005 , 280, 28742-8	5.4	22
65	Self-association of transmembrane alpha-helices in model membranes: importance of helix orientation and role of hydrophobic mismatch. <i>Journal of Biological Chemistry</i> , 2005 , 280, 39324-31	5.4	106
64	Small alcohols destabilize the KcsA tetramer via their effect on the membrane lateral pressure. <i>Biochemistry</i> , 2004 , 43, 5937-42	3.2	43

63	Transbilayer movement of phospholipids in biogenic membranes. <i>Biochemistry</i> , 2004 , 43, 2673-81	3.2	92
62	Stability of KcsA tetramer depends on membrane lateral pressure. <i>Biochemistry</i> , 2004 , 43, 4240-50	3.2	71
61	Photo-crosslinking analysis of preferential interactions between a transmembrane peptide and matching lipids. <i>Biochemistry</i> , 2004 , 43, 4482-9	3.2	30
60	Strength of integration of transmembrane alpha-helical peptides in lipid bilayers as determined by atomic force spectroscopy. <i>Biochemistry</i> , 2004 , 43, 14987-93	3.2	33
59	Tilt angles of transmembrane model peptides in oriented and non-oriented lipid bilayers as determined by 2H solid-state NMR. <i>Biophysical Journal</i> , 2004 , 86, 3709-21	2.9	162
58	Islet amyloid polypeptide-induced membrane leakage involves uptake of lipids by forming amyloid fibers. <i>FEBS Letters</i> , 2004 , 577, 117-20	3.8	211
57	Nonbilayer lipids affect peripheral and integral membrane proteins via changes in the lateral pressure profile. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2004 , 1666, 275-88	3.8	318
56	Phospholipid flop induced by transmembrane peptides in model membranes is modulated by lipid composition. <i>Biochemistry</i> , 2003 , 42, 231-7	3.2	109
55	Interfacial anchor properties of tryptophan residues in transmembrane peptides can dominate over hydrophobic matching effects in peptide-lipid interactions. <i>Biochemistry</i> , 2003 , 42, 5341-8	3.2	233
54	Interaction of the K ⁺ channel KcsA with membrane phospholipids as studied by ESI mass spectrometry. <i>FEBS Letters</i> , 2003 , 541, 28-32	3.8	46
53	Snorkeling of lysine side chains in transmembrane helices: how easy can it get?. <i>FEBS Letters</i> , 2003 , 544, 69-73	3.8	163
52	Sphingomyelin is much more effective than saturated phosphatidylcholine in excluding unsaturated phosphatidylcholine from domains formed with cholesterol. <i>FEBS Letters</i> , 2003 , 547, 101-6	3.8	86
51	Synthetic peptides as models for intrinsic membrane proteins. <i>FEBS Letters</i> , 2003 , 555, 134-8	3.8	127
50	Hydrophobic mismatch between helices and lipid bilayers. <i>Biophysical Journal</i> , 2003 , 84, 379-85	2.9	122
49	Protein-lipid interactions studied with designed transmembrane peptides: role of hydrophobic matching and interfacial anchoring. <i>Molecular Membrane Biology</i> , 2003 , 20, 271-84	3.4	245
48	Membrane interaction of the glycosyltransferase MurG: a special role for cardiolipin. <i>Journal of Bacteriology</i> , 2003 , 185, 3773-9	3.5	65
47	Domain formation in phosphatidylcholine bilayers containing transmembrane peptides: specific effects of flanking residues. <i>Biochemistry</i> , 2002 , 41, 2814-24	3.2	78
46	Lipid dependence of membrane anchoring properties and snorkeling behavior of aromatic and charged residues in transmembrane peptides. <i>Biochemistry</i> , 2002 , 41, 7190-8	3.2	99

45	The effects of hydrophobic mismatch between phosphatidylcholine bilayers and transmembrane alpha-helical peptides depend on the nature of interfacially exposed aromatic and charged residues. <i>Biochemistry</i> , 2002 , 41, 8396-404	3.2	89
44	Factors affecting gas-phase deuterium scrambling in peptide ions and their implications for protein structure determination. <i>Journal of the American Chemical Society</i> , 2002 , 124, 11191-8	16.4	99
43	Importance of hydrophobic matching for spontaneous insertion of a single-spanning membrane protein. <i>Biochemistry</i> , 2002 , 41, 4946-52	3.2	46
42	Components required for membrane assembly of newly synthesized K ⁺ channel KcsA. <i>FEBS Letters</i> , 2002 , 511, 51-8	3.8	24
41	Influence of hydrophobic mismatch and palmitoylation on the association of transmembrane alpha-helical peptides with detergent-resistant membranes. <i>FEBS Letters</i> , 2002 , 523, 79-84	3.8	48
40	Influence of lipids on membrane assembly and stability of the potassium channel KcsA. <i>FEBS Letters</i> , 2002 , 525, 33-8	3.8	71
39	Geometry and intrinsic tilt of a tryptophan-anchored transmembrane alpha-helix determined by (2)H NMR. <i>Biophysical Journal</i> , 2002 , 83, 1479-88	2.9	156
38	Characterization of the thermotropic behavior and lateral organization of lipid-peptide mixtures by a combined experimental and theoretical approach: effects of hydrophobic mismatch and role of flanking residues. <i>Biophysical Journal</i> , 2002 , 82, 1405-17	2.9	33
37	Hydrophobic Matching Mechanism Investigated by Molecular Dynamics Simulations. <i>Langmuir</i> , 2002 , 18, 1340-1351	4	79
36	Sensitivity of single membrane-spanning alpha-helical peptides to hydrophobic mismatch with a lipid bilayer: effects on backbone structure, orientation, and extent of membrane incorporation. <i>Biochemistry</i> , 2001 , 40, 5000-10	3.2	162
35	Anionic lipids stimulate Sec-independent insertion of a membrane protein lacking charged amino acid side chains. <i>EMBO Reports</i> , 2001 , 2, 403-8	6.5	38
34	Membrane-spanning peptides induce phospholipid flop: a model for phospholipid translocation across the inner membrane of E. coli. <i>Biochemistry</i> , 2001 , 40, 10500-6	3.2	76
33	Effect of nonbilayer lipids on membrane binding and insertion of the catalytic domain of leader peptidase. <i>Biochemistry</i> , 2001 , 40, 9677-84	3.2	44
32	Efficient membrane assembly of the KcsA potassium channel in Escherichia coli requires the protonmotive force. <i>EMBO Reports</i> , 2000 , 1, 340-6	6.5	31
31	The effect of peptide/lipid hydrophobic mismatch on the phase behavior of model membranes mimicking the lipid composition in Escherichia coli membranes. <i>Biophysical Journal</i> , 2000 , 78, 2475-85	2.9	53
30	Analysis of the role of interfacial tryptophan residues in controlling the topology of membrane proteins. <i>Biochemistry</i> , 2000 , 39, 6521-8	3.2	117
29	Visualization of highly ordered striated domains induced by transmembrane peptides in supported phosphatidylcholine bilayers. <i>Biochemistry</i> , 2000 , 39, 5852-8	3.2	108
28	Tryptophan-anchored transmembrane peptides promote formation of nonlamellar phases in phosphatidylethanolamine model membranes in a mismatch-dependent manner. <i>Biochemistry</i> , 2000 , 39, 3124-33	3.2	54

27	Different membrane anchoring positions of tryptophan and lysine in synthetic transmembrane alpha-helical peptides. <i>Journal of Biological Chemistry</i> , 1999 , 274, 20839-46	5.4	273
26	Membrane fusion and the lamellar-to-inverted-hexagonal phase transition in cardiolipin vesicle systems induced by divalent cations. <i>Biophysical Journal</i> , 1999 , 77, 2003-14	2.9	110
25	Peptide influences on lipids. <i>Novartis Foundation Symposium</i> , 1999 , 225, 170-83; discussion 183-7		1
24	Influence of lipid/peptide hydrophobic mismatch on the thickness of diacylphosphatidylcholine bilayers. A 2H NMR and ESR study using designed transmembrane alpha-helical peptides and gramicidin A. <i>Biochemistry</i> , 1998 , 37, 9333-45	3.2	242
23	Molecular ordering of interfacially localized tryptophan analogs in ester- and ether-lipid bilayers studied by 2H-NMR. <i>Biophysical Journal</i> , 1998 , 75, 1365-71	2.9	102
22	Chapter 13 Phospholipid Structure and Escherichia Coli Membranes. <i>Current Topics in Membranes</i> , 1997 , 477-515	2.2	13
21	Conformation of the acylation site of palmitoylgramicidin in lipid bilayers of dimyristoylphosphatidylcholine. <i>Biochemistry</i> , 1996 , 35, 3641-8	3.2	24
20	Induction of nonbilayer structures in diacylphosphatidylcholine model membranes by transmembrane alpha-helical peptides: importance of hydrophobic mismatch and proposed role of tryptophans. <i>Biochemistry</i> , 1996 , 35, 1037-45	3.2	259
19	PhoE signal peptide inserts into micelles as a dynamic helix-break-helix structure, which is modulated by the environment. A two-dimensional 1H NMR study. <i>Biochemistry</i> , 1995 , 34, 11617-24	3.2	71
18	Palmitoylation-induced conformational changes of specific side chains in the gramicidin transmembrane channel. <i>Biochemistry</i> , 1995 , 34, 9299-306	3.2	33
17	Effect of divalent cations on lipid organization of cardiolipin isolated from Escherichia coli strain AH930. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1994 , 1189, 225-32	3.8	58
16	Analysis of circular dichroism spectra of oriented protein-lipid complexes: toward a general application. <i>Biochemistry</i> , 1994 , 33, 14521-8	3.2	94
15	Orientation of the alpha-helices of apocytochrome c and derived fragments at membrane interfaces, as studied by circular dichroism. <i>Biochemistry</i> , 1994 , 33, 14529-35	3.2	12
14	A water-lipid interface induces a highly dynamic folded state in apocytochrome c and cytochrome c, which may represent a common folding intermediate. <i>Biochemistry</i> , 1992 , 31, 1636-43	3.2	97
13	Anionic phospholipids are essential for alpha-helix formation of the signal peptide of prePhoE upon interaction with phospholipid vesicles. <i>Biochemistry</i> , 1992 , 31, 1672-7	3.2	88
12	Acidic interaction of the colicin A pore-forming domain with model membranes of Escherichia coli lipids results in a large perturbation of acyl chain order and stabilization of the bilayer. <i>Biochemistry</i> , 1992 , 31, 11089-94	3.2	20
11	Orientation of the valine-1 side chain of the gramicidin transmembrane channel and implications for channel functioning. A 2H NMR study. <i>Biochemistry</i> , 1992 , 31, 11283-90	3.2	63
10	Effects of temperature variation and phenethyl alcohol addition on acyl chain order and lipid organization in Escherichia coli derived membrane systems. A 2H- and 31P-NMR study. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1992 , 1105, 253-62	3.8	45

9	Gramicidin and gramicidin-lipid interactions. <i>BBA - Biomembranes</i> , 1992 , 1113, 391-425		190
8	The membrane interaction of amphiphilic model peptides affects phosphatidylserine headgroup and acyl chain order and dynamics. Application of the "phospholipid headgroup electrometer" concept to phosphatidylserine. <i>Biochemistry</i> , 1991 , 30, 1155-62	3.2	19
7	The mitochondrial precursor protein apocytochrome c strongly influences the order of the headgroup and acyl chains of phosphatidylserine dispersions. A 2H and 31P NMR study. <i>Biochemistry</i> , 1990 , 29, 2312-21	3.2	18
6	Conformation of gramicidin in relation to its ability to form bilayers with lysophosphatidylcholine. <i>Biochemistry</i> , 1988 , 27, 7295-301	3.2	30
5	The membrane as an environment of minimal interconversion. A circular dichroism study on the solvent dependence of the conformational behavior of gramicidin in diacylphosphatidylcholine model membranes. <i>Biochemistry</i> , 1988 , 27, 4848-55	3.2	142
4	Gramicidin-induced hexagonal HII phase formation in negatively charged phospholipids and the effect of N- and C-terminal modification of gramicidin on its interaction with zwitterionic phospholipids. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1986 , 857, 13-27	3.8	33
3	Thermodynamic, motional, and structural aspects of gramicidin-induced hexagonal HII phase formation in phosphatidylethanolamine. <i>Biochemistry</i> , 1985 , 24, 7881-90	3.2	64
2	Importance of hydration for gramicidin-induced hexagonal HII phase formation in dioleoylphosphatidylcholine model membranes. <i>Biochemistry</i> , 1985 , 24, 7890-8	3.2	49
1	Membrane-catalyzed IAPP aggregation is dominated by secondary nucleation		1