Naser Elkum

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1446774/publications.pdf

Version: 2024-02-01

37	1,935	20	36
papers	citations	h-index	g-index
38	38	38	3352
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Genome-wide landscape establishes novel association signals for metabolic traits in the Arab population. Human Genetics, 2021, 140, 505-528.	3.8	13
2	Adult Diabetes and Prediabetes Prevalence in Kuwait: Data from the Cross-Sectional Kuwait Diabetes Epidemiology Program. Journal of Clinical Medicine, 2020, 9, 3420.	2.4	17
3	CSF total and oligomeric α-Synuclein along with TNF-α as risk biomarkers for Parkinson's disease: a study in LRRK2 mutation carriers. Translational Neurodegeneration, 2020, 9, 15.	8.0	32
4	PR3 levels are impaired in plasma and PBMCs from Arabs with cardiovascular diseases. PLoS ONE, 2020, 15, e0227606.	2.5	2
5	Genome-wide association study identifies novel risk variants from RPS6KA1, CADPS, VARS, and DHX58 for fasting plasma glucose in Arab population. Scientific Reports, 2020, 10, 152.	3.3	16
6	A teaching intervention increases the performance of handheld ultrasound devices for assessment of left ventricular ejection fraction. Heart Views, 2019, 20, 133.	0.2	1
7	DUSP1 Is a Potential Marker of Chronic Inflammation in Arabs with Cardiovascular Diseases. Disease Markers, 2018, 2018, 1-10.	1.3	8
8	Investigation of genetic variation and lifestyle determinants in vitamin D levels in Arab individuals. Journal of Translational Medicine, 2018, 16, 20.	4.4	9
9	Genome-wide association study identifies novel recessive genetic variants for high TGs in an Arab population. Journal of Lipid Research, 2018, 59, 1951-1966.	4.2	24
10	Genetic risk variants for metabolic traits in Arab populations. Scientific Reports, 2017, 7, 40988.	3.3	26
11	The <i>TCN2</i> variant of rs9606756 [lle23Val] acts as risk loci for obesityâ€related traits and mediates by interacting with Apoâ€A1. Obesity, 2017, 25, 1098-1108.	3.0	16
12	ANGPTL8/Betatrophin R59W variant is associated with higher glucose level in non-diabetic Arabs living in Kuwaits. Lipids in Health and Disease, 2016, 15, 26.	3.0	16
13	Physical exercise alleviates ER stress in obese humans through reduction in the expression and release of GRP78 chaperone. Metabolism: Clinical and Experimental, 2016, 65, 1409-1420.	3.4	45
14	Circulating angiopoietin-like protein 8 (betatrophin) association with HsCRP and metabolic syndrome. Cardiovascular Diabetology, 2016, 15, 25.	6.8	67
15	Model Comparison for Breast Cancer Prognosis Based on Clinical Data. PLoS ONE, 2016, 11, e0146413.	2.5	34
16	Circulating ANGPTL8/Betatrophin Is Increased in Obesity and Reduced after Exercise Training. PLoS ONE, 2016, 11, e0147367.	2.5	84
17	Value of routine blood cultures in febrile children presenting to the Emergency Department. Journal of Emergency Medicine, Trauma and Acute Care, 2016, 2016, .	0.1	0
18	Higher plasma betatrophin/ANGPTL8 level in Type 2 Diabetes subjects does not correlate with blood glucose or insulin resistance. Scientific Reports, 2015, 5, 10949.	3.3	108

#	Article	IF	Citations
19	Lack of associations between betatrophin/ANGPTL8 level and C-peptide in type 2 diabetic subjects. Cardiovascular Diabetology, 2015, 14, 112.	6.8	32
20	Gender-Specific Association of Oxidative Stress and Inflammation with Cardiovascular Risk Factors in Arab Population. Mediators of Inflammation, 2015, 2015, 1-11.	3.0	19
21	High adiponectin levels in lean Arab women compared to Asian women. Biomarker Research, 2015, 3, 7.	6.8	3
22	Obesity susceptibility loci in Qataris, a highly consanguineous Arabian population. Journal of Translational Medicine, 2015, 13, 119.	4.4	21
23	MAP kinase phosphatase DUSP1 is overexpressed in obese humans and modulated by physical exercise. American Journal of Physiology - Endocrinology and Metabolism, 2015, 308, E71-E83.	3.5	54
24	Vitamin D Insufficiency in Arabs and South Asians Positively Associates with Polymorphisms in GC and CYP2R1 Genes. PLoS ONE, 2014, 9, e113102.	2.5	68
25	Physical Exercise Reduces the Expression of RANTES and Its CCR5 Receptor in the Adipose Tissue of Obese Humans. Mediators of Inflammation, 2014, 2014, 1-13.	3.0	41
26	Gender Differences in Ghrelin Association with Cardiometabolic Risk Factors in Arab Population. International Journal of Endocrinology, 2014, 2014, 1-8.	1.5	13
27	Obesity is a significant risk factor for breast cancer in Arab women. BMC Cancer, 2014, 14, 788.	2.6	53
28	Evaluation of the usefulness of a D dimer test in combination with clinical pretest probability score in the prediction and exclusion of Venous Thromboembolism by medical residents. Thrombosis Journal, 2014, 12, 28.	2.1	17
29	Immunohistochemical profiling of the heat shock response in obese non-diabetic subjects revealed impaired expression of heat shock proteins in the adipose tissue. Lipids in Health and Disease, 2014, 13, 106.	3.0	43
30	Barriers in participant recruitment of diverse ethnicities in the state of Kuwait. International Journal for Equity in Health, 2013, 12, 93.	3.5	10
31	DNAJB3/HSP-40 Cochaperone Is Downregulated in Obese Humans and Is Restored by Physical Exercise. PLoS ONE, 2013, 8, e69217.	2.5	58
32	Proteomics Analysis of Human Obesity Reveals the Epigenetic Factor HDAC4 as a Potential Target for Obesity. PLoS ONE, 2013, 8, e75342.	2.5	75
33	The p53 codon 72 polymorphism is associated with risk and early onset of breast cancer among Saudi women. Oncology Letters, 2012, 3, 875-878.	1.8	18
34	FOXP3+ Tregs and B7-H1+/PD-1+T lymphocytes co-infiltrate the tumor tissues of high-risk breast cancer patients: Implication for immunotherapy. BMC Cancer, 2008, 8, 57.	2.6	178
35	Being 40 or younger is an independent risk factor for relapse in operable breast cancer patients: The Saudi Arabia experience. BMC Cancer, 2007, 7, 222.	2.6	76
36	Expression of B7-H1 in breast cancer patients is strongly associated with high proliferative Ki-67-expressing tumor cells. International Journal of Cancer, 2007, 121, 751-758.	5.1	132

#	ARTICLE	IF	CITATIONS
37	The B7-H1 (PD-L1) T Lymphocyte-Inhibitory Molecule Is Expressed in Breast Cancer Patients with Infiltrating Ductal Carcinoma: Correlation with Important High-Risk Prognostic Factors. Neoplasia, 2006, 8, 190-198.	5.3	505