

Javier F Caceres

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1449356/publications.pdf>

Version: 2024-02-01

81
papers

12,354
citations

41258

49
h-index

60497

81
g-index

93
all docs

93
docs citations

93
times ranked

13836
citing authors

#	ARTICLE	IF	CITATIONS
1	RNA splicing is a key mediator of tumour cell plasticity and a therapeutic vulnerability in colorectal cancer. <i>Nature Communications</i> , 2022, 13, 2791.	5.8	11
2	Nucleo-cytoplasmic shuttling of splicing factor SRSF1 is required for development and cilia function. <i>ELife</i> , 2021, 10, .	2.8	25
3	Identification of a localized nonsense-mediated decay pathway at the endoplasmic reticulum. <i>Genes and Development</i> , 2020, 34, 1075-1088.	2.7	37
4	The complex genetic landscape of familial MDS and AML reveals pathogenic germline variants. <i>Nature Communications</i> , 2020, 11, 1044.	5.8	81
5	Regulation of RUVBL1-RUVBL2 AAA-ATPases by the nonsense-mediated mRNA decay factor DHX34, as evidenced by Cryo-EM. <i>ELife</i> , 2020, 9, .	2.8	9
6	A slow transcription rate causes embryonic lethality and perturbs kinetic coupling of neuronal genes. <i>EMBO Journal</i> , 2019, 38, .	3.5	46
7	Rapid Depletion of DIS3, EXOSC10, or XRN2 Reveals the Immediate Impact of Exoribonucleolysis on Nuclear RNA Metabolism and Transcriptional Control. <i>Cell Reports</i> , 2019, 26, 2779-2791.e5.	2.9	61
8	Post-transcriptional control of miRNA biogenesis. <i>Rna</i> , 2019, 25, 1-16.	1.6	390
9	The Secretion of miR-200s by a PKC ζ /ADAR2 Signaling Axis Promotes Liver Metastasis in Colorectal Cancer. <i>Cell Reports</i> , 2018, 23, 1178-1191.	2.9	53
10	Structural basis for terminal loop recognition and stimulation of pri-miRNA-18a processing by hnRNP A1. <i>Nature Communications</i> , 2018, 9, 2479.	5.8	80
11	Genetic variation and RNA structure regulate microRNA biogenesis. <i>Nature Communications</i> , 2017, 8, 15114.	5.8	67
12	The RNA-binding landscape of RBM10 and its role in alternative splicing regulation in models of mouse early development. <i>RNA Biology</i> , 2017, 14, 45-57.	1.5	41
13	The RNA-binding profile of Acinus, a peripheral component of the exon junction complex, reveals its role in splicing regulation. <i>Rna</i> , 2016, 22, 1411-1426.	1.6	33
14	Mechanism and regulation of the nonsense-mediated decay pathway. <i>Nucleic Acids Research</i> , 2016, 44, 1483-1495.	6.5	415
15	The RNA helicase DHX34 functions as a scaffold for SMG1-mediated UPF1 phosphorylation. <i>Nature Communications</i> , 2016, 7, 10585.	5.8	39
16	DGCR8 Acts as an Adaptor for the Exosome Complex to Degrade Double-Stranded Structured RNAs. <i>Molecular Cell</i> , 2015, 60, 873-885.	4.5	68
17	Identification and characterization of novel factors that act in the nonsense-mediated mRNA decay pathway in nematodes, flies and mammals. <i>EMBO Reports</i> , 2015, 16, 71-78.	2.0	33
18	The RNA Helicase DHX34 Activates NMD by Promoting a Transition from the Surveillance to the Decay-Inducing Complex. <i>Cell Reports</i> , 2014, 8, 1845-1856.	2.9	65

#	ARTICLE	IF	CITATIONS
19	Control of mammalian retrotransposons by cellular RNA processing activities. <i>Mobile Genetic Elements</i> , 2014, 4, e28439.	1.8	31
20	The translational landscape of the splicing factor SRSF1 and its role in mitosis. <i>ELife</i> , 2014, 3, e02028.	2.8	96
21	The Microprocessor controls the activity of mammalian retrotransposons. <i>Nature Structural and Molecular Biology</i> , 2013, 20, 1173-1181.	3.6	105
22	Drosha Regulates Gene Expression Independently of RNA Cleavage Function. <i>Cell Reports</i> , 2013, 5, 1499-1510.	2.9	60
23	Cellular functions of the microprocessor. <i>Biochemical Society Transactions</i> , 2013, 41, 838-843.	1.6	40
24	DHX34 and NBAS form part of an autoregulatory NMD circuit that regulates endogenous RNA targets in human cells, zebrafish and <i>Caenorhabditis elegans</i> . <i>Nucleic Acids Research</i> , 2013, 41, 8319-8331.	6.5	80
25	Tissue-specific control of brain-enriched miR-7 biogenesis. <i>Genes and Development</i> , 2013, 27, 24-38.	2.7	131
26	DGCR8 HITS-CLIP reveals novel functions for the Microprocessor. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 760-766.	3.6	200
27	Dhx34 and Nbas function in the NMD pathway and are required for embryonic development in zebrafish. <i>Nucleic Acids Research</i> , 2011, 39, 3686-3694.	6.5	58
28	An Aptamer Targeting the Apical Loop Domain Modulates pri-miRNA Processing. <i>Angewandte Chemie - International Edition</i> , 2010, 49, 4674-4677.	7.2	49
29	Antagonistic role of hnRNP A1 and KSRP in the regulation of let-7a biogenesis. <i>Nature Structural and Molecular Biology</i> , 2010, 17, 1011-1018.	3.6	241
30	RNase-assisted RNA chromatography. <i>Rna</i> , 2010, 16, 1673-1678.	1.6	44
31	Stimulation of pri-miR-18a Processing by hnRNP A1. <i>Advances in Experimental Medicine and Biology</i> , 2010, 700, 28-35.	0.8	38
32	Cellular stress and RNA splicing. <i>Trends in Biochemical Sciences</i> , 2009, 34, 146-153.	3.7	181
33	Editing independent effects of ADARs on the miRNA/siRNA pathways. <i>EMBO Journal</i> , 2009, 28, 3145-3156.	3.5	161
34	The SR protein family of splicing factors: master regulators of gene expression. <i>Biochemical Journal</i> , 2009, 417, 15-27.	1.7	934
35	Hormonal Regulation of MicroRNA Biogenesis. <i>Molecular Cell</i> , 2009, 36, 172-173.	4.5	28
36	The Splicing Factor SF2/ASF Regulates Translation Initiation by Enhancing Phosphorylation of 4E-BP1. <i>Molecular Cell</i> , 2008, 30, 179-189.	4.5	233

#	ARTICLE	IF	CITATIONS
37	Posttranscriptional Regulation of miRNAs Harboring Conserved Terminal Loops. <i>Molecular Cell</i> , 2008, 32, 383-393.	4.5	316
38	Nonsense-Mediated mRNA Decay in <i>Caenorhabditis elegans</i> . <i>Methods in Enzymology</i> , 2008, 449, 149-164.	0.4	4
39	Spatial mapping of splicing factor complexes involved in exon and intron definition. <i>Journal of Cell Biology</i> , 2008, 181, 921-934.	2.3	53
40	Identification of Nuclear and Cytoplasmic mRNA Targets for the Shuttling Protein SF2/ASF. <i>PLoS ONE</i> , 2008, 3, e3369.	1.1	98
41	Mechanistic insights and identification of two novel factors in the <i>C. elegans</i> NMD pathway. <i>Genes and Development</i> , 2007, 21, 1075-1085.	2.7	140
42	Division of Labor: Minor Splicing in the Cytoplasm. <i>Cell</i> , 2007, 131, 645-647.	13.5	8
43	Stressful Splicing. <i>Molecular Cell</i> , 2007, 28, 180-181.	4.5	8
44	Identification and characterization of RED120: A conserved PWI domain protein with links to splicing and 3' end formation. <i>FEBS Letters</i> , 2007, 581, 3087-3097.	1.3	20
45	The multifunctional RNA-binding protein hnRNP A1 is required for processing of miR-18a. <i>Nature Structural and Molecular Biology</i> , 2007, 14, 591-596.	3.6	505
46	Cdk1 is sufficient to drive the mammalian cell cycle. <i>Nature</i> , 2007, 448, 811-815.	13.7	888
47	hnRNP A1 Relocalization to the Stress Granules Reflects a Role in the Stress Response. <i>Molecular and Cellular Biology</i> , 2006, 26, 5744-5758.	1.1	281
48	Multiple roles of arginine/serine-rich splicing factors in RNA processing. <i>Biochemical Society Transactions</i> , 2005, 33, 443-446.	1.6	140
49	Concerted regulation of nuclear and cytoplasmic activities of SR proteins by AKT. <i>Nature Structural and Molecular Biology</i> , 2005, 12, 1037-1044.	3.6	211
50	Proteomic Analysis of SRm160-containing Complexes Reveals a Conserved Association with Cohesin. <i>Journal of Biological Chemistry</i> , 2005, 280, 42227-42236.	1.6	28
51	Regulation of heterogenous nuclear ribonucleoprotein A1 transport by phosphorylation in cells stressed by osmotic shock. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 3605-3610.	3.3	144
52	Reversible phosphorylation differentially affects nuclear and cytoplasmic functions of splicing factor 2/alternative splicing factor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 15042-15047.	3.3	113
53	A Novel SR-Related Protein Is Required for the Second Step of Pre-mRNA Splicing. <i>Molecular and Cellular Biology</i> , 2005, 25, 2969-2980.	1.1	55
54	A rapid and efficient protocol to purify biologically active recombinant proteins from mammalian cells. <i>Protein Expression and Purification</i> , 2005, 42, 54-58.	0.6	38

#	ARTICLE	IF	CITATIONS
55	A novel role for shuttling SR proteins in mRNA translation. <i>Genes and Development</i> , 2004, 18, 755-768.	2.7	323
56	Pre-mRNA splicing: life at the centre of the central dogma. <i>Journal of Cell Science</i> , 2004, 117, 6261-6263.	1.2	26
57	The Ref/Aly proteins are dispensable for mRNA export and development in <i>Caenorhabditis elegans</i> . <i>Rna</i> , 2003, 9, 881-891.	1.6	92
58	Fibrillarin Is Essential for Early Development and Required for Accumulation of an Intron-Encoded Small Nucleolar RNA in the Mouse. <i>Molecular and Cellular Biology</i> , 2003, 23, 8519-8527.	1.1	91
59	An Evolutionarily Conserved Role for SRm160 in 3'â€²-End Processing That Functions Independently of Exon Junction Complex Formation. <i>Journal of Biological Chemistry</i> , 2003, 278, 44153-44160.	1.6	33
60	The Wilms' tumour protein (WT1) shuttles between nucleus and cytoplasm and is present in functional polysomes. <i>Human Molecular Genetics</i> , 2003, 13, 463-471.	1.4	130
61	Nuclear Export and Retention Signals in the RS Domain of SR Proteins. <i>Molecular and Cellular Biology</i> , 2002, 22, 6871-6882.	1.1	149
62	Alternative splicing: multiple control mechanisms and involvement in human disease. <i>Trends in Genetics</i> , 2002, 18, 186-193.	2.9	590
63	Multiple interactions between SRm160 and SR family proteins in enhancer-dependent splicing and development of <i>C. elegans</i> . <i>Current Biology</i> , 2001, 11, 1923-1933.	1.8	38
64	Distinctive Features of <i>Drosophila</i> Alternative Splicing Factor RS Domain: Implication for Specific Phosphorylation, Shuttling, and Splicing Activation. <i>Molecular and Cellular Biology</i> , 2001, 21, 1345-1359.	1.1	35
65	Serine-Arginine (SR) Protein-like Factors That Antagonize Authentic SR Proteins and Regulate Alternative Splicing. <i>Journal of Biological Chemistry</i> , 2001, 276, 48908-48914.	1.6	76
66	Large-scale identification of mammalian proteins localized to nuclear sub-compartments. <i>Human Molecular Genetics</i> , 2001, 10, 1995-2011.	1.4	108
67	Functional characterization of SR and SR-related genes in <i>Caenorhabditis elegans</i> . <i>EMBO Journal</i> , 2000, 19, 1625-1637.	3.5	142
68	Role of SR protein modular domains in alternative splicing specificity in vivo. <i>Nucleic Acids Research</i> , 2000, 28, 4822-4831.	6.5	41
69	The Mkk3/6-p38â€² Signaling Cascade Alters the Subcellular Distribution of Hnrnp A1 and Modulates Alternative Splicing Regulation. <i>Journal of Cell Biology</i> , 2000, 149, 307-316.	2.3	309
70	Selection of Alternative 5'â€² Splice Sites: Role of U1 snRNP and Models for the Antagonistic Effects of SF2/ASF and hnRNP A1. <i>Molecular and Cellular Biology</i> , 2000, 20, 8303-8318.	1.1	171
71	RNA Processing Marches on. <i>Cell</i> , 2000, 103, 703-709.	13.5	19
72	Coupling of Transcription with Alternative Splicing. <i>Molecular Cell</i> , 1999, 4, 251-258.	4.5	274

#	ARTICLE	IF	CITATIONS
73	A specific subset of SR proteins shuttles continuously between the nucleus and the cytoplasm. <i>Genes and Development</i> , 1998, 12, 55-66.	2.7	424
74	Serine Phosphorylation of SR Proteins Is Required for Their Recruitment to Sites of Transcription In Vivo. <i>Journal of Cell Biology</i> , 1998, 143, 297-307.	2.3	236
75	Role of the Modular Domains of SR Proteins in Subnuclear Localization and Alternative Splicing Specificity. <i>Journal of Cell Biology</i> , 1997, 138, 225-238.	2.3	360
76	The dynamics of a pre-mRNA splicing factor in living cells. <i>Nature</i> , 1997, 387, 523-527.	13.7	563
77	SRPK1 and Clk/Sty Protein Kinases Show Distinct Substrate Specificities for Serine/Arginine-rich Splicing Factors. <i>Journal of Biological Chemistry</i> , 1996, 271, 24569-24575.	1.6	172
78	Regulation of alternative splicing in vivo by overexpression of antagonistic splicing factors. <i>Science</i> , 1994, 265, 1706-1709.	6.0	594
79	Control of mouse U1a and U1b snRNA gene expression by differential transcription. <i>Nucleic Acids Research</i> , 1992, 20, 4247-4254.	6.5	14
80	Requirement of DNA topoisomerases for in vitro chromatin assembly by 3T6 mouse cell extracts. <i>FEBS Journal</i> , 1989, 181, 531-537.	0.2	5
81	A dual role for the RNA helicase DHX34 in NMD and pre-mRNA splicing and its function in hematopoietic differentiation. <i>Rna</i> , 0, , rna.079277.122.	1.6	4