

Marina Morigi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1451323/marina-morigi-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

91
papers

8,064
citations

47
h-index

89
g-index

91
ext. papers

8,803
ext. citations

7.7
avg, IF

5.48
L-index

#	Paper	IF	Citations
91	SARS-CoV-2 Spike Protein 1 Activates Microvascular Endothelial Cells and Complement System Leading to Platelet Aggregation.. <i>Frontiers in Immunology</i> , 2022 , 13, 827146	8.4	7
90	Shiga Toxin 2 Triggers C3a-Dependent Glomerular and Tubular Injury through Mitochondrial Dysfunction in Hemolytic Uremic Syndrome. <i>Cells</i> , 2022 , 11, 1755	7.9	1
89	Post-translational modifications by SIRT3 de-2-hydroxyisobutyrylase activity regulate glycolysis and enable nephrogenesis. <i>Scientific Reports</i> , 2021 , 11, 23580	4.9	1
88	C3a receptor blockade protects podocytes from injury in diabetic nephropathy. <i>JCI Insight</i> , 2020 , 5,	9.9	17
87	Protective Effects of Human Nonrenal and Renal Stromal Cells and Their Conditioned Media in a Rat Model of Chronic Kidney Disease. <i>Cell Transplantation</i> , 2020 , 29, 963689720965467	4	
86	Stem Cell Therapies in Kidney Diseases: Progress and Challenges. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	36
85	Complement Activation Contributes to the Pathophysiology of Shiga Toxin-Associated Hemolytic Uremic Syndrome. <i>Microorganisms</i> , 2019 , 7,	4.9	12
84	Shiga toxin triggers endothelial and podocyte injury: the role of complement activation. <i>Pediatric Nephrology</i> , 2019 , 34, 379-388	3.2	20
83	Sirtuins in Renal Health and Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 1799-1809	11.8	125
82	Therapeutic potential of stromal cells of non-renal or renal origin in experimental chronic kidney disease. <i>Stem Cell Research and Therapy</i> , 2018 , 9, 220	8.3	19
81	SGLT2 inhibitor dapagliflozin limits podocyte damage in proteinuric nondiabetic nephropathy. <i>JCI Insight</i> , 2018 , 3,	9.9	57
80	A Novel Method for Isolation of Pluripotent Stem Cells from Human Umbilical Cord Blood. <i>Stem Cells and Development</i> , 2017 , 26, 1258-1269	4.4	24
79	Human mesenchymal stromal cells transplanted into mice stimulate renal tubular cells and enhance mitochondrial function. <i>Nature Communications</i> , 2017 , 8, 983	17.4	85
78	Mesenchymal Stromal Cells for Acute Renal Injury 2017 , 1085-1095		
77	Mesenchymal Stem Cells in Kidney Repair. <i>Methods in Molecular Biology</i> , 2016 , 1416, 89-107	1.4	39
76	Functional Human Podocytes Generated in Organoids from Amniotic Fluid Stem Cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2016 , 27, 1400-11	12.7	44
75	Mitochondrial Sirtuin 3 and Renal Diseases. <i>Nephron</i> , 2016 , 134, 14-9	3.3	44

74	A previously unrecognized role of C3a in proteinuric progressive nephropathy. <i>Scientific Reports</i> , 2016 , 6, 28445	4.9	18
73	Renal progenitors derived from human iPSCs engraft and restore function in a mouse model of acute kidney injury. <i>Scientific Reports</i> , 2015 , 5, 8826	4.9	72
72	Mitochondrial-dependent Autoimmunity in Membranous Nephropathy of IgG4-related Disease. <i>EBioMedicine</i> , 2015 , 2, 456-66	8.8	17
71	Renal primordia activate kidney regenerative events in a rat model of progressive renal disease. <i>PLoS ONE</i> , 2015 , 10, e0120235	3.7	14
70	Sirtuin3 Dysfunction Is the Key Determinant of Skeletal Muscle Insulin Resistance by Angiotensin II. <i>PLoS ONE</i> , 2015 , 10, e0127172	3.7	13
69	Sirtuin 3-dependent mitochondrial dynamic improvements protect against acute kidney injury. <i>Journal of Clinical Investigation</i> , 2015 , 125, 715-26	15.9	244
68	Effects of MCP-1 inhibition by bindarit therapy in a rat model of polycystic kidney disease. <i>Nephron</i> , 2015 , 129, 52-61	3.3	35
67	Direct reprogramming of human bone marrow stromal cells into functional renal cells using cell-free extracts. <i>Stem Cell Reports</i> , 2015 , 4, 685-98	8	25
66	Recellularization of well-preserved acellular kidney scaffold using embryonic stem cells. <i>Tissue Engineering - Part A</i> , 2014 , 20, 1486-98	3.9	134
65	Errestin-1 drives endothelin-1-mediated podocyte activation and sustains renal injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2014 , 25, 523-33	12.7	54
64	Shiga toxin promotes podocyte injury in experimental hemolytic uremic syndrome via activation of the alternative pathway of complement. <i>Journal of the American Society of Nephrology: JASN</i> , 2014 , 25, 1786-98	12.7	39
63	Cell therapy for kidney injury: different options and mechanisms--mesenchymal and amniotic fluid stem cells. <i>Nephron Experimental Nephrology</i> , 2014 , 126, 59		38
62	Angiotensin II contributes to diabetic renal dysfunction in rodents and humans via Notch1/Snail pathway. <i>American Journal of Pathology</i> , 2013 , 183, 119-30	5.8	33
61	Transfer of growth factor receptor mRNA via exosomes unravels the regenerative effect of mesenchymal stem cells. <i>Stem Cells and Development</i> , 2013 , 22, 772-80	4.4	257
60	Mesenchymal stem cells and kidney repair. <i>Nephrology Dialysis Transplantation</i> , 2013 , 28, 788-93	4.3	82
59	A novel strategy to enhance mesenchymal stem cell migration capacity and promote tissue repair in an injury specific fashion. <i>Cell Transplantation</i> , 2013 , 22, 423-36	4	92
58	Localization of mesenchymal stromal cells dictates their immune or proinflammatory effects in kidney transplantation. <i>American Journal of Transplantation</i> , 2012 , 12, 2373-83	8.7	126
57	Mesenchymal stem cell therapy promotes renal repair by limiting glomerular podocyte and progenitor cell dysfunction in adriamycin-induced nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2012 , 303, F1370-81	4.3	71

56	In vivo maturation of functional renal organoids formed from embryonic cell suspensions. <i>Journal of the American Society of Nephrology: JASN</i> , 2012 , 23, 1857-68	12.7	125
55	Human amniotic fluid stem cell preconditioning improves their regenerative potential. <i>Stem Cells and Development</i> , 2012 , 21, 1911-23	4.4	103
54	Minimally manipulated whole human umbilical cord is a rich source of clinical-grade human mesenchymal stromal cells expanded in human platelet lysate. <i>Cytotherapy</i> , 2011 , 13, 786-801	4.8	93
53	Bone Marrow Mesenchymal Stem Cells in Organ Repair and Strategies to Optimize their Efficacy 2011 , 299-312		0
52	Inhibiting angiotensin-converting enzyme promotes renal repair by limiting progenitor cell proliferation and restoring the glomerular architecture. <i>American Journal of Pathology</i> , 2011 , 179, 628-38	5.8	90
51	MYO1E mutations and childhood familial focal segmental glomerulosclerosis. <i>New England Journal of Medicine</i> , 2011 , 365, 295-306	59.2	195
50	Embryonic stem cells, derived either after in vitro fertilization or nuclear transfer, prolong survival of semiallogeneic heart transplants. <i>Journal of Immunology</i> , 2011 , 186, 4164-74	5.3	8
49	Alternative pathway activation of complement by Shiga toxin promotes exuberant C3a formation that triggers microvascular thrombosis. <i>Journal of Immunology</i> , 2011 , 187, 172-80	5.3	186
48	Potential of mesenchymal stem cells in the repair of tubular injury. <i>Kidney International Supplements</i> , 2011 , 1, 90-93	6.3	11
47	Kidney regeneration. <i>Lancet, The</i> , 2010 , 375, 1310-7	4.0	113
46	Life-sparing effect of human cord blood-mesenchymal stem cells in experimental acute kidney injury. <i>Stem Cells</i> , 2010 , 28, 513-22	5.8	152
45	Shiga toxin-associated hemolytic uremic syndrome: pathophysiology of endothelial dysfunction. <i>Pediatric Nephrology</i> , 2010 , 25, 2231-40	3.2	137
44	Protein load impairs factor H binding promoting complement-dependent dysfunction of proximal tubular cells. <i>Kidney International</i> , 2009 , 75, 1050-9	9.9	24
43	Bone marrow-derived mesenchymal stem cells improve islet graft function in diabetic rats. <i>Transplantation Proceedings</i> , 2009 , 41, 1797-800	1.1	113
42	Disruption of the Ang II type 1 receptor promotes longevity in mice. <i>Journal of Clinical Investigation</i> , 2009 , 119, 524-30	15.9	374
41	Mesenchymal Stem Cells and Their Use in Acute Renal Injury 2009 , 216-220		
40	Complement-mediated dysfunction of glomerular filtration barrier accelerates progressive renal injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2008 , 19, 1158-67	12.7	54
39	Pretransplant infusion of mesenchymal stem cells prolongs the survival of a semiallogeneic heart transplant through the generation of regulatory T cells. <i>Journal of Immunology</i> , 2008 , 181, 3933-46	5.3	370

38	Fractalkine and CX3CR1 mediate leukocyte capture by endothelium in response to Shiga toxin. <i>Journal of Immunology</i> , 2008 , 181, 1460-9	5.3	35
37	Shiga-toxin-induced firm adhesion of human leukocytes to endothelium is in part mediated by heparan sulfate. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 3091-5	4.3	7
36	Human bone marrow mesenchymal stem cells accelerate recovery of acute renal injury and prolong survival in mice. <i>Stem Cells</i> , 2008 , 26, 2075-82	5.8	326
35	Insulin-like growth factor-1 sustains stem cell mediated renal repair. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2921-8	12.7	264
34	Permselective dysfunction of podocyte-podocyte contact upon angiotensin II unravels the molecular target for renoprotective intervention. <i>American Journal of Pathology</i> , 2006 , 168, 1073-85	5.8	74
33	Shigatoxin-induced endothelin-1 expression in cultured podocytes autocrinally mediates actin remodeling. <i>American Journal of Pathology</i> , 2006 , 169, 1965-75	5.8	85
32	The regenerative potential of stem cells in acute renal failure. <i>Cell Transplantation</i> , 2006 , 15 Suppl 1, S111-7	4	53
31	In response to protein load podocytes reorganize cytoskeleton and modulate endothelin-1 gene: implication for permselective dysfunction of chronic nephropathies. <i>American Journal of Pathology</i> , 2005 , 166, 1309-20	5.8	131
30	Activation of porcine endothelium in response to xenogeneic serum causes thrombosis independently of platelet activation. <i>Xenotransplantation</i> , 2005 , 12, 110-20	2.8	13
29	Genetics of rare diseases of the kidney: learning from mouse models. <i>Cytogenetic and Genome Research</i> , 2004 , 105, 479-84	1.9	4
28	Nitric oxide synthetic capacity in relation to dialysate temperature. <i>Blood Purification</i> , 2004 , 22, 203-9	3.1	17
27	Mesenchymal stem cells are renotropic, helping to repair the kidney and improve function in acute renal failure. <i>Journal of the American Society of Nephrology: JASN</i> , 2004 , 15, 1794-804	12.7	615
26	Vascular smooth muscle cells on hyaluronic acid: culture and mechanical characterization of an engineered vascular construct. <i>Tissue Engineering</i> , 2004 , 10, 699-710		55
25	Proteinuria and phenotypic change of proximal tubular cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14 Suppl 1, S36-41	12.7	71
24	Protein overload induces fractalkine upregulation in proximal tubular cells through nuclear factor kappaB- and p38 mitogen-activated protein kinase-dependent pathways. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, 2436-46	12.7	105
23	Effect of acetate-free biofiltration and bicarbonate hemodialysis on neutrophil activation. <i>American Journal of Kidney Diseases</i> , 2002 , 40, 783-93	7.4	57
22	Shiga toxin-2 triggers endothelial leukocyte adhesion and transmigration via NF-kappaB dependent up-regulation of IL-8 and MCP-1. <i>Kidney International</i> , 2002 , 62, 846-56	9.9	94
21	Transforming growth factor-beta1 is up-regulated by podocytes in response to excess intraglomerular passage of proteins: a central pathway in progressive glomerulosclerosis. <i>American Journal of Pathology</i> , 2002 , 161, 2179-93	5.8	116

20	Protein overload-induced NF-kappaB activation in proximal tubular cells requires H ₂ O ₂ through a PKC-dependent pathway. <i>Journal of the American Society of Nephrology: JASN</i> , 2002 , 13, 1179-89	12.7	133
19	Verotoxin-1-induced up-regulation of adhesive molecules renders microvascular endothelial cells thrombogenic at high shear stress. <i>Blood</i> , 2001 , 98, 1828-35	2.2	81
18	Influence of donor age on bovine pancreatic islet isolation. <i>Transplantation</i> , 2000 , 70, 1032-7	1.8	12
17	Shear stress-induced cytoskeleton rearrangement mediates NF-kappaB-dependent endothelial expression of ICAM-1. <i>Microvascular Research</i> , 2000 , 60, 182-8	3.7	25
16	Xenogeneic serum promotes leukocyte-endothelium interaction under flow through two temporally distinct pathways: role of complement and nuclear factor-kappaB. <i>Journal of the American Society of Nephrology: JASN</i> , 1999 , 10, 2197-207	12.7	20
15	Xenogeneic human serum promotes leukocyte adhesion to porcine endothelium under flow conditions, possibly through the activation of the transcription factor NF-kappa B. <i>Xenotransplantation</i> , 1998 , 5, 57-60	2.8	12
14	Bindarit retards renal disease and prolongs survival in murine lupus autoimmune disease. <i>Kidney International</i> , 1998 , 53, 726-34	9.9	63
13	Protein overload stimulates RANTES production by proximal tubular cells depending on NF-kappa B activation. <i>Kidney International</i> , 1998 , 53, 1608-15	9.9	324
12	Identification of a novel gene--SSK1--in human endothelial cells exposed to shear stress. <i>Biochemical and Biophysical Research Communications</i> , 1998 , 246, 881-7	3.4	6
11	Leukocyte-endothelial interaction is augmented by high glucose concentrations and hyperglycemia in a NF-kB-dependent fashion. <i>Journal of Clinical Investigation</i> , 1998 , 101, 1905-15	15.9	316
10	Fluid Shear Stress Modulates von Willebrand Factor Release From Human Vascular Endothelium. <i>Blood</i> , 1997 , 90, 1558-1564	2.2	110
9	Mycophenolate mofetil limits renal damage and prolongs life in murine lupus autoimmune disease. <i>Kidney International</i> , 1997 , 51, 1583-9	9.9	114
8	Fluid Shear Stress Modulates von Willebrand Factor Release From Human Vascular Endothelium. <i>Blood</i> , 1997 , 90, 1558-1564	2.2	5
7	Cyclosporine enhances leukocyte adhesion to vascular endothelium under physiologic flow conditions. <i>American Journal of Kidney Diseases</i> , 1996 , 28, 23-31	7.4	24
6	Proximal tubular cell synthesis and secretion of endothelin-1 on challenge with albumin and other proteins. <i>American Journal of Kidney Diseases</i> , 1995 , 26, 934-41	7.4	211
5	Nitric oxide synthesis by cultured endothelial cells is modulated by flow conditions. <i>Circulation Research</i> , 1995 , 76, 536-43	15.7	371
4	Turnover necrosis factor stimulates endothelin-1 gene expression in cultured bovine endothelial cells. <i>Mediators of Inflammation</i> , 1992 , 1, 263-6	4.3	4
3	Ticlopidine prevents renal disease progression in rats with reduced renal mass. <i>Kidney International</i> , 1990 , 37, 934-42	9.9	24

2	Platelet activating factor (PAF) as a mediator of injury in nephrotoxic nephritis. <i>Kidney International</i> , 1987 , 31, 1248-56	9.9	43
1	SARS-CoV-2 Spike Protein 1 Activates Microvascular Endothelial Cells and Complement System Leading to Thrombus Formation. <i>SSRN Electronic Journal</i> ,	1	1