

Shintaro Ichikawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1458226/publications.pdf>

Version: 2024-02-01

66
papers

1,548
citations

331670

21
h-index

330143

37
g-index

66
all docs

66
docs citations

66
times ranked

1908
citing authors

#	ARTICLE	IF	CITATIONS
1	An Intra-individual Comparison between Free-breathing Dynamic MR Imaging of the Liver Using Stack-of-stars Acquisition and the Breath-holding Method Using Cartesian Sampling or View-sharing. <i>Magnetic Resonance in Medical Sciences</i> , 2023, 22, 221-231.	2.0	6
2	MRI-based radiomics analysis for differentiating phyllodes tumors of the breast from fibroadenomas. <i>European Radiology</i> , 2022, 32, 4090-4100.	4.5	13
3	Distinction Between Hepatocellular Carcinoma and Hypervascular Liver Metastases in Non-cirrhotic Patients Using Gadoxetate Disodium-Enhanced Magnetic Resonance Imaging. <i>Canadian Association of Radiologists Journal</i> , 2022, 73, 639-646.	2.0	3
4	Accelerated Acquisition of High-resolution Diffusion-weighted Imaging of the Brain with a Multi-shot Echo-planar Sequence: Deep-learning-based Denoising. <i>Magnetic Resonance in Medical Sciences</i> , 2021, 20, 99-105.	2.0	24
5	Optimal Combination of Features on Gadoxetate Disodium-enhanced MR Imaging for Non-invasive Differential Diagnosis of Hepatocellular Carcinoma: The JAMP-HCC Study. <i>Magnetic Resonance in Medical Sciences</i> , 2021, 20, 47-59.	2.0	7
6	Optimal target b-value on computed diffusion-weighted magnetic resonance imaging for visualization of pancreatic ductal adenocarcinoma and focal autoimmune pancreatitis. <i>Abdominal Radiology</i> , 2021, 46, 636-646.	2.1	6
7	Gadolinium-Based Contrast Agent Toxicity and Accumulation. , 2021, , 77-87.		1
8	Key Imaging Findings for the Prospective Diagnosis of Rare Diseases of the Gallbladder and Cystic Duct. <i>Korean Journal of Radiology</i> , 2021, 22, 1462.	3.4	2
9	Multi-arterial phase MRI depicts inconsistent arterial phase hyperenhancement (APHE) subtypes in liver observations of patients at risk for hepatocellular carcinoma. <i>European Radiology</i> , 2021, 31, 7594-7604.	4.5	4
10	Uptake of gadoteric acid in hepatobiliary phase magnetic resonance imaging and transporter expression in hypovascular hepatocellular nodules. <i>European Journal of Radiology</i> , 2021, 138, 109669.	2.6	1
11	Diagnostic performance and image quality of low-tube voltage and low-contrast medium dose protocol with hybrid iterative reconstruction for hepatic dynamic CT. <i>British Journal of Radiology</i> , 2021, 94, 20210601.	2.2	8
12	Preoperative CT Findings for Predicting Acute Exacerbation of Interstitial Pneumonia After Lung Cancer Surgery: A Multicenter Case-Control Study. <i>American Journal of Roentgenology</i> , 2021, 217, 859-869.	2.2	2
13	Pathologic, Molecular, and Prognostic Radiologic Features of Hepatocellular Carcinoma. <i>Radiographics</i> , 2021, 41, 1611-1631.	3.3	32
14	CNN color-coded difference maps accurately display longitudinal changes in liver MRI-PDFF. <i>European Radiology</i> , 2021, 31, 5041-5049.	4.5	1
15	Transient Respiratory-motion Artifact and Scan Timing during the Arterial Phase of Gadoxetate Disodium-enhanced MR Imaging: The Benefit of Shortened Acquisition and Multiple Arterial Phase Acquisition. <i>Magnetic Resonance in Medical Sciences</i> , 2021, 20, 280-289.	2.0	10
16	Imaging findings of immunoglobulin G4-related disease: from the head to the pelvis. <i>Singapore Medical Journal</i> , 2021, 62, 574-581.	0.6	1
17	Motion Artifact Reduction Using a Convolutional Neural Network for Dynamic Contrast Enhanced MR Imaging of the Liver. <i>Magnetic Resonance in Medical Sciences</i> , 2020, 19, 64-76.	2.0	87
18	MRI-based risk factors of hepatocellular carcinoma in patients with chronic liver disease: A prospective observational study. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 389-396.	3.4	9

#	ARTICLE	IF	CITATIONS
19	Predicting Patients With Insufficient Liver Enhancement in the Hepatobiliary Phase Before the Injection of Gadoteric Acid: A Practical Approach Using the Bayesian Method. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 62-69.	3.4	4
20	Utility of Stack-of-stars Acquisition for Hepatobiliary Phase Imaging without Breath-holding. <i>Magnetic Resonance in Medical Sciences</i> , 2020, 19, 99-107.	2.0	9
21	Clinical Evaluation of Respiratory-triggered 3D MRCP with Navigator Echoes Compared to Breath-hold Acquisition Using Compressed Sensing and/or Parallel Imaging. <i>Magnetic Resonance in Medical Sciences</i> , 2020, 19, 318-323.	2.0	9
22	Imaging features of hepatic inflammatory pseudotumor: distinction from colorectal liver metastasis using gadoteric acid-enhanced magnetic resonance imaging. <i>Abdominal Radiology</i> , 2020, 45, 2400-2408.	2.1	10
23	Distinguishing intrahepatic mass-forming biliary carcinomas from hepatocellular carcinoma by computed tomography and magnetic resonance imaging using the Bayesian method: a bi-center study. <i>European Radiology</i> , 2020, 30, 5992-6002.	4.5	14
24	Diffusion-weighted MRI-based Virtual Elastography for the Assessment of Liver Fibrosis. <i>Radiology</i> , 2020, 295, 127-135.	7.3	37
25	Imaging Findings of Acute Abdomen due to Complications of Meckel Diverticulum. <i>Canadian Association of Radiologists Journal</i> , 2020, 71, 149-153.	2.0	3
26	Neuroendocrine carcinoma of uterine cervix findings shown by MRI for staging and survival analysis - Japan multicenter study. <i>Oncotarget</i> , 2020, 11, 3675-3686.	1.8	8
27	Magnetic resonance elastography can predict development of hepatocellular carcinoma with longitudinally acquired two-point data. <i>European Radiology</i> , 2019, 29, 1013-1021.	4.5	24
28	MR-guided Focused Ultrasound for Uterine Fibroids: A Preliminary Study of Relationship between the Treatment Outcomes and Factors of MR Images Including Elastography. <i>Magnetic Resonance in Medical Sciences</i> , 2019, 18, 82-87.	2.0	7
29	Fully automated convolutional neural network-based affine algorithm improves liver registration and lesion co-localization on hepatobiliary phase T1-weighted MR images. <i>European Radiology Experimental</i> , 2019, 3, 43.	3.4	12
30	Diffusion-weighted image improves detectability of magnetic resonance cholangiopancreatography for pancreatic ductal adenocarcinoma concomitant with intraductal papillary mucinous neoplasm. <i>Medicine (United States)</i> , 2019, 98, e18039.	1.0	9
31	Improving the Quality of Diffusion-weighted Imaging of the Left Hepatic Lobe Using Weighted Averaging of Signals from Multiple Excitations. <i>Magnetic Resonance in Medical Sciences</i> , 2019, 18, 225-232.	2.0	11
32	Linear gadolinium-based contrast agent (gadodiamide and gadopentetate dimeglumine)-induced high signal intensity on unenhanced T1-weighted images in pediatric patients. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 49, 1046-1052.	3.4	5
33	Dose-dependence of transient respiratory motion artifacts on gadoteric acid-enhanced arterial phase MR images. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 433-438.	3.4	7
34	Ring-Like Enhancement of Hepatocellular Carcinoma in Gadoteric Acid-Enhanced Multiphasic Hepatic Arterial Phase Imaging With Differential Subsampling With Cartesian Ordering. <i>Investigative Radiology</i> , 2018, 53, 191-199.	6.2	16
35	Magnetic resonance elastography is as accurate as liver biopsy for liver fibrosis staging. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 1268-1275.	3.4	51
36	Feasibility of Combination Therapy with Nab-paclitaxel Plus Gemcitabine in Patients with Recurrent Pancreatic Cancer. <i>Anticancer Research</i> , 2018, 38, 6537-6542.	1.1	4

#	ARTICLE	IF	CITATIONS
37	Histological Grading of Hepatocellular Carcinomas with Intravoxel Incoherent Motion Diffusion-weighted Imaging: Inconsistent Results Depending on the Fitting Method. <i>Magnetic Resonance in Medical Sciences</i> , 2018, 17, 168-173.	2.0	14
38	Prognostic value of preoperative fluorodeoxyglucose positron emission tomography/computed tomography in patients with potentially resectable pancreatic cancer. <i>Abdominal Radiology</i> , 2018, 43, 3381-3389.	2.1	6
39	Noninvasive hepatic fibrosis staging using mr elastography: The usefulness of the bayesian prediction method. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 46, 375-382.	3.4	17
40	Magnetic resonance elastography for prediction of radiation-induced liver disease after stereotactic body radiation therapy. <i>Hepatology</i> , 2017, 66, 664-665.	7.3	8
41	Contrast Agent-induced High Signal Intensity in Dentate Nucleus on Unenhanced T1-Weighted Images. <i>Investigative Radiology</i> , 2017, 52, 389-395.	6.2	26
42	Diffusion and Intravoxel Incoherent Motion MR Imaging-based Virtual Elastography: A Hypothesis-generating Study in the Liver. <i>Radiology</i> , 2017, 285, 609-619.	7.3	44
43	Outcome of hypovascular hepatic nodules with positive uptake of gadoxetic acid in patients with cirrhosis. <i>European Radiology</i> , 2017, 27, 518-525.	4.5	10
44	Comparison of diagnostic accuracies of two- and three-dimensional MR elastography of the liver. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 1163-1170.	3.4	44
45	Added Value of a Gadoxetic Acid-enhanced Hepatocyte-phase Image to the LI-RADS System for Diagnosing Hepatocellular Carcinoma. <i>Magnetic Resonance in Medical Sciences</i> , 2016, 15, 49-59.	2.0	55
46	Usefulness of MR elastography for detecting clinical progression of cirrhosis from child-pugh class A to B in patients with type C viral hepatitis. <i>Journal of Magnetic Resonance Imaging</i> , 2016, 44, 715-722.	3.4	21
47	Morphophenotypic changes in human multistep hepatocarcinogenesis with translational implications. <i>Journal of Hepatology</i> , 2016, 64, 87-93.	3.7	40
48	Was Hypervascular Hepatocellular Carcinoma Visible on Previous Gadoxetic Acid-Enhanced Magnetic Resonance Images?. <i>Liver Cancer</i> , 2015, 4, 154-162.	7.7	20
49	Hepatitis activity should be considered a confounder of liver stiffness measured with MR elastography. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1203-1208.	3.4	42
50	MRI-based staging of hepatic fibrosis: Comparison of intravoxel incoherent motion diffusion-weighted imaging with magnetic resonance elastography. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 42, 204-210.	3.4	68
51	Validity and Reliability of Magnetic Resonance Elastography for Staging Hepatic Fibrosis in Patients with Chronic Hepatitis B. <i>Magnetic Resonance in Medical Sciences</i> , 2015, 14, 211-221.	2.0	17
52	Comparison of the diagnostic accuracies of magnetic resonance elastography and transient elastography for hepatic fibrosis. <i>Magnetic Resonance Imaging</i> , 2015, 33, 26-30.	1.8	84
53	Association of splenic MR elastographic findings with gastroesophageal varices in patients with chronic liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 117-124.	3.4	42
54	Presence of a hypovascular hepatic nodule showing hypointensity on hepatocyte-phase image is a risk factor for hypervascular hepatocellular carcinoma. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 293-297.	3.4	30

#	ARTICLE	IF	CITATIONS
55	Computed Tomography (CT) Venography With Dual-Energy CT. Journal of Computer Assisted Tomography, 2014, 38, 797-801.	0.9	8
56	Primer on magnetic resonance imaging of the liver. Clinical Liver Disease, 2014, 4, 120-123.	2.1	1
57	Liver stiffness measured by magnetic resonance elastography as a risk factor for hepatocellular carcinoma: a preliminary case-control study. European Radiology, 2013, 23, 156-162.	4.5	59
58	Intravoxel incoherent motion imaging of the kidney: alterations in diffusion and perfusion in patients with renal dysfunction. Magnetic Resonance Imaging, 2013, 31, 414-417.	1.8	90
59	Intravoxel incoherent motion imaging of focal hepatic lesions. Journal of Magnetic Resonance Imaging, 2013, 37, 1371-1376.	3.4	47
60	Early Hepatocellular Carcinomas Showing Isointensity or Hyperintensity in Gadoteric Acid-Enhanced, Hepatocyte-Phase Magnetic Resonance Images. Journal of Computer Assisted Tomography, 2013, 37, 466-469.	0.9	8
61	MR-based Measurements of Portal Vein Flow and Liver Stiffness for Predicting Gastroesophageal Varices. Magnetic Resonance in Medical Sciences, 2013, 12, 77-86.	2.0	18
62	Magnetic Resonance Elastography as a Predictor of Insufficient Liver Enhancement on Gadoteric Acid-Enhanced Hepatocyte-Phase Magnetic Resonance Imaging in Patients With Type C Hepatitis and Child-Pugh Class A Disease. Investigative Radiology, 2012, 47, 566-570.	6.2	15
63	Magnetic Resonance Elastography for Staging Liver Fibrosis in Chronic Hepatitis C. Magnetic Resonance in Medical Sciences, 2012, 11, 291-297.	2.0	110
64	Dilution method of gadolinium ethoxybenzyl diethylenetriaminepentaacetic acid (Gd-EOB-DTPA)-enhanced magnetic resonance imaging (MRI). Journal of Magnetic Resonance Imaging, 2009, 30, 849-854.	3.4	67
65	Delay before the hepatocyte phase of Gd-EOB-DTPA-enhanced MR imaging: Is it possible to shorten the examination time?. European Radiology, 2009, 19, 2623-2629.	4.5	80
66	Computed tomography and magnetic resonance imaging findings of gynaecologic emergencies: A pictorial essay. Journal of Medical Imaging and Radiation Oncology, 0, , .	1.8	0