

Chang-Qin Liu

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1467864/chang-qin-liu-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

52
papers

1,209
citations

17
h-index

34
g-index

55
ext. papers

1,464
ext. citations

4.6
avg, IF

3.84
L-index

#	Paper	IF	Citations
52	Plasma exosomal β synuclein is likely CNS-derived and increased in Parkinson's disease. <i>Acta Neuropathologica</i> , 2014 , 128, 639-650	14.3	348
51	Cerebrospinal fluid β synuclein predicts cognitive decline in Parkinson disease progression in the DATATOP cohort. <i>American Journal of Pathology</i> , 2014 , 184, 966-975	5.8	98
50	Inhibition of Wnt/ β catenin signaling downregulates P-glycoprotein and reverses multi-drug resistance of cholangiocarcinoma. <i>Cancer Science</i> , 2013 , 104, 1303-8	6.9	77
49	β synuclein in cerebrospinal fluid of Alzheimer's disease and mild cognitive impairment. <i>Journal of Alzheimer's Disease</i> , 2013 , 36, 679-88	4.3	62
48	CSF tau and tau/A β 2 predict cognitive decline in Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2015 , 21, 271-6	3.6	57
47	Longitudinal assessment of tau and amyloid beta in cerebrospinal fluid of Parkinson disease. <i>Acta Neuropathologica</i> , 2013 , 126, 671-82	14.3	56
46	Ebselen treatment prevents islet apoptosis, maintains intranuclear Pdx-1 and MafA levels, and preserves β cell mass and function in ZDF rats. <i>Diabetes</i> , 2013 , 62, 3582-8	0.9	56
45	Phosphorylated β synuclein in Parkinson's disease: correlation depends on disease severity. <i>Acta Neuropathologica Communications</i> , 2015 , 3, 7	7.3	53
44	Puerarin improves insulin resistance and modulates adipokine expression in rats fed a high-fat diet. <i>European Journal of Pharmacology</i> , 2010 , 649, 398-402	5.3	51
43	Chinese lacto-vegetarian diet exerts favorable effects on metabolic parameters, intima-media thickness, and cardiovascular risks in healthy men. <i>Nutrition in Clinical Practice</i> , 2012 , 27, 392-8	3.6	37
42	Elevated circulating irisin is associated with lower risk of insulin resistance: association and path analyses of obese Chinese adults. <i>BMC Endocrine Disorders</i> , 2016 , 16, 44	3.3	23
41	Relationship of carotid intima-media thickness and duration of vegetarian diet in Chinese male vegetarians. <i>Nutrition and Metabolism</i> , 2011 , 8, 63	4.6	23
40	Serum uric acid is independently and linearly associated with risk of nonalcoholic fatty liver disease in obese Chinese adults. <i>Scientific Reports</i> , 2016 , 6, 38605	4.9	20
39	Association of serum irisin and body composition with chronic kidney disease in obese Chinese adults: a cross-sectional study. <i>BMC Nephrology</i> , 2015 , 16, 16	2.7	19
38	Metabolomic profilings of urine and serum from high fat-fed rats via 1H NMR spectroscopy and pattern recognition. <i>Applied Biochemistry and Biotechnology</i> , 2013 , 169, 1250-61	3.2	18
37	Activation of nuclear factor- κ B pathway is responsible for tumor necrosis factor- α -induced up-regulation of endothelin B2 receptor expression in vascular smooth muscle cells in vitro. <i>Toxicology Letters</i> , 2012 , 209, 107-12	4.4	18
36	Fetuin-B links nonalcoholic fatty liver disease to type 2 diabetes via inducing insulin resistance: Association and path analyses. <i>Cytokine</i> , 2018 , 108, 145-150	4	17

35	Both interleukin-23A polymorphism and serum interleukin-23 expression are associated with Graves disease risk. <i>Cellular Immunology</i> , 2015 , 294, 39-43	4.4	17
34	Genetic association between IL-17F gene polymorphisms and the pathogenesis of Graves disease in the Han Chinese population. <i>Gene</i> , 2013 , 512, 300-4	3.8	17
33	Evaluation of the three-in-one team-based care model on hierarchical diagnosis and treatment patterns among patients with diabetes: a retrospective cohort study using Xiamen regional electronic health records. <i>BMC Health Services Research</i> , 2017 , 17, 779	2.9	14
32	Hepatic fat content is a determinant of metabolic phenotypes and increased carotid intima-media thickness in obese adults. <i>Scientific Reports</i> , 2016 , 6, 21894	4.9	13
31	Serum fetuin-B is positively associated with intrahepatic triglyceride content and increases the risk of insulin resistance in obese Chinese adults: A cross-sectional study. <i>Journal of Diabetes</i> , 2018 , 10, 581-588	3.8	13
30	An association of interleukin-10 gene polymorphisms with Graves disease in two Chinese populations. <i>Endocrine</i> , 2011 , 40, 90-4	4	12
29	Risk of tuberculosis in patients with immune-mediated diseases on biological therapies: a population-based study in a tuberculosis endemic region. <i>Rheumatology</i> , 2019 , 58, 803-810	3.9	12
28	Serum Uric Acid Is Independently Associated with Risk of Obstructive Sleep Apnea-Hypopnea Syndrome in Chinese Patients with Type 2 Diabetes. <i>Disease Markers</i> , 2019 , 2019, 4578327	3.2	10
27	Correlations of non-alcoholic fatty liver disease and serum uric acid with subclinical atherosclerosis in obese Chinese adults. <i>Journal of Diabetes</i> , 2017 , 9, 586-595	3.8	8
26	Associations of apnea hypopnea index and educational attainments with microvascular complications in patients with T2DM. <i>Endocrine</i> , 2020 , 67, 363-373	4	7
25	Lipid accumulation product independently correlate with hepatic steatosis quantified by controlled attenuation parameter in women with polycystic ovary syndrome. <i>Endocrine Connections</i> , 2020 , 9, 154-162	3.5	6
24	Neck circumference is a good predictor for insulin resistance in women with polycystic ovary syndrome. <i>Fertility and Sterility</i> , 2021 , 115, 753-760	4.8	6
23	Fetuin-B Links Nonalcoholic Fatty Liver Disease to Chronic Kidney Disease in Obese Chinese Adults: A Cross-Sectional Study. <i>Annals of Nutrition and Metabolism</i> , 2019 , 74, 287-295	4.5	5
22	Lipid accumulation product (LAP) was independently associated with obstructive sleep apnea in patients with type 2 diabetes mellitus. <i>BMC Endocrine Disorders</i> , 2020 , 20, 179	3.3	5
21	Effect of exenatide after short-time intensive insulin therapy on glycaemic remission maintenance in type 2 diabetes patients: a randomized controlled trial. <i>Scientific Reports</i> , 2017 , 7, 2383	4.9	4
20	Independent Association of Serum Fibroblast Growth Factor 21 Levels With Impaired Liver Enzymes in Hyperthyroid Patients. <i>Frontiers in Endocrinology</i> , 2018 , 9, 800	5.7	4
19	Common genetic variants in the FETUB locus, genetically predicted fetuin-B levels, and risk of insulin resistance in obese Chinese adults. <i>Medicine (United States)</i> , 2017 , 96, e9234	1.8	3
18	Association of Fetuin-B with Subclinical Atherosclerosis in Obese Chinese Adults. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 418-428	4	3

17	Independent associations of thyroid-related hormones with hepatic steatosis and insulin resistance in euthyroid overweight/obese Chinese adults. <i>BMC Gastroenterology</i> , 2021 , 21, 431	3	2
16	Dysmyelination by Oligodendrocyte-Specific Ablation of Ninj2 Contributes to Depressive-Like Behaviors. <i>Advanced Science</i> , 2021 , e2103065	13.6	2
15	The Relationship of Serum 25-Hydroxyvitamin-D Level with Severity of Obstructive Sleep Apnea in Patients with Type 2 Diabetes Mellitus. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 1391-1398	3.4	2
14	Maternal postload 1-hour glucose level during pregnancy and offspring's overweight/obesity status in preschool age. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	2
13	Correlation Between Daily Energy Intake from Fat with Insulin Resistance in Patients with Polycystic Ovary Syndrome. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2021 , 14, 295-303	3.4	2
12	Higher Apnea-Hypopnea Index (AHI) and Oxygen Desaturation Index (ODI) Were Independently Associated with Increased Risks of Hypertension in Patients with T2DM: A Cross-Sectional Study. <i>International Journal of Hypertension</i> , 2021 , 2021, 8887944	2.4	2
11	The rs4686434 variant in the fetuin B (FETUB) locus is associated with intrahepatic triglyceride content in obese Chinese adults. <i>Journal of Diabetes</i> , 2018 , 10, 916-925	3.8	1
10	Benefits and harms of hemithyroidectomy, total or near-total thyroidectomy in 1-4 cm differentiated thyroid cancer. <i>Clinical Endocrinology</i> , 2021 , 95, 668-676	3.4	1
9	DDB1 binds histone reader BRWD3 to activate the transcriptional cascade in adipogenesis and promote onset of obesity. <i>Cell Reports</i> , 2021 , 35, 109281	10.6	1
8	The Mediator subunit MED20 organizes the early adipogenic complex to promote development of adipose tissues and diet-induced obesity. <i>Cell Reports</i> , 2021 , 36, 109314	10.6	1
7	Joint associations of metabolically healthy abdominal obesity and non-alcoholic fatty liver disease with prediabetes and diabetes in Chinese adults. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	1
6	Comparative effectiveness of team-based care with a clinical decision support system versus team-based care alone on cardiovascular risk reduction among patients with diabetes: Rationale and design of the D4C trial. <i>American Heart Journal</i> , 2021 , 238, 45-58	4.9	0
5	Association between intrahepatic triglyceride content in subjects with metabolically healthy abdominal obesity and risks of pre-diabetes plus diabetes: an observational study.. <i>BMJ Open</i> , 2022 , 12, e057820	3	0
4	Nurse-led care versus usual care on cardiovascular risk factors for patients with type 2 diabetes: a systematic review and meta-analysis.. <i>BMJ Open</i> , 2022 , 12, e058533	3	0
3	Relationship between diabetes and risk of gastric cancer: A systematic review and meta-analysis of cohort studies.. <i>Diabetes Research and Clinical Practice</i> , 2022 , 109866	7.4	0
2	Visceral Adipose Tissue Indices Independently Correlated with Obstructive Sleep Apnea in Patients with Type 2 Diabetes.. <i>Journal of Diabetes Research</i> , 2022 , 2022, 4950528	3.9	
1	Coexistence of ovarian serous papillary cystadenofibroma and type A insulin resistance syndrome in a 14-year-old girl: A case report. <i>World Journal of Clinical Cases</i> , 2020 , 8, 3334-3340	1.6	