

Natalia I Dmitrieva

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1473143/publications.pdf>

Version: 2024-02-01

47
papers

2,566
citations

236833

25
h-index

223716

46
g-index

47
all docs

47
docs citations

47
times ranked

3420
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Middle age serum sodium levels in the upper part of normal range and risk of heart failure. <i>European Heart Journal</i> , 2022, 43, 3335-3348. | 1.0 | 19 |
| 2 | Chronic habitual hypohydration that elevates serum sodium above 142 mmol/l is a risk factor for accelerated cognitive decline and dementia, suggesting lifelong optimal hydration as preventive measure. <i>Alzheimer's and Dementia</i> , 2021, 17, . | 0.4 | 0 |
| 3 | Mutations that prevent caspase cleavage of RIPK1 cause autoinflammatory disease. <i>Nature</i> , 2020, 577, 103-108. | 13.7 | 198 |
| 4 | STAT3 modulates reprogramming efficiency of human somatic cells; Insights from autosomal dominant Hyper IgE syndrome caused by STAT3 mutations. <i>Biology Open</i> , 2020, 9, . | 0.6 | 3 |
| 5 | Generation of human induced pluripotent stem cells (NIHTVBi004-A, NIHTVBi005-A, NIHTVBi006-A,) Tj ETQq1 1 0.784314 rgBT /Overl 45, 101821. | 0.3 | 1 |
| 6 | Impaired angiogenesis and extracellular matrix metabolism in autosomal-dominant hyper-IgE syndrome. <i>Journal of Clinical Investigation</i> , 2020, 130, 4167-4181. | 3.9 | 13 |
| 7 | Generation of human induced pluripotent stem cell lines (NIHTVBi011-A, NIHTVBi012-A, NIHTVBi013-A) from autosomal dominant Hyper IgE syndrome (AD-HIES) patients carrying STAT3 mutation. <i>Stem Cell Research</i> , 2019, 41, 101586. | 0.3 | 5 |
| 8 | Generation of human induced pluripotent stem cells from individuals with a homozygous CCR5 ^{Δ32} mutation. <i>Stem Cell Research</i> , 2019, 38, 101481. | 0.3 | 6 |
| 9 | Suboptimal hydration remodels metabolism, promotes degenerative diseases, and shortens life. <i>JCI Insight</i> , 2019, 4, . | 2.3 | 25 |
| 10 | Cross-Sectional Positive Association of Serum Lipids and Blood Pressure With Serum Sodium Within the Normal Reference Range of 135–145 mmol/L. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017, 37, 598-606. | 1.1 | 22 |
| 11 | Increased activity of TNAP compensates for reduced adenosine production and promotes ectopic calcification in the genetic disease ACDC. <i>Science Signaling</i> , 2016, 9, ra121. | 1.6 | 65 |
| 12 | Elevated Sodium and Dehydration Stimulate Inflammatory Signaling in Endothelial Cells and Promote Atherosclerosis. <i>PLoS ONE</i> , 2015, 10, e0128870. | 1.1 | 66 |
| 13 | Secretion of von Willebrand factor by endothelial cells links sodium to hypercoagulability and thrombosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 6485-6490. | 3.3 | 86 |
| 14 | Global discovery of high-NaCl-induced changes of protein phosphorylation. <i>American Journal of Physiology - Cell Physiology</i> , 2014, 307, C442-C454. | 2.1 | 16 |
| 15 | DNA double-strand breaks induced by high NaCl occur predominantly in gene deserts. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 20796-20801. | 3.3 | 48 |
| 16 | Mre11 is expressed in mammalian mitochondria where it binds to mitochondrial DNA. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2011, 301, R632-R640. | 0.9 | 43 |
| 17 | Increased Insensible Water Loss Contributes to Aging Related Dehydration. <i>PLoS ONE</i> , 2011, 6, e20691. | 1.1 | 26 |
| 18 | Mediator of DNA Damage Checkpoint 1 (MDC1) Contributes to High NaCl-Induced Activation of the Osmoprotective Transcription Factor TonEBP/OREBP. <i>PLoS ONE</i> , 2010, 5, e12108. | 1.1 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Knockout of Ku86 accelerates cellular senescence induced by high NaCl. <i>Aging</i> , 2009, 1, 245-253. | 1.4 | 3 |
| 20 | Analysis of DNA breaks, DNA damage response, and apoptosis produced by high NaCl. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F1678-F1688. | 1.3 | 36 |
| 21 | MKP-1 inhibits high NaCl-induced activation of p38 but does not inhibit the activation of TonEBP/OREBP: Opposite roles of p38 ^Δ and p38 ^Δ . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 5620-5625. | 3.3 | 34 |
| 22 | Helper B Cells Promote Cytotoxic T Cell Survival and Proliferation Independently of Antigen Presentation through CD27/CD70 Interactions. <i>Journal of Immunology</i> , 2008, 180, 1362-1372. | 0.4 | 77 |
| 23 | High NaCl Promotes Cellular Senescence. <i>Cell Cycle</i> , 2007, 6, 3108-3113. | 1.3 | 46 |
| 24 | Cellular Response to Hyperosmotic Stresses. <i>Physiological Reviews</i> , 2007, 87, 1441-1474. | 13.1 | 635 |
| 25 | Osmotic Stress and DNA Damage. <i>Methods in Enzymology</i> , 2007, 428, 241-252. | 0.4 | 20 |
| 26 | Effects of expression of p53 and Gadd45 on osmotic tolerance of renal inner medullary cells. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, F341-F349. | 1.3 | 12 |
| 27 | The Saltiness of the Sea Breaks DNA in Marine Invertebrates: Possible Implications for Animal Evolution. <i>Cell Cycle</i> , 2006, 5, 1320-1323. | 1.3 | 24 |
| 28 | Hypertonic stress response. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2005, 569, 65-74. | 0.4 | 66 |
| 29 | Ataxia telangiectasia-mutated, a DNA damage-inducible kinase, contributes to high NaCl-induced nuclear localization of transcription factor TonEBP/OREBP. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F506-F511. | 1.3 | 38 |
| 30 | DNA damage and osmotic regulation in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F2-F7. | 1.3 | 23 |
| 31 | Pax2 expression occurs in renal medullary epithelial cells in vivo and in cell culture, is osmoregulated, and promotes osmotic tolerance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 503-508. | 3.3 | 60 |
| 32 | Ku86 preserves chromatin integrity in cells adapted to high NaCl. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 10730-10735. | 3.3 | 29 |
| 33 | Cells adapted to high NaCl have many DNA breaks and impaired DNA repair both in cell culture and in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 2317-2322. | 3.3 | 109 |
| 34 | Cell Cycle-dependent Expression of Thyroid Hormone Receptor- β^2 Is a Mechanism for Variable Hormone Sensitivity. <i>Molecular Biology of the Cell</i> , 2004, 15, 1895-1903. | 0.9 | 27 |
| 35 | Living with DNA Breaks is an Everyday Reality for Cells Adapted to High NaCl. <i>Cell Cycle</i> , 2004, 3, 559-561. | 1.3 | 16 |
| 36 | From The Cover: High urea and NaCl carbonylate proteins in renal cells in culture and in vivo, and high urea causes 8-oxoguanine lesions in their DNA. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 9491-9496. | 3.3 | 151 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Living with DNA breaks is an everyday reality for cells adapted to high NaCl. <i>Cell Cycle</i> , 2004, 3, 561-3. | 1.3 | 8 |
| 38 | High NaCl causes Mre11 to leave the nucleus, disrupting DNA damage signaling and repair. <i>American Journal of Physiology - Renal Physiology</i> , 2003, 285, F266-F274. | 1.3 | 64 |
| 39 | Toxicity of Acetaminophen, Salicylic Acid, and Caffeine for First-Passage Rat Renal Inner Medullary Collecting Duct Cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003, 306, 35-42. | 1.3 | 15 |
| 40 | Rapid activation of G2/M checkpoint after hypertonic stress in renal inner medullary epithelial (IME) cells is protective and requires p38 kinase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 184-189. | 3.3 | 79 |
| 41 | Proliferation and osmotic tolerance of renal inner medullary epithelial cells in vivo and in cell culture. <i>American Journal of Physiology - Renal Physiology</i> , 2002, 283, F302-F308. | 1.3 | 33 |
| 42 | Mitochondrial dysfunction is an early event in high-NaCl-induced apoptosis of mIMCD3 cells. <i>American Journal of Physiology - Renal Physiology</i> , 2002, 282, F981-F990. | 1.3 | 54 |
| 43 | Rate of increase of osmolality determines osmotic tolerance of mouse inner medullary epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2002, 283, F792-F798. | 1.3 | 25 |
| 44 | p53 Protects renal inner medullary cells from hypertonic stress by restricting DNA replication. <i>American Journal of Physiology - Renal Physiology</i> , 2001, 281, F522-F530. | 1.3 | 45 |
| 45 | Cell cycle delay and apoptosis in response to osmotic stress. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2001, 130, 411-420. | 0.8 | 73 |
| 46 | Protection of Renal Inner Medullary Epithelial Cells from Apoptosis by Hypertonic Stress-induced p53 Activation. <i>Journal of Biological Chemistry</i> , 2000, 275, 18243-18247. | 1.6 | 99 |
| 47 | Involvement of endogenous digitalis-like factors involuntary selection of alcohol by rats. <i>Life Sciences</i> , 1999, 64, PL219-PL225. | 2.0 | 15 |