Zouher Amzil

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1475147/publications.pdf

Version: 2024-02-01

	172207	233125
2,230	29	45
citations	h-index	g-index
62	60	1776
62	62	1776
docs citations	times ranked	citing authors
	citations 62	2,230 29 citations h-index 62 62

#	Article	IF	CITATIONS
1	Ovatoxin-a and Palytoxin Accumulation in Seafood in Relation to Ostreopsis cf. ovata Blooms on the French Mediterranean Coast. Marine Drugs, 2012, 10, 477-496.	2.2	125
2	Domoic acid accumulation in French shellfish in relation to toxic species of Pseudo-nitzschia multiseries and P. pseudodelicatissima. Toxicon, 2001, 39, 1245-1251.	0.8	100
3	First Evidence of Palytoxin and 42-Hydroxy-palytoxin in the Marine Cyanobacterium Trichodesmium. Marine Drugs, 2011, 9, 543-560.	2.2	99
4	Toxicity screening of 13 Gambierdiscus strains using neuro-2a and erythrocyte lysis bioassays. Harmful Algae, 2017, 63, 173-183.	2.2	98
5	Complex Toxin Profile of French Mediterranean Ostreopsis cf. ovata Strains, Seafood Accumulation and Ovatoxins Prepurification. Marine Drugs, 2014, 12, 2851-2876.	2.2	78
6	Pinnatoxin G is responsible for atypical toxicity in mussels (Mytilus galloprovincialis) and clams (Venerupis decussata) from Ingril, a French Mediterranean lagoon. Toxicon, 2013, 75, 16-26.	0.8	74
7	Report on the First Detection of Pectenotoxin-2, Spirolide-A and Their Derivatives in French Shellfish. Marine Drugs, 2007, 5, 168-179.	2.2	73
8	Î ² -N-methylamino-l-alanine (BMAA) and isomers: Distribution in different food web compartments of Thau lagoon, French Mediterranean Sea. Marine Environmental Research, 2015, 110, 8-18.	1.1	73
9	First report on azaspiracid and yessotoxin groups detection in French shellfish. Toxicon, 2008, 52, 39-48.	0.8	68
10	Characterization of ovatoxin-h, a new ovatoxin analog, and evaluation of chromatographic columns for ovatoxin analysis and purification. Journal of Chromatography A, 2015, 1388, 87-101.	1.8	61
11	Production of BMAA and DAB by diatoms (Phaeodactylum tricornutum , Chaetoceros sp., Chaetoceros) Tj ETQq1 Algae, 2016, 58, 45-50.		14 rgBT /Ov∈ 61
12	Toxicity and Growth Assessments of Three Thermophilic Benthic Dinoflagellates (Ostreopsis cf. ovata,) Tj ETQq0 0 2016, 8, 297.	0 0 rgBT /Ov 1.5	verlock 10 T 59
13	Influence of Environmental Factors on the Paralytic Shellfish Toxin Content and Profile of Alexandrium catenella (Dinophyceae) Isolated from the Mediterranean Sea. Marine Drugs, 2013, 11, 1583-1601.	2.2	57
14	Beta-N-Methylamino-l-Alanine: LC-MS/MS Optimization, Screening of Cyanobacterial Strains and Occurrence in Shellfish from Thau, a French Mediterranean Lagoon. Marine Drugs, 2014, 12, 5441-5467.	2.2	56
15	Systematic detection of BMAA (\hat{l}^2 -N-methylamino-l-alanine) and DAB (2,4-diaminobutyric acid) in mollusks collected in shellfish production areas along the French coasts. Toxicon, 2016, 110, 35-46.	0.8	54
16	Tectus niloticus (Tegulidae, Gastropod) as a Novel Vector of Ciguatera Poisoning: Detection of Pacific Ciguatoxins in Toxic Samples from Nuku Hiva Island (French Polynesia). Toxins, 2018, 10, 2.	1.5	54
17	Evidence of the bioaccumulation of ciguatoxins in giant clams (Tridacna maxima) exposed to Gambierdiscus spp. cells. Harmful Algae, 2016, 57, 78-87.	2.2	53
18	Exposure to toxic Alexandrium minutum activates the detoxifying and antioxidant systems in gills of the oyster Crassostrea gigas. Harmful Algae, 2015, 48, 55-62.	2.2	45

#	Article	IF	CITATIONS
19	Viability, growth and toxicity ofAlexandrium catenellaandAlexandrium minutum(Dinophyceae) following ingestion and gut passage in the oysterCrassostrea gigas. Aquatic Living Resources, 2007, 20, 51-57.	0.5	43
20	Three decades of data on phytoplankton and phycotoxins on the French coast: Lessons from REPHY and REPHYTOX. Harmful Algae, 2021, 102, 101733.	2.2	43
21	Paralytic shellfish poison outbreaks in the Penz \tilde{A} © estuary: Environmental factors affecting toxin uptake in the oyster, Crassostrea gigas. Aquatic Living Resources, 2004, 17, 207-214.	0.5	42
22	Toxicological Investigations on the Sea Urchin Tripneustes gratilla (Toxopneustidae, Echinoid) from Anaho Bay (Nuku Hiva, French Polynesia): Evidence for the Presence of Pacific Ciguatoxins. Marine Drugs, 2018, 16, 122.	2.2	42
23	Demonstrated transfer of cyanobacteria and cyanotoxins along a freshwater-marine continuum in France. Harmful Algae, 2019, 87, 101639.	2.2	38
24	Effects of Organic and Inorganic Nitrogen on the Growth and Production of Domoic Acid by Pseudo-nitzschia multiseries and P. australis (Bacillariophyceae) in Culture. Marine Drugs, 2015, 13, 7067-7086.	2.2	36
25	Quantitative analysis of azaspiracids in Azadinium spinosum cultures. Analytical and Bioanalytical Chemistry, 2012, 403, 833-846.	1.9	35
26	Extended evaluation of polymeric and lipophilic sorbents for passive sampling of marine toxins. Toxicon, 2014, 91, 57-68.	0.8	34
27	Okadaic acid and PP2A cellular immunolocalization in Prorocentrum lima (Dinophyceae). Phycologia, 1999, 38, 41-46.	0.6	32
28	New insights on the species-specific allelopathic interactions between macrophytes and marine HAB dinoflagellates. PLoS ONE, 2017, 12, e0187963.	1.1	32
29	Modelling the accumulation of PSP toxins in Thau Lagoon oysters (Crassostrea gigas) from trials using mixed cultures ofAlexandrium catenellaandThalassiosira weissflogii. Aquatic Living Resources, 2007, 20, 59-67.	0.5	31
30	Exposure to the Neurotoxic Dinoflagellate, Alexandrium catenella, Induces Apoptosis of the Hemocytes of the Oyster, Crassostrea gigas. Marine Drugs, 2013, 11, 4799-4814.	2.2	30
31	Ostreopsis lenticularis Y. Fukuyo (Dinophyceae, Gonyaulacales) from French Polynesia (South Pacific) Tj ETQq1 1	. 0 <u>.78</u> 431	4 rgBT /Overl
32	Detoxification of Pacific oyster <i>Crassostrea gigas</i> fed on diets of <i>Skeletonema costatum</i> with and without silt, following PSP contamination by <i>Alexandrium minutum</i> Aquatic Living Resources, 2008, 21, 13-20.	0.5	28
33	Passive Sampling and High Resolution Mass Spectrometry for Chemical Profiling of French Coastal Areas with a Focus on Marine Biotoxins. Environmental Science & Environmental Science & 2016, 50, 8522-8529.	4.6	28
34	Interactions between Scientists, Managers and Policy Makers in the Framework of the French MediOs Project on ⟨i⟩Ostreopsis⟨ i⟩(2008–2010). Cryptogamie, Algologie, 2012, 33, 137-142.	0.3	27
35	Physiological and Metabolic Responses of Freshwater and Brackish-Water Strains of Microcystis aeruginosa Acclimated to a Salinity Gradient: Insight into Salt Tolerance. Applied and Environmental Microbiology, 2019, 85, .	1.4	27
36	Toxin content of Ostreopsis cf. ovata depends on bloom phases, depth and macroalgal substrate in the NW Mediterranean Sea. Harmful Algae, 2020, 92, 101727.	2.2	23

#	Article	IF	CITATIONS
37	Effects of copper and butyltin compounds on the growth, photosynthetic activity and toxin production of two HAB dinoflagellates: The planktonic Alexandrium catenella and the benthic Ostreopsis cf. ovata. Aquatic Toxicology, 2018, 196, 154-167.	1.9	22
38	Oyster transcriptome response to Alexandrium exposure is related to saxitoxin load and characterized by disrupted digestion, energy balance, and calcium and sodium signaling. Aquatic Toxicology, 2018, 199, 127-137.	1.9	19
39	Monitoring the Emergence of Algal Toxins in Shellfish: First Report on Detection of Brevetoxins in French Mediterranean Mussels. Marine Drugs, 2021, 19, 393.	2.2	19
40	Influence of sudden salinity variation on the physiology and domoic acid production by two strains of ⟨i⟩Pseudoâ€nitzschia australis⟨/i⟩. Journal of Phycology, 2019, 55, 186-195.	1.0	18
41	Alexandrium ostenfeldii growth and spirolide production in batch culture and photobioreactor. Harmful Algae, 2011, 10, 794-803.	2.2	17
42	<i>Ostreopsis</i> cf. <i>ovata</i> ii>in the French Mediterranean Coast: Molecular Characterisation and Toxin Profile. Cryptogamie, Algologie, 2012, 33, 89-98.	0.3	17
43	Spirolide uptake and detoxification by Crassostrea gigas exposed to the toxic dinoflagellate Alexandrium ostenfeldii. Aquaculture, 2012, 358-359, 108-115.	1.7	17
44	Exposure to the Paralytic Shellfish Toxin Producer Alexandrium catenella Increases the Susceptibility of the Oyster Crassostrea gigas to Pathogenic Vibrios. Toxins, 2016, 8, 24.	1.5	17
45	Acclimation of the Marine Diatom <i>Pseudoâ€nitzschia australis</i> to Different Salinity Conditions: Effects on Growth, Photosynthetic Activity, and Domoic Acid Content ¹ . Journal of Phycology, 2020, 56, 97-109.	1.0	17
46	Paralytic Toxins Accumulation and Tissue Expression of \hat{I}_{\pm} -Amylase and Lipase Genes in the Pacific Oyster Crassostrea gigas Fed with the Neurotoxic Dinoflagellate Alexandrium catenella. Marine Drugs, 2012, 10, 2519-2534.	2,2	15
47	Microalgae and Toxins. , 2018, , 263-305.		15
48	Tissue Distribution and Elimination of Ciguatoxins in Tridacna maxima (Tridacnidae, Bivalvia) Fed Gambierdiscus polynesiensis. Toxins, 2018, 10, 189.	1.5	15
49	Salt Shock Responses of Microcystis Revealed through Physiological, Transcript, and Metabolomic Analyses. Toxins, 2020, 12, 192.	1.5	15
50	Guidance Level for Brevetoxins in French Shellfish. Marine Drugs, 2021, 19, 520.	2.2	15
51	Chemodiversity of Brevetoxins and Other Potentially Toxic Metabolites Produced by Karenia spp. and Their Metabolic Products in Marine Organisms. Marine Drugs, 2021, 19, 656.	2.2	15
52	Molecular Characterization of Voltage-Gated Sodium Channels and Their Relations with Paralytic Shellfish Toxin Bioaccumulation in the Pacific Oyster Crassostrea gigas. Marine Drugs, 2017, 15, 21.	2.2	13
53	First Characterization of Ostreopsis cf. ovata (Dinophyceae) and Detection of Ovatoxins during a Multispecific and Toxic Ostreopsis Bloom on French Atlantic Coast. Marine Drugs, 2022, 20, 461.	2.2	13
54	A Feedback Mechanism to Control Apoptosis Occurs in the Digestive Gland of the Oyster Crassostrea gigas Exposed to the Paralytic Shellfish Toxins Producer Alexandrium catenella. Marine Drugs, 2014, 12, 5035-5054.	2.2	12

#	Article	IF	CITATIONS
55	Modelling paralytic shellfish toxins (PST) accumulation in Crassostrea gigas by using Dynamic Energy Budgets (DEB). Journal of Sea Research, 2019, 143, 152-164.	0.6	12
56	Taxonomy and toxicity of a bloom-forming Ostreopsis species (Dinophyceae, Gonyaulacales) in Tahiti island (South Pacific Ocean): one step further towards resolving the identity of O. siamensis Harmful Algae, 2020, 98, 101888.	2.2	12
57	Combined effects of temperature and light intensity on growth, metabolome and ovatoxin content of a Mediterranean Ostreopsis cf. ovata strain. Harmful Algae, 2021, 106, 102060.	2.2	6
58	Winter accumulation of paralytic shellfish toxins in digestive glands of mussels from Arcachon and Toulon (France) without detectable toxic plankton species revealed by interference in the mouse bioassay for lipophilic toxins. Natural Toxins, 1999, 7, 271-277.	1.0	5
59	Toxicity of palytoxin, purified ovatoxin-a, ovatoxin-d and extracts of Ostreopsis cf. ovata on the Caco-2 intestinal barrier model. Environmental Toxicology and Pharmacology, 2022, 94, 103909.	2.0	3
60	Winter accumulation of paralytic shellfish toxins in digestive glands of mussels from Arcachon and Toulon (France) without detectable toxic plankton species revealed by interference in the mouse bioassay for lipophilic toxins. Natural Toxins, 1999, 7, 271-277.	1.0	1