

Seho Park

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1480071/seho-park-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140 papers	1,826 citations	24 h-index	35 g-index
149 ext. papers	2,199 ext. citations	3.8 avg, IF	4.62 L-index

#	Paper	IF	Citations
140	Ki-67 and breast cancer prognosis: does it matter if Ki-67 level is examined using preoperative biopsy or postoperative specimen?. <i>Breast Cancer Research and Treatment</i> , 2022 , 192, 343	4.4	4
139	Abstract P2-11-13: Distinctive clinicopathological features of premenopausal women with intermediate 21-gene recurrence score. <i>Cancer Research</i> , 2022 , 82, P2-11-13-P2-11-13	10.1	
138	Abstract OT1-04-02: The NAUTILUS trial (No Axillary sUrgical Treatment In clinically Lymph node negative patients after UltraSonography): A prospective multicenter randomized phase III trial (NCT04303715). <i>Cancer Research</i> , 2022 , 82, OT1-04-02-OT1-04-02	10.1	
137	Abstract P1-17-09: Efficacy of limited dose modifications for palbociclib-related grade 3 neutropenia in hormone receptor positive metastatic breast cancer. <i>Cancer Research</i> , 2022 , 82, P1-17-09-P1-17-09	10.1	
136	Abstract P2-08-13: The prognostic significance of Ki67 and progesterone receptor; could they really have prognostic importance in early breast cancer with estrogen receptor positive, HER-2 negative?. <i>Cancer Research</i> , 2022 , 82, P2-08-13-P2-08-13	10.1	
135	No axillary surgical treatment for lymph node-negative patients after ultra-sonography [NAUTILUS]: protocol of a prospective randomized clinical trial.. <i>BMC Cancer</i> , 2022 , 22, 189	4.8	1
134	Abstract P4-07-14: Long-term prognosis of minimal residual disease in breast cancer patients with breast or nodal pathologic complete response (pCR) after neoadjuvant chemotherapy (NAC). <i>Cancer Research</i> , 2022 , 82, P4-07-14-P4-07-14	10.1	
133	Abstract PD6-07: Whole genome sequencing-based circulating tumor DNA profiling of metastatic breast cancer patients for molecular characterization and therapy response prediction. <i>Cancer Research</i> , 2022 , 82, PD6-07-PD6-07	10.1	
132	Abstract P1-07-03: Clinical treatment score post-5 years (CTS5) and the prediction of survival benefit from extended endocrine therapy for breast cancer patients under and over 50 years of age. <i>Cancer Research</i> , 2022 , 82, P1-07-03-P1-07-03	10.1	
131	Abstract P2-13-14: Pattern of recurrence after pathologic complete response after neoadjuvant chemotherapy in patients with early HER2-positive breast cancer: Real-world evidence. <i>Cancer Research</i> , 2022 , 82, P2-13-14-P2-13-14	10.1	
130	RT-induced dynamic changes in the lymphocyte-to-monocyte ratio in patients with breast cancer indicate poor prognosis.. <i>Breast Cancer Research and Treatment</i> , 2022 , 193, 637	4.4	0
129	Relationship Between Breast and Axillary Pathologic Complete Response According to Clinical Nodal Stage: A Nationwide Study From Korean Breast Cancer Society.. <i>Journal of Breast Cancer</i> , 2022 , 25, 94-105	3	0
128	Genomic landscape of extraordinary responses in metastatic breast cancer. <i>Communications Biology</i> , 2021 , 4, 449	6.7	1
127	On-treatment derived neutrophil-to-lymphocyte ratio and response to palbociclib and letrozole: Analysis of a multicenter retrospective cohort and the PALOMA-2 study.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 1066-1066	2.2	
126	Clinicopathological Features and Prognosis of Lobular Carcinoma In Situ. <i>Journal of Breast Disease</i> , 2021 , 9, 10-15	0.3	
125	Real-World Clinical Outcomes of Biosimilar Trastuzumab (CT-P6) in HER2-Positive Early-Stage and Metastatic Breast Cancer. <i>Frontiers in Oncology</i> , 2021 , 11, 689587	5.3	3
124	Preoperative Routine Laboratory Markers for Predicting Postoperative Recurrence and Death in Patients with Breast Cancer. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1

123	Time-Sequencing of the Neutrophil-to-Lymphocyte Ratio to Predict Prognosis of Triple-Negative Breast Cancer. <i>Cancers</i> , 2021 , 13,	6.6	2
122	Change in Shoulder Function in the Early Recovery Phase after Breast Cancer Surgery: A Prospective Observational Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
121	Survival Rates of Breast Cancer Patients Aged 40 to 49 Years according to Detection Modality in Korea: Screening Ultrasound versus Mammography. <i>Korean Journal of Radiology</i> , 2021 , 22, 159-167	6.9	2
120	Axillary response according to neoadjuvant single or dual human epidermal growth factor receptor 2 (HER2) blockade in clinically node-positive, HER2-positive breast cancer. <i>International Journal of Cancer</i> , 2021 , 149, 1585-1592	7.5	2
119	Effects of a partnership-based, needs-tailored self-management support intervention for post-treatment breast cancer survivors: A randomized controlled trial. <i>Psycho-Oncology</i> , 2021 ,	3.9	1
118	Design of a randomized controlled trial of a partnership-based, needs-tailored self-management support intervention for post-treatment breast cancer survivors. <i>BMC Cancer</i> , 2020 , 20, 367	4.8	4
117	Palbociclib use with grade 3 neutropenia in hormone receptor-positive metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 183, 107-116	4.4	0
116	Dietary advanced glycation end products and the risk of postmenopausal breast cancer in the National Institutes of Health-AARP Diet and Health Study. <i>Cancer</i> , 2020 , 126, 2648-2657	6.4	12
115	Validating Breast Cancer Risk Prediction Models in the Korean Cancer Prevention Study-II Biobank. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 1271-1277	4	2
114	Optimal treatment of pseudoangiomatous stromal hyperplasia of the breast. <i>Asian Journal of Surgery</i> , 2020 , 43, 735-741	1.6	5
113	Intermediate HER2 expression is associated with poor prognosis in estrogen receptor-positive breast cancer patients aged 55 years and older. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 687-697	4.4	6
112	Radiotherapy-Induced High Neutrophil-to-Lymphocyte Ratio is a Negative Prognostic Factor in Patients with Breast Cancer. <i>Cancers</i> , 2020 , 12,	6.6	9
111	Prognostic value of neutrophil-to-lymphocyte ratio in human epidermal growth factor receptor 2-negative breast cancer patients who received neoadjuvant chemotherapy. <i>Scientific Reports</i> , 2020 , 10, 13078	4.9	8
110	Conserving the lymphatics from the arm using fluorescence imaging in patients with breast cancer at high risk of postoperative lymphedema: a pilot study. <i>Gland Surgery</i> , 2020 , 9, 629-636	2.2	2
109	Validation of Clinical Treatment Score post-5 years (CTS5) risk stratification in premenopausal breast cancer patients and Ki-67 labelling index. <i>Scientific Reports</i> , 2020 , 10, 16850	4.9	6
108	Post-Operative Complications and Nipple Necrosis Rates Between Conventional and Robotic Nipple-Sparing Mastectomy. <i>Frontiers in Oncology</i> , 2020 , 10, 594388	5.3	6
107	Effect of primary tumor resection on overall survival in patients with stage IV breast cancer. <i>Breast Journal</i> , 2019 , 25, 908-915	1.2	4
106	Childbirth in young Korean women with previously treated breast cancer: The SMARTSHIP study. <i>Breast Cancer Research and Treatment</i> , 2019 , 176, 419-427	4.4	6

105	Single dose of dexamethasone is not associated with postoperative recurrence and mortality in breast cancer patients: a propensity-matched cohort study. <i>BMC Cancer</i> , 2019 , 19, 251	4.8	15
104	Association of white blood cell count with breast cancer burden varies according to menopausal status, body mass index, and hormone receptor status: a case-control study. <i>Scientific Reports</i> , 2019 , 9, 5762	4.9	6
103	Early experiences with robot-assisted prosthetic breast reconstruction. <i>Archives of Plastic Surgery</i> , 2019 , 46, 79-83	1.6	18
102	Retrospectively validating the results of the ACOSOG Z0011 trial in a large Asian Z0011-eligible cohort. <i>Breast Cancer Research and Treatment</i> , 2019 , 175, 203-215	4.4	7
101	Adjuvant chemotherapy and survival among patients 70 years of age and younger with node-negative breast cancer and the 21-gene recurrence score of 26-30. <i>Breast Cancer Research</i> , 2019 , 21, 110	8.3	8
100	Tramadol use is associated with enhanced postoperative outcomes in breast cancer patients: a retrospective clinical study with in vitro confirmation. <i>British Journal of Anaesthesia</i> , 2019 , 123, 865-876	5.4	15
99	Postoperative Cancer Surveillance Following Oncoplastic Surgery with Latissimus Dorsi Flap: a Matched Case-Control Study. <i>Annals of Surgical Oncology</i> , 2019 , 26, 4681-4691	3.1	1
98	Aromatase inhibitor use is a risk factor of carotid plaque presence in endocrine-responsive breast cancer patients. <i>Korean Journal of Internal Medicine</i> , 2019 , 34, 579-587	2.5	1
97	A Phase II Study to Evaluate the Safety and Efficacy of Pegteograstim in Korean Breast Cancer Patients Receiving Dose-Dense Doxorubicin/Cyclophosphamide. <i>Cancer Research and Treatment</i> , 2019 , 51, 812-818	5.2	2
96	Is the frozen section examination for sentinel lymph node necessary in early breast cancer patients?. <i>Annals of Surgical Treatment and Research</i> , 2019 , 97, 49-57	2	11
95	Establishment of chemosensitivity tests in triple-negative and BRCA-mutated breast cancer patient-derived xenograft models. <i>PLoS ONE</i> , 2019 , 14, e0225082	3.7	6
94	Prevalence of breast cancer-related risk factors in underweight premenopausal women: the Korea National Health and Nutrition Examination Survey IV-VI. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 515-524	4.4	4
93	Establishment of chemosensitivity tests in triple-negative and BRCA-mutated breast cancer patient-derived xenograft models 2019 , 14, e0225082		
92	Establishment of chemosensitivity tests in triple-negative and BRCA-mutated breast cancer patient-derived xenograft models 2019 , 14, e0225082		
91	Establishment of chemosensitivity tests in triple-negative and BRCA-mutated breast cancer patient-derived xenograft models 2019 , 14, e0225082		
90	Establishment of chemosensitivity tests in triple-negative and BRCA-mutated breast cancer patient-derived xenograft models 2019 , 14, e0225082		
89	Can Biannual Ultrasound Surveillance Detect Smaller Second Cancers or Detect Cancers Earlier in Patients with Breast Cancer History?. <i>Ultrasound in Medicine and Biology</i> , 2018 , 44, 1355-1363	3.5	2
88	Oncologic Safety of Gonadotropin-Releasing Hormone Agonist for Ovarian Function Protection During Breast Cancer Chemotherapy. <i>Clinical Breast Cancer</i> , 2018 , 18, e1165-e1172	3	1

87	Locoregional Treatment of the Primary Tumor in Patients With De Novo Stage IV Breast Cancer: A Radiation Oncologist's Perspective. <i>Clinical Breast Cancer</i> , 2018 , 18, e167-e178	3	20
86	Impact of Micrometastatic Axillary Nodes on Survival of Breast Cancer Patients with Tumors ≥ 1 cm. <i>World Journal of Surgery</i> , 2018 , 42, 3969-3978	3.3	4
85	Feasibility of Charcoal Tattooing of Cytology-Proven Metastatic Axillary Lymph Node at Diagnosis and Sentinel Lymph Node Biopsy after Neoadjuvant Chemotherapy in Breast Cancer Patients. <i>Cancer Research and Treatment</i> , 2018 , 50, 801-812	5.2	45
84	Relationship between resting heart rate and metabolic risk factors in breast cancer patients. <i>Clinica Chimica Acta</i> , 2018 , 486, 104-109	6.2	7
83	Association between Changes in Serum 25-Hydroxyvitamin D Levels and Survival in Patients with Breast Cancer Receiving Neoadjuvant Chemotherapy. <i>Journal of Breast Cancer</i> , 2018 , 21, 134-141	3	5
82	Role of adding ovarian function suppression to tamoxifen in young women with hormone-sensitive breast cancer who remain premenopausal or resume menstruation after chemotherapy: The ASTRRA study.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 502-502	2.2	13
81	External validation of IBTR! 2.0 nomogram for prediction of ipsilateral breast tumor recurrence. <i>Radiation Oncology Journal</i> , 2018 , 36, 139-146	2.5	4
80	CD44/CD24 and aldehyde dehydrogenase 1 in estrogen receptor-positive early breast cancer treated with tamoxifen: CD24 positivity is a poor prognosticator. <i>Oncotarget</i> , 2018 , 9, 2622-2630	3.3	10
79	Gasless Robot-Assisted Nipple-Sparing Mastectomy: A Case Report. <i>Journal of Breast Cancer</i> , 2018 , 21, 334-338	3	21
78	Effects of hormone receptor status on the durable response of trastuzumab-based therapy in metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2017 , 163, 255-262	4.4	3
77	Preoperative prediction of the size of pure ductal carcinoma in situ using three imaging modalities as compared to histopathological size: does magnetic resonance imaging add value?. <i>Breast Cancer Research and Treatment</i> , 2017 , 164, 437-444	4.4	6
76	Chemotherapy-induced irreversible alopecia in early breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2017 , 163, 527-533	4.4	11
75	The clinical implication of the number of lymph nodes harvested during sentinel lymph node biopsy and its effects on survival outcome in patients with node-negative breast cancer. <i>American Journal of Surgery</i> , 2017 , 214, 726-732	2.7	12
74	Meeting Highlights: The Second Consensus Conference for Breast Cancer Treatment in Korea. <i>Journal of Breast Cancer</i> , 2017 , 20, 228-233	3	2
73	Effects of systemic lidocaine versus magnesium administration on postoperative functional recovery and chronic pain in patients undergoing breast cancer surgery: A prospective, randomized, double-blind, comparative clinical trial. <i>PLoS ONE</i> , 2017 , 12, e0173026	3.7	23
72	Detection of Circulating Tumor Cells in Breast Cancer Patients Using Cytokeratin-19 Real-Time RT-PCR. <i>Yonsei Medical Journal</i> , 2017 , 58, 19-26	3	13
71	Factors predictive of occult nipple-areolar complex involvement in patients with carcinoma in situ of the breast. <i>Journal of Surgical Oncology</i> , 2017 , 116, 1046-1055	2.8	6
70	Outcome following sentinel lymph node biopsy-guided decisions in breast cancer patients with conversion from positive to negative axillary lymph nodes after neoadjuvant chemotherapy. <i>Breast Cancer Research and Treatment</i> , 2017 , 166, 473-480	4.4	21

69	Different patterns in the risk of newly developed fatty liver and lipid changes with tamoxifen versus aromatase inhibitors in postmenopausal women with early breast cancer: A propensity score-matched cohort study. <i>European Journal of Cancer</i> , 2017 , 82, 103-114	7.5	17
68	Does the type of anesthesia really affect the recurrence-free survival after breast cancer surgery?. <i>Oncotarget</i> , 2017 , 8, 90477-90487	3.3	44
67	The Effects of Perioperative Anesthesia and Analgesia on Immune Function in Patients Undergoing Breast Cancer Resection: A Prospective Randomized Study. <i>International Journal of Medical Sciences</i> , 2017 , 14, 970-976	3.7	49
66	Next-generation sequencing of in breast cancer patients: potential effects on clinical decision-making using rapid, high-accuracy genetic results. <i>Annals of Surgical Treatment and Research</i> , 2017 , 92, 331-339	2	16
65	Comparison of characteristics in patients with both thyroid and breast cancer: Based on order of incidence. <i>Korean Journal of Clinical Oncology</i> , 2017 , 13, 1-9	0.1	5
64	Oncologic safety of breast-conserving surgery in breast cancer patients under the age of 35. <i>Korean Journal of Clinical Oncology</i> , 2017 , 13, 32-38	0.1	1
63	Lack of both androgen receptor and forkhead box A1 (FOXA1) expression is a poor prognostic factor in estrogen receptor-positive breast cancers. <i>Oncotarget</i> , 2017 , 8, 82940-82955	3.3	6
62	Resting heart rate as a prognostic factor for mortality in patients with breast cancer. <i>Breast Cancer Research and Treatment</i> , 2016 , 159, 375-84	4.4	27
61	Trends in the Application of Postmastectomy Radiotherapy for Breast Cancer With 1 to 3 Positive Axillary Nodes and Tumors ≤ cm in the Modern Treatment Era: A Retrospective Korean Breast Cancer Society Report. <i>Medicine (United States)</i> , 2016 , 95, e3592	1.8	4
60	No effect of tumor-infiltrating lymphocytes (TILs) on prognosis in patients with early triple-negative breast cancer: Validation of recommendations by the International TILs Working Group 2014. <i>Journal of Surgical Oncology</i> , 2016 , 114, 17-21	2.8	8
59	Association Between Insulin Resistance and Luminal B Subtype Breast Cancer in Postmenopausal Women. <i>Medicine (United States)</i> , 2016 , 95, e2825	1.8	18
58	Magnetic Resonance Imaging after Completion of Neoadjuvant Chemotherapy Can Accurately Discriminate between No Residual Carcinoma and Residual Ductal Carcinoma In Situ in Patients with Triple-Negative Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0149347	3.7	12
57	Risk Factors Associated with Discordant Ki-67 Levels between Preoperative Biopsy and Postoperative Surgical Specimens in Breast Cancers. <i>PLoS ONE</i> , 2016 , 11, e0151054	3.7	8
56	ZNF224, Krüppel like zinc finger protein, induces cell growth and apoptosis-resistance by down-regulation of p21 and p53 via miR-663a. <i>Oncotarget</i> , 2016 , 7, 31177-90	3.3	22
55	Comparisons of Oncologic Outcomes between Triple-Negative Breast Cancer (TNBC) and Non-TNBC among Patients Treated with Breast-Conserving Therapy. <i>Yonsei Medical Journal</i> , 2016 , 57, 1192-8	3	15
54	Incidence of Febrile Neutropenia in Korean Female Breast Cancer Patients Receiving Preoperative or Postoperative Doxorubicin/Cyclophosphamide Followed by Docetaxel Chemotherapy. <i>Journal of Breast Cancer</i> , 2016 , 19, 76-82	3	13
53	Preoperative Magnetic Resonance Imaging and Survival Outcomes in T1-2 Breast Cancer Patients Who Receive Breast-Conserving Therapy. <i>Journal of Breast Cancer</i> , 2016 , 19, 423-428	3	9
52	Expression of T-Lymphocyte Markers in Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer. <i>Journal of Breast Cancer</i> , 2016 , 19, 385-393	3	2

51	Intratumoral Agreement of High-Resolution Magic Angle Spinning Magnetic Resonance Spectroscopic Profiles in the Metabolic Characterization of Breast Cancer. <i>Medicine (United States)</i> , 2016 , 95, e3398	1.8	13
50	The role of the addition of ovarian suppression to tamoxifen in young women with hormone-sensitive breast cancer who remain premenopausal or regain menstruation after chemotherapy (ASTRRA): study protocol for a randomized controlled trial and progress. <i>BMC Cancer</i> , 2016 , 16, 319	4.8	16
49	Serum 25-hydroxyvitamin D deficiency and increased risk of breast cancer among Korean women: a case-control study. <i>Breast Cancer Research and Treatment</i> , 2015 , 152, 147-154	4.4	26
48	Prognostic Nomogram for Prediction of Axillary Pathologic Complete Response After Neoadjuvant Chemotherapy in Cytologically Proven Node-Positive Breast Cancer. <i>Medicine (United States)</i> , 2015 , 94, e1720	1.8	36
47	The effect of oncologists' exercise recommendations on the level of exercise and quality of life in survivors of breast and colorectal cancer: A randomized controlled trial. <i>Cancer</i> , 2015 , 121, 2740-8	6.4	82
46	Evaluation of the Survival Benefit of Different Chemotherapy Regimens in Patients with T1-2N0 Triple-Negative Breast Cancer. <i>Journal of Breast Cancer</i> , 2015 , 18, 271-8	3	11
45	Local control of brain metastasis: treatment outcome of focal brain treatments in relation to subtypes. <i>Journal of Breast Cancer</i> , 2015 , 18, 29-35	3	6
44	Characteristics and Survival of Breast Cancer Patients with Multiple Synchronous or Metachronous Primary Cancers. <i>Yonsei Medical Journal</i> , 2015 , 56, 1213-20	3	23
43	Anaplastic lymphoma kinase gene copy number gain in inflammatory breast cancer (IBC): prevalence, clinicopathologic features and prognostic implication. <i>PLoS ONE</i> , 2015 , 10, e0120320	3.7	12
42	Evaluation with 3.0-T MR imaging: predicting the pathological response of triple-negative breast cancer treated with anthracycline and taxane neoadjuvant chemotherapy. <i>Acta Radiologica</i> , 2015 , 56, 1069-77	2	11
41	The long term outcome of clinical trial-based treatment comparing standard treatment for metastatic breast cancer.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 11106-11106	2.2	
40	Expression of growth factor receptor family before and after targeted therapy in human epidermal growth factor receptor-2 positive breast cancer tissues. <i>Korean Journal of Clinical Oncology</i> , 2015 , 11, 12-19	0.1	1
39	Human leukocyte antigen-G (HLA-G) polymorphism and expression in breast cancer patients. <i>PLoS ONE</i> , 2014 , 9, e98284	3.7	48
38	Subcellular localization of Mdm2 expression and prognosis of breast cancer. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 842-51	4.2	19
37	Clinical Outcome of Immediate Breast Reconstruction after Mastectomy for Breast Cancer. <i>Journal of Breast Disease</i> , 2014 , 2, 44-50	0.3	2
36	Elevated levels of serum tumor markers CA 15-3 and CEA are prognostic factors for diagnosis of metastatic breast cancers. <i>Breast Cancer Research and Treatment</i> , 2013 , 141, 477-84	4.4	30
35	S-1 combined with docetaxel following doxorubicin plus cyclophosphamide as neoadjuvant therapy in breast cancer: phase II trial. <i>BMC Cancer</i> , 2013 , 13, 583	4.8	6
34	Lack of either estrogen or progesterone receptor expression is associated with poor survival outcome among luminal A breast cancer subtype. <i>Annals of Surgical Oncology</i> , 2013 , 20, 1505-13	3.1	26

33	Prolonged clinical benefit from the maintenance hormone therapy in patients with metastatic breast cancer. <i>Breast</i> , 2013 , 22, 1205-9	3.6	5
32	A nomogram for predicting underestimation of invasiveness in ductal carcinoma in situ diagnosed by preoperative needle biopsy. <i>Breast</i> , 2013 , 22, 869-73	3.6	27
31	Sentinel lymph node biopsy after neoadjuvant chemotherapy in patients with cytologically proven node-positive breast cancer at diagnosis. <i>Annals of Surgical Oncology</i> , 2013 , 20, 2858-65	3.1	51
30	Risk predictors of underestimation and the need for sentinel node biopsy in patients diagnosed with ductal carcinoma in situ by preoperative needle biopsy. <i>Journal of Surgical Oncology</i> , 2013 , 107, 388-92	2.8	44
29	Oncologic safety of breast-conserving surgery compared to mastectomy in patients receiving neoadjuvant chemotherapy for locally advanced breast cancer. <i>Journal of Surgical Oncology</i> , 2013 , 108, 531-6	2.8	45
28	Efficacy of a combined alendronate and calcitriol agent (Maxmarvil®) in Korean postmenopausal women with early breast cancer receiving aromatase inhibitor: a double-blind, randomized, placebo-controlled study. <i>Endocrine Journal</i> , 2013 , 60, 167-72	2.9	17
27	Characteristics and outcomes according to molecular subtypes of breast cancer as classified by a panel of four biomarkers using immunohistochemistry. <i>Breast</i> , 2012 , 21, 50-7	3.6	141
26	Breast cancers presenting luminal B subtype features show higher discordant human epidermal growth factor receptor 2 results between immunohistochemistry and fluorescence in situ hybridization. <i>Cancer</i> , 2012 , 118, 914-23	6.4	12
25	Higher expression of androgen receptor is a significant predictor for better endocrine-responsiveness in estrogen receptor-positive breast cancers. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 311-20	4.4	30
24	Cyclooxygenase-2 expression in proliferative Ki-67-positive breast cancers is associated with poor outcomes. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 741-51	4.4	23
23	Homeodomain-interacting protein kinase 1 (HIPK1) expression in breast cancer tissues. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 1138-45	2.8	3
22	Overexpression of class III beta tubulin and amplified HER2 gene predict good response to paclitaxel and trastuzumab therapy. <i>PLoS ONE</i> , 2012 , 7, e45127	3.7	13
21	Association between genetic polymorphisms of CYP2D6 and outcomes in breast cancer patients with tamoxifen treatment. <i>Journal of Korean Medical Science</i> , 2011 , 26, 1007-13	4.7	28
20	Clinical Significance of Age at the Time of Diagnosis among Young Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2011 , 14, 314-21	3	8
19	The impact of local and regional recurrence on distant metastasis and survival in patients treated with breast conservation therapy. <i>Journal of Breast Cancer</i> , 2011 , 14, 191-7	3	12
18	The feasibility of endoscopy-assisted breast conservation surgery for patients with early breast cancer. <i>Journal of Breast Cancer</i> , 2011 , 14, 52-7	3	22
17	How many sentinel lymph nodes are enough for accurate axillary staging in t1-2 breast cancer?. <i>Journal of Breast Cancer</i> , 2011 , 14, 296-300	3	45
16	Dermatofibrosarcoma protuberans arising on the skin of the breast. <i>Breast Journal</i> , 2011 , 17, 93-5	1.2	2

15	Clinical significance of progesterone receptor and HER2 status in estrogen receptor-positive, operable breast cancer with adjuvant tamoxifen. <i>Journal of Cancer Research and Clinical Oncology</i> , 2011 , 137, 1123-30	4.9	24
14	Factors influencing the outcome of breast cancer patients with 10 or more metastasized axillary lymph nodes. <i>International Journal of Clinical Oncology</i> , 2011 , 16, 473-81	4.2	11
13	The impact of caveolin protein expression in tumor stroma on prognosis of breast cancer. <i>Tumor Biology</i> , 2011 , 32, 787-99	2.9	40
12	Comparison of treatment outcome between breast-conservation surgery with radiation and total mastectomy without radiation in patients with one to three positive axillary lymph nodes. <i>International Journal of Radiation Oncology Biology Physics</i> , 2011 , 80, 1446-52	4	11
11	The impact of a focally positive resection margin on the local control in patients treated with breast-conserving therapy. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 600-8	2.8	19
10	Clinicopathologic features and outcomes of metaplastic breast carcinoma: comparison with invasive ductal carcinoma of the breast. <i>Yonsei Medical Journal</i> , 2010 , 51, 864-9	3	47
9	Clinicopathological characteristics of mucinous carcinoma of the breast in Korea: comparison with invasive ductal carcinoma-not otherwise specified. <i>Journal of Korean Medical Science</i> , 2010 , 25, 361-8	4.7	27
8	Paclitaxel combined with ifosfamide in anthracycline- and docetaxel-pretreated metastatic breast cancer: activity independence of prior docetaxel resistance. <i>Cancer Chemotherapy and Pharmacology</i> , 2010 , 66, 425-31	3.5	
7	Clinicopathological features of infiltrating lobular carcinomas comparing with infiltrating ductal carcinomas: a case control study. <i>World Journal of Surgical Oncology</i> , 2010 , 8, 34	3.4	34
6	Clinical Analysis of Medullary Carcinoma of the Breast. <i>Journal of Breast Cancer</i> , 2009 , 12, 47	3	2
5	The Prognostic Significance of Survivin Expression in Breast Cancer. <i>Journal of Breast Cancer</i> , 2009 , 12, 285	3	2
4	Clinicopathologic Characteristics of Apocrine Breast Carcinoma. [<i>Chapchi</i>] <i>Journal Taehan Oekwa Hakhoe</i> , 2009 , 77, 43		1
3	Changes of Coregulators, MAP Kinase Activity and p27/kip1 with Estrogen or Antiestrogen Treatment in Breast Cancer Cell Line. <i>Journal of Breast Cancer</i> , 2008 , 11, 56	3	
2	Clinicopathological Characteristics and Prognostic Factors of Papillary Carcinoma of the Breast. <i>Journal of Breast Cancer</i> , 2008 , 11, 77	3	
1	Clinicopathological characteristics of male breast cancer. <i>Yonsei Medical Journal</i> , 2008 , 49, 978-86	3	25