Yan-Jin Liu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1480870/publications.pdf

Version: 2024-02-01

1307594 1474206 9 169 7 9 citations g-index h-index papers 9 9 9 305 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Highly sensitive fluorogenic sensing of L-Cysteine in live cells using gelatin-stabilized gold nanoparticles decorated graphene nanosheets. Sensors and Actuators B: Chemical, 2018, 259, 339-346.	7.8	50
2	The methanol extract of Euonymus laxiflorus, Rubia lanceolata and Gardenia jasminoides inhibits xanthine oxidase and reduce serum uric acid level in rats. Food and Chemical Toxicology, 2014, 70, 179-184.	3.6	33
3	Body weight management effect of burdock (Arctium lappa L.) root is associated with the activation of AMP-activated protein kinase in human HepG2 cells. Food Chemistry, 2012, 134, 1320-1326.	8.2	23
4	Targeting \hat{I}^2 -tubulin/CCT- \hat{I}^2 complex induces apoptosis and suppresses migration and invasion of highly metastatic lung adenocarcinoma. Carcinogenesis, 2020, 41, 699-710.	2.8	17
5	Disrupting CCT-β : β-tubulin selectively kills CCT-β overexpressed cancer cells through MAPKs activation Cell Death and Disease, 2017, 8, e3052-e3052.	·6.3	15
6	CCT327 enhances TRAIL-induced apoptosis through the induction of death receptors and downregulation of cell survival proteins in TRAIL-resistant human leukemia cells. Oncology Reports, 2014, 32, 1257-1264.	2.6	11
7	Synergistic inhibition of leukemia WEHI-3 cell growth by arsenic trioxide and Hedyotis diffusa Willd extract in vitro and in vivo. Experimental and Therapeutic Medicine, 2017, 13, 3388-3396.	1.8	10
8	Hypolipidemic activity of Taraxacum mongolicum associated with the activation of AMP-activated protein kinase in human HepG2 cells. Food and Function, 2014, 5, 1755.	4.6	8
9	Econazoleâ€induced Ca ²⁺ fluxes and apoptosis in human oral cancer cells. Drug Development Research, 2010, 71, 240-248.	2.9	2