

# Imre Romics

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/1482564/imre-romics-publications-by-citations.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60  
papers

998  
citations

21  
h-index

29  
g-index

75  
ext. papers

1,172  
ext. citations

3.1  
avg, IF

3.41  
L-index

| #  | Paper  | IF   | Citations |
|----|--|------|-----------|
| 60 | Urinary incontinence and voiding dysfunction after radical retropubic prostatectomy (prospective urodynamic study). <i>Neurourology and Urodynamics</i> , <b>2006</b> , 25, 2-7  | 2.3  | 110       |
| 59 | Matrix metalloproteinase-7 as a marker of metastasis and predictor of poor survival in bladder cancer. <i>Cancer Science</i> , <b>2010</b> , 101, 1300-8   | 6.9  | 76        |
| 58 | Gene network and canonical pathway analysis in prostate cancer: a microarray study. <i>Experimental and Molecular Medicine</i> , <b>2008</b> , 40, 176-85  | 12.8 | 49        |
| 57 | Treatment and outcome of vaseline-induced sclerosing lipogranuloma of the penis. <i>Urology</i> , <b>2008</b> , 71, 1132-7   | 1.6  | 41        |
| 56 | Genotyping NAT2 with only two SNPs (rs1041983 and rs1801280) outperforms the tagging SNP rs1495741 and is equivalent to the conventional 7-SNP NAT2 genotype. <i>Pharmacogenetics and Genomics</i> , <b>2011</b> , 21, 673-8 | 1.9  | 38        |
| 55 | Long-term psychological and sexual outcomes of severe penile hypospadias repair. <i>Journal of Sexual Medicine</i> , <b>2011</b> , 8, 1529-39  | 1.1  | 37        |
| 54 | Elevated serum matrix metalloproteinase 7 levels predict poor prognosis after radical prostatectomy. <i>International Journal of Cancer</i> , <b>2011</b> , 128, 1486-92   | 7.5  | 36        |
| 53 | Serum endostatin levels correlate with enhanced extracellular matrix degradation and poor patients prognosis in bladder cancer. <i>International Journal of Cancer</i> , <b>2012</b> , 130, 2922-9                           | 7.5  | 34        |
| 52 | Analysis of risk factors for urinary incontinence after radical prostatectomy. <i>Urologia Internationalis</i> , <b>2007</b> , 78, 202-7   | 1.9  | 33        |
| 51 | Angiogenic switch of angiotensin-Tie2 system and its prognostic value in bladder cancer. <i>Clinical Cancer Research</i> , <b>2008</b> , 14, 8253-62   | 12.9 | 32        |
| 50 | Prognostic factors and survival of renal clear cell carcinoma patients with bone metastases. <i>Pathology and Oncology Research</i> , <b>2010</b> , 16, 29-38  | 2.6  | 30        |
| 49 | High insulin-like growth factor mRNA-binding protein 3 (IMP3) protein expression is associated with poor survival in muscle-invasive bladder cancer. <i>BJU International</i> , <b>2012</b> , 110, E308-17                   | 5.6  | 29        |
| 48 | Rs710521[A] on chromosome 3q28 close to TP63 is associated with increased urinary bladder cancer risk. <i>Archives of Toxicology</i> , <b>2010</b> , 84, 967-78  | 5.8  | 29        |
| 47 | The prognostic value of cadherin switch in bladder cancer. <i>Oncology Reports</i> , <b>2010</b> , 23, 1125-32   | 3.5  | 28        |
| 46 | Detection of bladder cancer from the urine using fluorescence in situ hybridization technique. <i>Pathology and Oncology Research</i> , <b>2007</b> , 13, 187-94   | 2.6  | 27        |
| 45 | Management of congenital penile curvature. <i>Journal of Urology</i> , <b>2008</b> , 179, 1495-8   | 2.5  | 26        |
| 44 | Rs11892031[A] on chromosome 2q37 in an intronic region of the UGT1A locus is associated with urinary bladder cancer risk. <i>Archives of Toxicology</i> , <b>2012</b> , 86, 1369-78  | 5.8  | 24        |

|    |  |      |    |
|----|--|------|----|
| 43 | Overexpression of CD24, c-myc and phospholipase 2A in prostate cancer tissue samples obtained by needle biopsy. <i>Pathology and Oncology Research</i> , <b>2009</b> , 15, 279-83  | 2.6  | 24 |
| 42 | Serum levels of angiogenic factors and their prognostic relevance in bladder cancer. <i>Pathology and Oncology Research</i> , <b>2009</b> , 15, 193-201  | 2.6  | 24 |
| 41 | Urinary matrix metalloproteinase-7 level is associated with the presence of metastasis in bladder cancer. <i>BJU International</i> , <b>2011</b> , 107, 1069-73  | 5.6  | 22 |
| 40 | beta-catenin expression and claudin expression pattern as prognostic factors of prostatic cancer progression. <i>BJU International</i> , <b>2010</b> , 105, 716-22   | 5.6  | 21 |
| 39 | Expression of claudins and their prognostic significance in noninvasive urothelial neoplasms of the human urinary bladder. <i>Journal of Histochemistry and Cytochemistry</i> , <b>2011</b> , 59, 932-41   | 3.4  | 21 |
| 38 | The application of adjuvant autologous intravesical macrophage cell therapy vs. BCG in non-muscle invasive bladder cancer: a multicenter, randomized trial. <i>Journal of Translational Medicine</i> , <b>2010</b> , 8, 54   | 8.5  | 15 |
| 37 | Serum Chromogranin A as a Complementary Marker for the Prediction of Prostate Cancer-Specific Survival. <i>Pathology and Oncology Research</i> , <b>2017</b> , 23, 643-650   | 2.6  | 13 |
| 36 | European Study of Radical Prostatectomy: time trends in Europe, 1993-2005. <i>BJU International</i> , <b>2007</b> , 100 Suppl 2, 22-5  | 5.6  | 12 |
| 35 | Urinary bladder cancer risk in relation to a single nucleotide polymorphism (rs2854744) in the insulin-like growth factor-binding protein-3 (IGFBP3) gene. <i>Archives of Toxicology</i> , <b>2012</b> , 86, 195-203   | 5.8  | 11 |
| 34 | A special terpene combination (Rowatinex <sup>®</sup> ) improves stone clearance after extracorporeal shockwave lithotripsy in urolithiasis patients: results of a placebo-controlled randomised controlled trial. <i>Urologia Internationalis</i> , <b>2011</b> , 86, 102-9 | 1.9  | 11 |
| 33 | Claudins and ki-67: potential markers to differentiate low- and high-grade transitional cell carcinomas of the urinary bladder. <i>Journal of Histochemistry and Cytochemistry</i> , <b>2011</b> , 59, 1022-30   | 3.4  | 11 |
| 32 | Genetic profile of bone metastases in renal cell carcinoma. <i>European Urology</i> , <b>2004</b> , 45, 320-4  | 10.2 | 11 |
| 31 | Signet-ring cell carcinoma arising from the urinary bladder. <i>Canadian Journal of Urology</i> , <b>2008</b> , 15, 4266-8.8   |      | 11 |
| 30 | Circulating angiostatin, bFGF, and Tie2/TEK levels and their prognostic impact in bladder cancer. <i>Urology</i> , <b>2012</b> , 80, 737.e13-8   | 1.6  | 10 |
| 29 | Expression of p21(waf1/cip1), p27 (kip1), p63 and androgen receptor in low and high Gleason score prostate cancer. <i>Pathology and Oncology Research</i> , <b>2008</b> , 14, 307-11   | 2.6  | 9  |
| 28 | The influence of expertise of the surgical pathologist to undergrading, upgrading, and understaging of prostate cancer in patients undergoing subsequent radical prostatectomy. <i>International Urology and Nephrology</i> , <b>2014</b> , 46, 371-7                        | 2.3  | 7  |
| 27 | A novel videomicroscopic technique for studying rat ureteral peristalsis in vivo. <i>World Journal of Urology</i> , <b>2009</b> , 27, 265-70   | 4    | 7  |
| 26 | The role of alpha-adrenoreceptors in the treatment of urological diseases. <i>Neurochemistry International</i> , <b>2007</b> , 51, 328-31  | 4.4  | 7  |

|    |  |       |
|----|--|-------|
| 25 | T2a transitional cell carcinoma of the bladder: long-term experience with intravesical immunoprophylaxis with bacillus Calmette-Guerin. <i>Journal of Urology</i> , <b>2003</b> , 169, 931-4; discussion 934-5 <sup>2.5</sup>  | 7     |
| 24 | Cyclophosphamide is contraindicated in patients with a history of transitional cell carcinoma. <i>Clinical Rheumatology</i> , <b>2005</b> , 24, 319-23   | 3.9 7 |
| 23 | Clinical symptoms related to renal cell carcinoma are independent prognostic factors for intraoperative complications and overall survival. <i>International Urology and Nephrology</i> , <b>2009</b> , 41, 835-42 <sup>2.3</sup>  | 5     |
| 22 | The technique of ultrasound guided prostate biopsy. <i>World Journal of Urology</i> , <b>2004</b> , 22, 353-6  | 4 5   |
| 21 | Improving Stone Clearance After Extracorporeal Shock Wave Lithotripsy in Urolithiasis Patients by a Special Terpene Combination (Rowatinex <sup>®</sup> ): Results of a Placebo-Controlled, Randomized Trial. <i>European Urology Supplements</i> , <b>2010</b> , 9, 819-825 | 0.9 4 |
| 20 | Changes in Erythrocyte Transmethylation Ability are Predictive Factors for Tumor Prognosis in Prostate Cancer. <i>Croatica Chemica Acta</i> , <b>2011</b> , 84, 127-131  | 0.8 4 |
| 19 | E-cadherin expression in transitional cell carcinomas. <i>Pathology and Oncology Research</i> , <b>2006</b> , 12, 73-7   | 2.6 4 |
| 18 | Changes of protein expression in prostate cancer having lost its androgen sensitivity. <i>International Urology and Nephrology</i> , <b>2015</b> , 47, 1149-54   | 2.3 3 |
| 17 | Penis covering and simultaneous urethral replacement by scrotal skin for severe penile and urethral necrosis. <i>International Urology and Nephrology</i> , <b>2009</b> , 41, 537-40   | 2.3 3 |
| 16 | A novel approach to the treatment of benign prostatic hyperplasia. <i>BJU International</i> , <b>2006</b> , 97, 1252-5   | 5.6 3 |
| 15 | Improving diagnostic accuracy of prostate carcinoma by systematic random map-biopsy. <i>Pathology and Oncology Research</i> , <b>2000</b> , 6, 111-3   | 2.6 3 |
| 14 | A case of bilateral testicular lymphoma. <i>Pathology and Oncology Research</i> , <b>1999</b> , 5, 152-4   | 2.6 3 |
| 13 | Phytotherapy Adds to the Therapeutic Armamentarium for the Treatment of Mild-To-Moderate Lower Urinary Tract Symptoms in Men. <i>Urologia Internationalis</i> , <b>2020</b> , 104, 333-342   | 1.9 3 |
| 12 | Voiding symptoms and urodynamic findings in patients with modified ileal neobladder. <i>Pathology and Oncology Research</i> , <b>2009</b> , 15, 307-13   | 2.6 2 |
| 11 | Real-time flow resistance compensation for urodynamic examinations of the upper urinary tract through single lumen catheters. <i>International Journal of Urology</i> , <b>2008</b> , 15, 110-3  | 2.3 2 |
| 10 | A novel quantitative method for measuring obstruction in the upper urinary tract: the obstruction coefficient S. <i>International Journal of Urology</i> , <b>2008</b> , 15, 499-504   | 2.3 2 |
| 9  | How Can Cancer-Associated Anemia Be Moderated with Nutritional Factors and How Do Beta Vulgaris L. Ssp. Esculenta Var. Rubra Modify the Transmethylation Reaction in Erythrocytes in Cancerous Patients? <b>2012</b> ,   | 1     |
| 8  | Mucoepidermoid lung tumor appearing as an abscess on the scrotum. <i>Urologia Internationalis</i> , <b>2009</b> , 82, 122-4  | 1.9 1 |

|   |  |      |   |
|---|--|------|---|
| 7 | Re: B.J. De Castro and K. C. Baker: Effect of flexible cystoscopy on serum prostate-specific antigen values (Urology 2009; 73:237-240). <i>Urology</i> , <b>2009</b> , 74, 234         | 1.6  | 1 |
| 6 | Chromosomal Aneuploidy in Azoospermic Men. <i>International Journal of Human Genetics</i> , <b>2006</b> , 6, 171-176   |      | 1 |
| 5 | Bilateral renal cell carcinoma in a horseshoe kidney. <i>Pathology and Oncology Research</i> , <b>2002</b> , 8, 270-1  | 2.6  | 1 |
| 4 | The guestbook of the urology department of Semmelweis University in Budapest: a mirror of international contacts. <i>Journal of Urology</i> , <b>2007</b> , 178, 409-13                | 2.5  | 0 |
| 3 | Diagnostic and treatment options in a papillary pelvic tumor patient with solitary kidney refusing nephrectomy. <i>European Journal of Radiology Extra</i> , <b>2009</b> , 72, e17-e19 |      |   |
| 2 | Words of wisdom. Re: Post-prostatectomy urinary incontinence: a confluence of 3 factors. Loughlin KR, Prasad MM. <i>European Urology</i> , <b>2010</b> , 58, 463-4                     | 10.2 |   |
| 1 | Re: A new postoperative predictor of time to urinary continence after laparoscopic radical prostatectomy: the urine loss ratio. <i>European Urology</i> , <b>2007</b> , 52, 1529-30    | 10.2 |   |