

Liming Xiong

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1484762/publications.pdf>

Version: 2024-02-01

81
papers

14,416
citations

53660

45
h-index

66788

78
g-index

90
all docs

90
docs citations

90
times ranked

14521
citing authors

#	ARTICLE	IF	CITATIONS
1	Plant abiotic stress response and nutrient use efficiency. <i>Science China Life Sciences</i> , 2020, 63, 635-674.	2.3	689
2	LOWER TEMPERATURE 1 Enhances ABA Responses and Plant Drought Tolerance by Modulating the Stability and Localization of C2-Domain ABA-Related Proteins in Arabidopsis. <i>Molecular Plant</i> , 2019, 12, 1243-1258.	3.9	28
3	Subcellular Localization and Functions of Plant lncRNAs in Drought and Salt Stress Tolerance. <i>Methods in Molecular Biology</i> , 2019, 1933, 173-186.	0.4	13
4	Arabidopsis proteome and the mass spectral assay library. <i>Scientific Data</i> , 2019, 6, 278.	2.4	39
5	Spliceosomal protein U1A is involved in alternative splicing and salt stress tolerance in Arabidopsis thaliana. <i>Nucleic Acids Research</i> , 2018, 46, 1777-1792.	6.5	57
6	The caseinolytic protease complex component CLPC1 in Arabidopsis maintains proteome and RNA homeostasis in chloroplasts. <i>BMC Plant Biology</i> , 2018, 18, 192.	1.6	9
7	A Nucleus-Localized Long Non-Coding RNA Enhances Drought and Salt Stress Tolerance. <i>Plant Physiology</i> , 2017, 175, 1321-1336.	2.3	251
8	AtLSG1-2 Regulates Leaf Growth by Affecting Cell Proliferation and the Onset of Endoreduplication and Synergistically Interacts with AtNMD3 during Cell Proliferation Process. <i>Frontiers in Plant Science</i> , 2017, 8, 337.	1.7	2
9	Arabidopsis <i>YAK1</i> regulates abscisic acid response and drought resistance. <i>FEBS Letters</i> , 2016, 590, 2201-2209.	1.3	28
10	The RNA Polymerase II C-Terminal Domain Phosphatase-Like Protein FIERY2/CPL1 Interacts with eIF4AIII and Is Essential for Nonsense-Mediated mRNA Decay in Arabidopsis. <i>Plant Cell</i> , 2016, 28, 770-785.	3.1	21
11	Two domain-disrupted <i>hda6</i> alleles have opposite epigenetic effects on transgenes and some endogenous targets. <i>Scientific Reports</i> , 2015, 5, 17832.	1.6	8
12	Arabidopsis <i>Yak1</i> protein (<i>AtYak1</i>) is a dual specificity protein kinase. <i>FEBS Letters</i> , 2015, 589, 3321-3327.	1.3	18
13	Arabidopsis flower specific defense gene expression patterns affect resistance to pathogens. <i>Frontiers in Plant Science</i> , 2015, 6, 79.	1.7	17
14	The RNA-binding protein HOS5 and serine/arginine-rich proteins RS40 and RS41 participate in miRNA biogenesis in Arabidopsis. <i>Nucleic Acids Research</i> , 2015, 43, 8283-8298.	6.5	67
15	Environmental Stress and Pre-mRNA Splicing. <i>Molecular Plant</i> , 2015, 8, 1302-1303.	3.9	21
16	Proteomic identification of early salicylate- and flg22-responsive redox-sensitive proteins in Arabidopsis. <i>Scientific Reports</i> , 2015, 5, 8625.	1.6	41
17	The Arabidopsis gene <i>DIG6</i> encodes a large 60S subunit nuclear export GTPase 1 that is involved in ribosome biogenesis and affects multiple auxin-regulated development processes. <i>Journal of Experimental Botany</i> , 2015, 66, 6863-6875.	2.4	21
18	<i>AtMYB7</i> , a subgroup 4 <i>R2R3MYB</i> , negatively regulates ABA-induced inhibition of seed germination by blocking the expression of the <i>ZIP5</i> transcription factor <i>ABI5</i> . <i>Plant, Cell and Environment</i> , 2015, 38, 559-571.	2.8	66

#	ARTICLE	IF	CITATIONS
19	The Arabidopsis Vacuolar Sorting Receptor1 Is Required for Osmotic Stress-Induced Abscisic Acid Biosynthesis. <i>Plant Physiology</i> , 2014, 167, 137-152.	2.3	41
20	Dynamic regulation of genome-wide pre-mRNA splicing and stress tolerance by the Sm-like protein LSM5 in Arabidopsis. <i>Genome Biology</i> , 2014, 15, R1.	13.9	1,501
21	Genome-wide analysis of alternative splicing of pre-mRNA under salt stress in Arabidopsis. <i>BMC Genomics</i> , 2014, 15, 431.	1.2	234
22	The Putative E3 Ubiquitin Ligase ECERIFERUM9 Regulates Abscisic Acid Biosynthesis and Response during Seed Germination and Postgermination Growth in Arabidopsis. <i>Plant Physiology</i> , 2014, 165, 1255-1268.	2.3	42
23	GSA1/ARG1 protects root gravitropism in Arabidopsis under ammonium stress. <i>New Phytologist</i> , 2013, 200, 97-111.	3.5	35
24	Arabidopsis cysteine-rich receptor-like kinase 45 functions in the responses to abscisic acid and abiotic stresses. <i>Plant Physiology and Biochemistry</i> , 2013, 67, 189-198.	2.8	57
25	A KH-Domain RNA-Binding Protein Interacts with FIERY2/CTD Phosphatase-Like 1 and Splicing Factors and Is Important for Pre-mRNA Splicing in Arabidopsis. <i>PLoS Genetics</i> , 2013, 9, e1003875.	1.5	88
26	Arabidopsis Plastid AMOS1/EGY1 Integrates Abscisic Acid Signaling to Regulate Global Gene Expression Response to Ammonium Stress. <i>Plant Physiology</i> , 2012, 160, 2040-2051.	2.3	92
27	Genome-Wide Transcriptional Reprogramming Under Drought Stress. , 2012, , 273-289.		3
28	A plant microRNA regulates the adaptation of roots to drought stress. <i>FEBS Letters</i> , 2012, 586, 1742-1747.	1.3	118
29	A Nucleotide Metabolite Controls Stress-Responsive Gene Expression and Plant Development. <i>PLoS ONE</i> , 2011, 6, e26661.	1.1	45
30	Shoot-supplied ammonium targets the root auxin influx carrier AUX1 and inhibits lateral root emergence in Arabidopsis. <i>Plant, Cell and Environment</i> , 2011, 34, 933-946.	2.8	90
31	Genetic interaction of two abscisic acid signaling regulators, HY5 and FIERY1, in mediating lateral root formation. <i>Plant Signaling and Behavior</i> , 2011, 6, 123-125.	1.2	19
32	The Plant Cuticle Is Required for Osmotic Stress Regulation of Abscisic Acid Biosynthesis and Osmotic Stress Tolerance in Arabidopsis. <i>Plant Cell</i> , 2011, 23, 1971-1984.	3.1	147
33	The bifunctional abiotic stress signalling regulator and endogenous RNA silencing suppressor FIERY1 is required for lateral root formation. <i>Plant, Cell and Environment</i> , 2010, 33, 2180-2190.	2.8	41
34	Genetic analysis of pathway regulation for enhancing branched-chain amino acid biosynthesis in plants. <i>Plant Journal</i> , 2010, 63, 573-583.	2.8	57
35	myo-Inositol-1-phosphate Synthase Is Required for Polar Auxin Transport and Organ Development. <i>Journal of Biological Chemistry</i> , 2010, 285, 24238-24247.	1.6	62
36	Alternative splicing of anciently exonized 5S rRNA regulates plant transcription factor TFIIIA. <i>Genome Research</i> , 2009, 19, 913-921.	2.4	34

#	ARTICLE	IF	CITATIONS
37	Localized auxin biosynthesis and postembryonic root development in Arabidopsis. <i>Plant Signaling and Behavior</i> , 2009, 4, 752-754.	1.2	8
38	The short-rooted vitamin B6-deficient mutant pdx1 has impaired local auxin biosynthesis. <i>Planta</i> , 2009, 229, 1303-1310.	1.6	22
39	Enhancement of vitamin B ₆ levels in seeds through metabolic engineering. <i>Plant Biotechnology Journal</i> , 2009, 7, 673-681.	4.1	30
40	Plants, endosymbionts and parasites. <i>Communicative and Integrative Biology</i> , 2008, 1, 62-65.	0.6	20
41	Role of HY5 in abscisic acid response in seeds and seedlings. <i>Plant Signaling and Behavior</i> , 2008, 3, 986-988.	1.2	20
42	Integration of light and abscisic acid signaling during seed germination and early seedling development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 4495-4500.	3.3	251
43	Abscisic Acid In Plant Response And Adaptation To Drought And Salt Stress. , 2007, , 193-221.		13
44	Stress Signal Transduction: components, pathways and network integration. , 2006, , 3-29.		11
45	Identification of Drought Tolerance Determinants by Genetic Analysis of Root Response to Drought Stress and Abscisic Acid. <i>Plant Physiology</i> , 2006, 142, 1065-1074.	2.3	366
46	Pyridoxine is required for post-embryonic root development and tolerance to osmotic and oxidative stresses. <i>Plant Journal</i> , 2005, 44, 396-408.	2.8	163
47	A DEAD Box RNA Helicase Is Essential for mRNA Export and Important for Development and Stress Responses in Arabidopsis. <i>Plant Cell</i> , 2005, 17, 256-267.	3.1	322
48	A single amino acid substitution in the Arabidopsis FIERY1/HOS2 protein confers cold signaling specificity and lithium tolerance. <i>Plant Journal</i> , 2004, 40, 536-545.	2.8	58
49	Regulation of Abscisic Acid Biosynthesis. <i>Plant Physiology</i> , 2003, 133, 29-36.	2.3	708
50	The Arabidopsis salt overly sensitive 4 Mutants Uncover a Critical Role for Vitamin B6 in Plant Salt Tolerance. <i>Plant Cell</i> , 2002, 14, 575-588.	3.1	191
51	Regulation of Osmotic Stress-responsive Gene Expression by the LOS6/ABA1 Locus in Arabidopsis. <i>Journal of Biological Chemistry</i> , 2002, 277, 8588-8596.	1.6	382
52	RNA helicase-like protein as an early regulator of transcription factors for plant chilling and freezing tolerance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 11507-11512.	3.3	275
53	Salt Tolerance. <i>The Arabidopsis Book</i> , 2002, 1, e0048.	0.5	63
54	An Arabidopsis mutation in translation elongation factor 2 causes superinduction of CBF/DREB1 transcription factor genes but blocks the induction of their downstream targets under low temperatures. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 7786-7791.	3.3	144

#	ARTICLE	IF	CITATIONS
55	A Mitochondrial Complex I Defect Impairs Cold-Regulated Nuclear Gene Expression. <i>Plant Cell</i> , 2002, 14, 1235-1251.	3.1	233
56	C-terminal domain phosphatase-like family members (AtCPLs) differentially regulate <i>Arabidopsis thaliana</i> abiotic stress signaling, growth, and development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 10893-10898.	3.3	146
57	Cell Signaling during Cold, Drought, and Salt Stress. <i>Plant Cell</i> , 2002, 14, S165-S183.	3.1	1,874
58	A Calcium Sensor and Its Interacting Protein Kinase Are Global Regulators of Abscisic Acid Signaling in <i>Arabidopsis</i> . <i>Developmental Cell</i> , 2002, 3, 233-244.	3.1	278
59	Repression of stress-responsive genes by FIERY2, a novel transcriptional regulator in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 10899-10904.	3.3	137
60	Molecular and genetic aspects of plant responses to osmotic stress. <i>Plant, Cell and Environment</i> , 2002, 25, 131-139.	2.8	702
61	Title is missing!. <i>Nutrient Cycling in Agroecosystems</i> , 2002, 63, 91-98.	1.1	4
62	LOS2, a genetic locus required for cold-responsive gene transcription encodes a bi-functional enolase. <i>EMBO Journal</i> , 2002, 21, 2692-2702.	3.5	303
63	Modulation of Abscisic Acid Signal Transduction and Biosynthesis by an Sm-like Protein in <i>Arabidopsis</i> . <i>Developmental Cell</i> , 2001, 1, 771-781.	3.1	311
64	The <i>Arabidopsis</i> LOS5/ABA3 Locus Encodes a Molybdenum Cofactor Sulfurase and Modulates Cold Stress- and Osmotic Stress-Responsive Gene Expression. <i>Plant Cell</i> , 2001, 13, 2063.	3.1	11
65	Abiotic stress signal transduction in plants: Molecular and genetic perspectives. <i>Physiologia Plantarum</i> , 2001, 112, 152-166.	2.6	219
66	The <i>Arabidopsis</i> HOS1 gene negatively regulates cold signal transduction and encodes a RING finger protein that displays cold-regulated nucleo-cytoplasmic partitioning. <i>Genes and Development</i> , 2001, 15, 912-924.	2.7	392
67	The <i>Arabidopsis</i> LOS5/ABA3 Locus Encodes a Molybdenum Cofactor Sulfurase and Modulates Cold Stress- and Osmotic Stress-Responsive Gene Expression. <i>Plant Cell</i> , 2001, 13, 2063-2083.	3.1	492
68	FIERY1 encoding an inositol polyphosphate 1-phosphatase is a negative regulator of abscisic acid and stress signaling in <i>Arabidopsis</i> . <i>Genes and Development</i> , 2001, 15, 1971-1984.	2.7	343
69	The <i>Arabidopsis</i> LOS5/ABA3 Locus Encodes a Molybdenum Cofactor Sulfurase and Modulates Cold Stress- and Osmotic Stress-Responsive Gene Expression. <i>Plant Cell</i> , 2001, 13, 2063-2083.	3.1	440
70	Interaction of Osmotic Stress, Temperature, and Abscisic Acid in the Regulation of Gene Expression in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 1999, 119, 205-212.	2.3	172
71	Cold-regulated gene expression and freezing tolerance in an <i>Arabidopsis thaliana</i> mutant. <i>Plant Journal</i> , 1999, 17, 301-308.	2.8	93
72	HOS5-a negative regulator of osmotic stress-induced gene expression in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 1999, 19, 569-578.	2.8	72

#	ARTICLE	IF	CITATIONS
73	High Throughput Screening of Signal Transduction Mutants With Luciferase Imaging. <i>Plant Molecular Biology Reporter</i> , 1999, 17, 159-170.	1.0	20
74	Genetic Analysis of Salt Tolerance in Arabidopsis: Evidence for a Critical Role of Potassium Nutrition. <i>Plant Cell</i> , 1998, 10, 1181-1191.	3.1	607
75	HOS1, a Genetic Locus Involved in Cold-Responsive Gene Expression in Arabidopsis. <i>Plant Cell</i> , 1998, 10, 1151-1161.	3.1	276
76	HOS1, a Genetic Locus Involved in Cold-Responsive Gene Expression in Arabidopsis. <i>Plant Cell</i> , 1998, 10, 1151.	3.1	20
77	Genetic Analysis of Osmotic and Cold Stress Signal Transduction in Arabidopsis: Interactions and Convergence of Abscisic Acid-Dependent and Abscisic Acid-Independent Pathways. <i>Plant Cell</i> , 1997, 9, 1935.	3.1	85
78	Enhanced plant growth by uniform placement of superphosphate with rock phosphate in acidic soils. <i>Communications in Soil Science and Plant Analysis</i> , 1996, 27, 2837-2850.	0.6	11
79	Influence of phosphate on cadmium adsorption by soils. <i>Fertilizer Research</i> , 1995, 40, 31-40.	0.5	7
80	Magnesium influence on plant uptake of phosphorus in a calcareous soil. <i>Journal of Plant Nutrition</i> , 1995, 18, 1251-1261.	0.9	4
81	An evaluation of the agronomic potential of partially acidulated rock phosphates in calcareous soil. <i>Fertilizer Research</i> , 1994, 38, 205-212.	0.5	8