Aleksandra Wlaź

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1488962/publications.pdf

Version: 2024-02-01

		566801	642321
38	637	15	23
papers	citations	h-index	g-index
38	38	38	858
30	30	30	030
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	The Interaction of Selective A1 and A2A Adenosine Receptor Antagonists with Magnesium and Zinc Ions in Mice: Behavioural, Biochemical and Molecular Studies. International Journal of Molecular Sciences, 2021, 22, 1840.	1.8	5
2	Limbal Approach Phacovitrectomy to Treat Cataract with Clinically Significant Asteroid Hyalosisâ€"Presentation of the Technique and Preliminary Results. Journal of Clinical Medicine, 2021, 10, 3338.	1.0	1
3	Intraocular lens power calculations in eyes with pseudoexfoliation syndrome. Scientific Reports, 2021, 11, 19071.	1.6	5
4	Propofol and Sevoflurane Anesthesia in Early Childhood Do Not Influence Seizure Threshold in Adult Rats. International Journal of Environmental Research and Public Health, 2021, 18, 12367.	1.2	1
5	The role of epithelial thickness mapping in corneal refractive surgery. OphthaTherapy Therapies in Ophthalmology, 2021, 8, 196-201.	0.1	O
6	Ligands of the CB2 cannabinoid receptors augment activity of the conventional antidepressant drugs in the behavioural tests in mice. Behavioural Brain Research, 2020, 378, 112297.	1.2	10
7	Influence of the CB1 and CB2 cannabinoid receptor ligands on the activity of atypical antidepressant drugs in the behavioural tests in mice. Pharmacology Biochemistry and Behavior, 2020, 188, 172833.	1.3	11
8	Predictive Value of Bleb Vascularity after Mitomycin C Augmented Trabeculectomy. Journal of Clinical Medicine, 2020, 9, 3501.	1.0	0
9	Influence of the endocannabinoid system on the antidepressant activity of bupropion and moclobemide in the behavioural tests in mice. Pharmacological Reports, 2020, 72, 1562-1572.	1.5	8
10	Zinc signaling and epilepsy., 2019, 193, 156-177.		52
11	Influence of the CB1 cannabinoid receptors on the activity of the monoaminergic system in the behavioural tests in mice. Brain Research Bulletin, 2019, 150, 179-185.	1.4	9
12	Agomelatine and tianeptine antidepressant activity in mice behavioral despair tests is enhanced by DMPX, a selective adenosine A2A receptor antagonist, but not DPCPX, a selective adenosine A1 receptor antagonist. Pharmacological Reports, 2019, 71, 676-681.	1.5	16
13	Phacotrabeculectomy using collagen matrix implant (Ologen \hat{A}°) versus mitomycin C: a prospective randomized controlled trial. Acta Ophthalmologica, 2019, 97, e817-e826.	0.6	9
14	Antidepressant-Like Activity of Typical Antidepressant Drugs in the Forced Swim Test and Tail Suspension Test in Mice Is Augmented by DMPX, an Adenosine A2A Receptor Antagonist. Neurotoxicity Research, 2019, 35, 344-352.	1.3	32
15	CB1 cannabinoid receptor ligands augment the antidepressant-like activity of biometals (magnesium) Tj ETQq1 I	l 0 <mark>,7,8</mark> 431	4 rgBT /Overlo
16	Arvanil, olvanil, AM 1172 and LY 2183240 (various cannabinoid CB1 receptor agonists) increase the threshold for maximal electroshock-induced seizures in mice. Pharmacological Reports, 2018, 70, 106-109.	1.5	6
17	The influence of selective A1 and A2A receptor antagonists on the antidepressant-like activity of moclobemide, venlafaxine and bupropion in mice. Journal of Pharmacy and Pharmacology, 2018, 70, 1200-1208.	1.2	10
18	Combination of phenobarbital with phenytoin and pregabalin produces synergy in the mouse tonic-clonic seizure model: An isobolographic analysis. Epilepsy Research, 2018, 145, 116-122.	0.8	10

#	Article	IF	Citations
19	DPCPX, a selective adenosine A1 receptor antagonist, enhances the antidepressant-like effects of imipramine, escitalopram, and reboxetine in mice behavioral tests. Naunyn-Schmiedeberg's Archives of Pharmacology, 2018, 391, 1361-1371.	1.4	18
20	Withdrawal of caffeine after its chronic administration modifies the antidepressant-like activity of atypical antidepressants in mice. Changes in cortical expression of Comt, Slc6a15 and Adora1 genes. Psychopharmacology, 2018, 235, 2423-2434.	1.5	6
21	Chronic treatment with caffeine and its withdrawal modify the antidepressant-like activity of selective serotonin reuptake inhibitors in the forced swim and tail suspension tests in mice. Effects on Comt, Slc6a15 and Adora1 gene expression. Toxicology and Applied Pharmacology, 2017, 337, 95-103.	1.3	11
22	Caffeine enhances the antidepressant-like activity of common antidepressant drugs in the forced swim test in mice. Naunyn-Schmiedeberg's Archives of Pharmacology, 2016, 389, 211-221.	1.4	46
23	Synergistic antidepressant-like effect of the joint administration of caffeine and NMDA receptor ligands in the forced swim test in mice. Journal of Neural Transmission, 2016, 123, 463-472.	1.4	10
24	Anxiogenic- and antidepressant-like behavior in corneally kindled rats. Pharmacological Reports, 2015, 67, 349-352.	1.5	6
25	Synergistic interaction of levetiracetam with gabapentin in the mouse 6 Hz psychomotor seizure model – a type II isobolographic analysis. Current Issues in Pharmacy and Medical Sciences, 2015, 28, 204-207.	0.1	1
26	Effects of WIN 55,212-2 (a synthetic cannabinoid CB1 and CB2 receptor agonist) on the anticonvulsant activity of various novel antiepileptic drugs against 6Hz-induced psychomotor seizures in mice. Pharmacology Biochemistry and Behavior, 2015, 130, 53-58.	1.3	20
27	The influence of caffeine on the activity of moclobemide, venlafaxine, bupropion and milnacipran in the forced swim test in mice. Life Sciences, 2015, 136, 13-18.	2.0	15
28	Activity and Safety of Inhaled Itraconazole Nanosuspension in a Model Pulmonary Aspergillus fumigatus Infection in Inoculated Young Quails. Mycopathologia, 2015, 180, 35-42.	1.3	22
29	Effects of WIN 55,212-2 (a non-selective cannabinoid CB1 and CB2 receptor agonist) on the protective action of various classical antiepileptic drugs in the mouse 6ÂHz psychomotor seizure model. Journal of Neural Transmission, 2014, 121, 707-715.	1.4	25
30	The effects of ifenprodil on the activity of antidepressant drugs in the forced swim test in mice. Pharmacological Reports, 2014, 66, 1031-1036.	1.5	12
31	Interactions of levetiracetam with carbamazepine, phenytoin, topiramate and vigabatrin in the mouse 6Hz psychomotor seizure model – A type II isobolographic analysis. European Journal of Pharmacology, 2014, 723, 410-418.	1.7	28
32	Effects of WIN 55,212-2 mesylate on the anticonvulsant action of lamotrigine, oxcarbazepine, pregabalin and topiramate against maximal electroshock-induced seizures in mice. European Journal of Pharmacology, 2013, 720, 247-254.	1.7	30
33	Effects of ifenprodil on the antidepressant-like activity of NMDA ligands in the forced swim test in mice. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2013, 46, 29-35.	2.5	25
34	NMDA and AMPA receptors are involved in the antidepressant-like activity of tianeptine in the forced swim test in mice. Pharmacological Reports, 2011, 63, 1526-1532.	1.5	32
35	A complex interaction between glycine/NMDA receptors and serotonergic/noradrenergic antidepressants in the forced swim test in mice. Journal of Neural Transmission, 2011, 118, 1535-1546.	1.4	46
36	Isobolographic characterization of interactions of levetiracetam with the various antiepileptic drugs in the mouse 6Hz psychomotor seizure model. Epilepsy Research, 2009, 86, 163-174.	0.8	45

#	Article	IF	CITATIONS
37	Characterization of acute adverse-effect profiles of selected antiepileptic drugs in the grip-strength test in mice. Pharmacological Reports, 2009, 61, 737-742.	1.5	22
38	D-serine, a selective glycine/N-methyl-D-aspartate receptor agonist, antagonizes the antidepressant-like effects of magnesium and zinc in mice. Pharmacological Reports, 2008, 60, 996-1000.	1.5	24