

Ángel Cabezas Serisa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1491966/publications.pdf>

Version: 2024-02-01

17
papers

327
citations

933447

10
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

693
citing authors

#	ARTICLE	IF	CITATIONS
1	Hypothalamic-pituitary-adrenal axis function and exposure to stress factors and cannabis use in recent-onset psychosis. <i>World Journal of Biological Psychiatry</i> , 2020, 21, 564-571.	2.6	9
2	Association between anti-thyroid antibodies and negative symptoms in early psychosis. <i>Microbial Biotechnology</i> , 2020, 14, 470-475.	1.7	19
3	The Role of Sleep Quality, Trait Anxiety and Hypothalamic-Pituitary-Adrenal Axis Measures in Cognitive Abilities of Healthy Individuals. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7600.	2.6	18
4	Glycated Haemoglobin Is Associated With Poorer Cognitive Performance in Patients With Recent-Onset Psychosis. <i>Frontiers in Psychiatry</i> , 2020, 11, 455.	2.6	4
5	Clinical and cognitive correlates of childhood attention-deficit/hyperactivity disorder in first-episode psychosis: A controlled study. <i>European Neuropsychopharmacology</i> , 2020, 36, 90-99.	0.7	6
6	The impact of sex and cannabis on clinical features in first-admitted patients with psychosis. <i>European Neuropsychopharmacology</i> , 2020, 36, 235-243.	0.7	15
7	Cognitive Biases Questionnaire for Psychosis (CBQp): Spanish Validation and Relationship With Cognitive Insight in Psychotic Patients. <i>Frontiers in Psychiatry</i> , 2020, 11, 596625.	2.6	6
8	Relationship between ANKK1 rs1800497 polymorphism, overweight and executive dysfunction in early psychosis. <i>Schizophrenia Research</i> , 2019, 209, 278-280.	2.0	2
9	Gene-environment interaction between the brain-derived neurotrophic factor <sc>Val66Met</sc> polymorphism, psychosocial stress and dietary intake in early psychosis. <i>Microbial Biotechnology</i> , 2018, 12, 811-820.	1.7	11
10	Sex differences in the relationship between prolactin levels and impaired processing speed in early psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2018, 52, 585-595.	2.3	11
11	F87. SERUM PROLACTIN LEVELS AND COGNITIVE OUTCOME IN FIRST EPISODE PSYCHOSIS: A PROSPECTIVE 1-YEAR FOLLOW-UP STUDY. <i>Schizophrenia Bulletin</i> , 2018, 44, S253-S254.	4.3	0
12	Improvement in cognitive biases after group psychoeducation and metacognitive training in recent-onset psychosis: A randomized crossover clinical trial. <i>Psychiatry Research</i> , 2018, 270, 720-723.	3.3	16
13	Hypothalamic-pituitary-adrenal axis measures and cognitive abilities in early psychosis: Are there sex differences?. <i>Psychoneuroendocrinology</i> , 2016, 72, 54-62.	2.7	17
14	Sex differences in the effect of childhood trauma on the clinical expression of early psychosis. <i>Comprehensive Psychiatry</i> , 2016, 68, 86-96.	3.1	73
15	Free thyroxine levels are associated with cognitive abilities in subjects with early psychosis. <i>Schizophrenia Research</i> , 2015, 166, 37-42.	2.0	21
16	Clinical correlates of obsessive-compulsive symptom dimensions in at-risk mental states and psychotic disorders at early stages. <i>Psychiatry Research</i> , 2015, 228, 363-367.	3.3	10
17	Stress biomarkers as predictors of transition to psychosis in at-risk mental states: Roles for cortisol, prolactin and albumin. <i>Journal of Psychiatric Research</i> , 2015, 60, 163-169.	3.1	89