

Ruey-Hwa Chen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1497040/ruey-hwa-chen-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

86

papers

5,793

citations

40

h-index

75

g-index

87

ext. papers

6,438

ext. citations

10.8

avg, IF

5.37

L-index

#	Paper	IF	Citations
86	Long noncoding RNA BCRP3 stimulates VPS34 and autophagy activities to promote protein homeostasis and cell survival.. <i>Journal of Biomedical Science</i> , 2022 , 29, 30	13.3	0
85	Autophagy and cancer metabolism-The two-way interplay. <i>IUBMB Life</i> , 2021 ,	4.7	1
84	Regulation of autophagy by VPS34 branched ubiquitination controls proteostasis and liver metabolism. <i>Molecular and Cellular Oncology</i> , 2021 , 8, 1915076	1.2	
83	VPS34 K29/K48 branched ubiquitination governed by UBE3C and TRABID regulates autophagy, proteostasis and liver metabolism. <i>Nature Communications</i> , 2021 , 12, 1322	17.4	9
82	Usp11 controls cortical neurogenesis and neuronal migration through Sox11 stabilization. <i>Science Advances</i> , 2021 , 7,	14.3	6
81	Tumor suppressor death-associated protein kinase 1 inhibits necroptosis by p38 MAPK activation. <i>Cell Death and Disease</i> , 2020 , 11, 305	9.8	10
80	Cullin 3 and Its Role in Tumorigenesis. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1217, 187-2106	10	9
79	Integrative analyses of noncoding RNAs reveal the potential mechanisms augmenting tumor malignancy in lung adenocarcinoma. <i>Nucleic Acids Research</i> , 2020 , 48, 1175-1191	20.1	15
78	Branched Ubiquitination: Detection Methods, Biological Functions and Chemical Synthesis. <i>Molecules</i> , 2020 , 25,	4.8	6
77	WD40 protein Wuho controls germline homeostasis via TRIM-NHL tumor suppressor Mei-p26 in. <i>Development (Cambridge)</i> , 2020 , 147,	6.6	1
76	Long non-coding RNA HOXB-AS3 promotes myeloid cell proliferation and its higher expression is an adverse prognostic marker in patients with acute myeloid leukemia and myelodysplastic syndrome. <i>BMC Cancer</i> , 2019 , 19, 617	4.8	27
75	Reactivation of PTEN tumor suppressor for cancer treatment through inhibition of a MYC-WWP1 inhibitory pathway. <i>Science</i> , 2019 , 364,	33.3	115
74	Suppression of autophagy during mitosis via CUL4-RING ubiquitin ligases-mediated WIP1 polyubiquitination and proteasomal degradation. <i>Autophagy</i> , 2019 , 15, 1917-1934	10.2	25
73	LncRNA NORAD is repressed by the YAP pathway and suppresses lung and breast cancer metastasis by sequestering S100P. <i>Oncogene</i> , 2019 , 38, 5612-5626	9.2	61
72	BIK ubiquitination by the E3 ligase Cul5-ASB11 determines cell fate during cellular stress. <i>Journal of Cell Biology</i> , 2019 , 218, 3002-3018	7.3	8
71	miR-103/107 prolong Wnt/ β -catenin signaling and colorectal cancer stemness by targeting Axin2. <i>Scientific Reports</i> , 2019 , 9, 9687	4.9	32
70	Ubiquitin-mediated regulation of autophagy. <i>Journal of Biomedical Science</i> , 2019 , 26, 80	13.3	68

69	PSPC1-interchanged interactions with PTK6 and E-catenin synergize oncogenic subcellular translocations and tumor progression. <i>Nature Communications</i> , 2019 , 10, 5716	17.4	12
68	PSPC1 mediates TGF- β autocrine signalling and Smad2/3 target switching to promote EMT, stemness and metastasis. <i>Nature Cell Biology</i> , 2018 , 20, 479-491	23.4	88
67	Extracellular domain of EpCAM enhances tumor progression through EGFR signaling in colon cancer cells. <i>Cancer Letters</i> , 2018 , 433, 165-175	9.9	29
66	MicroRNA-140-5p inhibits hepatocellular carcinoma by directly targeting the unique isomerase Pin1 to block multiple cancer-driving pathways. <i>Scientific Reports</i> , 2017 , 7, 45915	4.9	36
65	Ubiquitination of tumor suppressor PML regulates prometastatic and immunosuppressive tumor microenvironment. <i>Journal of Clinical Investigation</i> , 2017 , 127, 2982-2997	15.9	25
64	PML degradation fosters an immunosuppressive and pro-metastatic tumor microenvironment. <i>Molecular and Cellular Oncology</i> , 2017 , 4, e1364212	1.2	1
63	Tumour suppressor death-associated protein kinase targets cytoplasmic HIF-1 α for Th17 suppression. <i>Nature Communications</i> , 2016 , 7, 11904	17.4	15
62	Cul3-KLHL20 ubiquitin ligase: physiological functions, stress responses, and disease implications. <i>Cell Division</i> , 2016 , 11, 5	2.8	11
61	KLHL20 links the ubiquitin-proteasome system to autophagy termination. <i>Autophagy</i> , 2016 , 12, 890-1	10.2	6
60	Cul3-KLHL20 Ubiquitin Ligase Governs the Turnover of ULK1 and VPS34 Complexes to Control Autophagy Termination. <i>Molecular Cell</i> , 2016 , 61, 84-97	17.6	138
59	Cullin 3 Ubiquitin Ligases in Cancer Biology: Functions and Therapeutic Implications. <i>Frontiers in Oncology</i> , 2016 , 6, 113	5.3	46
58	Neural activity and CaMKII protect mitochondria from fragmentation in aging <i>Caenorhabditis elegans</i> neurons. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 8768-73	11.5	32
57	KLHL39 suppresses colon cancer metastasis by blocking KLHL20-mediated PML and DAPK ubiquitination. <i>Oncogene</i> , 2015 , 34, 5141-51	9.2	18
56	The effect of resveratrol on protecting corneal epithelial cells from cytotoxicity caused by moxifloxacin and benzalkonium chloride. <i>Investigative Ophthalmology and Visual Science</i> , 2015 , 56, 1575-84		19
55	K33-Linked Polyubiquitination of Coronin 7 by Cul3-KLHL20 Ubiquitin E3 Ligase Regulates Protein Trafficking. <i>Molecular Cell</i> , 2014 , 54, 586-600	17.6	95
54	Fibroblast activation protein (FAP) is essential for the migration of bone marrow mesenchymal stem cells through RhoA activation. <i>PLoS ONE</i> , 2014 , 9, e88772	3.7	39
53	Small GTPase Rab37 targets tissue inhibitor of metalloproteinase 1 for exocytosis and thus suppresses tumour metastasis. <i>Nature Communications</i> , 2014 , 5, 4804	17.4	37
52	USP11 regulates PML stability to control Notch-induced malignancy in brain tumours. <i>Nature Communications</i> , 2014 , 5, 3214	17.4	62

51	SCP phosphatases suppress renal cell carcinoma by stabilizing PML and inhibiting mTOR/HIF signaling. <i>Cancer Research</i> , 2014 , 74, 6935-46	10.1	21
50	The functions and regulations of DAPK in cancer metastasis. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014 , 19, 364-70	5.4	31
49	Regulation of inflammation by DAPK. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014 , 19, 357-63	5.4	30
48	SEN1 deSUMOylates and regulates Pin1 protein activity and cellular function. <i>Cancer Research</i> , 2013 , 73, 3951-62	10.1	60
47	miR-103/107 promote metastasis of colorectal cancer by targeting the metastasis suppressors DAPK and KLF4. <i>Cancer Research</i> , 2012 , 72, 3631-41	10.1	246
46	The role of PML ubiquitination in human malignancies. <i>Journal of Biomedical Science</i> , 2012 , 19, 81	13.3	24
45	C. elegans EIF-3.K promotes programmed cell death through CED-3 caspase. <i>PLoS ONE</i> , 2012 , 7, e36584	3.7	7
44	Activity-dependent retrograde laminin A signaling regulates synapse growth at Drosophila neuromuscular junctions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 17699-704	11.5	27
43	Neurofibromin mediates FAK signaling in confining synapse growth at Drosophila neuromuscular junctions. <i>Journal of Neuroscience</i> , 2012 , 32, 16971-81	6.6	20
42	Structural and functional roles of Daxx SIM phosphorylation in SUMO paralog-selective binding and apoptosis modulation. <i>Molecular Cell</i> , 2011 , 42, 62-74	17.6	112
41	Death-associated protein kinase 1 phosphorylates Pin1 and inhibits its prolyl isomerase activity and cellular function. <i>Molecular Cell</i> , 2011 , 42, 147-59	17.6	123
40	Tumor suppressor death-associated protein kinase is required for full IL-1 β production. <i>Blood</i> , 2011 , 117, 960-70	2.2	54
39	DAPK activates MARK1/2 to regulate microtubule assembly, neuronal differentiation, and tau toxicity. <i>Cell Death and Differentiation</i> , 2011 , 18, 1507-20	12.7	56
38	Latent membrane protein 1 of Epstein-Barr virus regulates death-associated protein kinase 1 in lymphoblastoid cell line. <i>Virology</i> , 2011 , 413, 19-25	3.6	4
37	A Cullin3-KLHL20 Ubiquitin ligase-dependent pathway targets PML to potentiate HIF-1 signaling and prostate cancer progression. <i>Cancer Cell</i> , 2011 , 20, 214-28	24.3	124
36	PDZ-RhoGEF ubiquitination by Cullin3-KLHL20 controls neurotrophin-induced neurite outgrowth. <i>Journal of Cell Biology</i> , 2011 , 193, 985-94	7.3	39
35	The Cullin 3 substrate adaptor KLHL20 mediates DAPK ubiquitination to control interferon responses. <i>EMBO Journal</i> , 2010 , 29, 1748-61	13	87
34	The tumor suppressor death-associated protein kinase targets to TCR-stimulated NF-kappa B activation. <i>Journal of Immunology</i> , 2008 , 180, 3238-49	5.3	44

33	elF3k regulates apoptosis in epithelial cells by releasing caspase 3 from keratin-containing inclusions. <i>Journal of Cell Science</i> , 2008 , 121, 2382-93	5.3	25
32	Breast tumor kinase phosphorylates p190RhoGAP to regulate rho and ras and promote breast carcinoma growth, migration, and invasion. <i>Cancer Research</i> , 2008 , 68, 7779-87	10.1	64
31	Fak56 functions downstream of integrin alphaPS3betanu and suppresses MAPK activation in neuromuscular junction growth. <i>Neural Development</i> , 2008 , 3, 26	3.9	27
30	The tumor suppressor DAPK is reciprocally regulated by tyrosine kinase Src and phosphatase LAR. <i>Molecular Cell</i> , 2007 , 27, 701-16	17.6	58
29	The tumor suppressor DAP-kinase links cell adhesion and cytoskeleton reorganization to cell death regulation. <i>Journal of Biomedical Science</i> , 2006 , 13, 193-9	13.3	15
28	The tumor suppressor DAPK inhibits cell motility by blocking the integrin-mediated polarity pathway. <i>Journal of Cell Biology</i> , 2006 , 172, 619-31	7.3	93
27	Role of breast tumour kinase in the in vitro differentiation of HaCaT cells. <i>British Journal of Dermatology</i> , 2005 , 153, 282-9	4	25
26	Bidirectional signals transduced by DAPK-ERK interaction promote the apoptotic effect of DAPK. <i>EMBO Journal</i> , 2005 , 24, 294-304	13	166
25	Daxx mediates the small ubiquitin-like modifier-dependent transcriptional repression of Smad4. <i>Journal of Biological Chemistry</i> , 2005 , 280, 10164-73	5.4	90
24	Pentoxifylline attenuates tubulointerstitial fibrosis by blocking Smad3/4-activated transcription and profibrogenic effects of connective tissue growth factor. <i>Journal of the American Society of Nephrology: JASN</i> , 2005 , 16, 2702-13	12.7	125
23	Brk activates rac1 and promotes cell migration and invasion by phosphorylating paxillin. <i>Molecular and Cellular Biology</i> , 2004 , 24, 10558-72	4.8	131
22	Antibiotics induce apoptosis of human peritoneal mesothelial cells. <i>Nephrology</i> , 2003 , 8, 142-9	2.2	13
21	Uncoordinated regulation of stress fibers and focal adhesions by DAP kinase. <i>Journal of Cell Science</i> , 2003 , 116, 4777-90	5.3	69
20	Pentoxifylline inhibits platelet-derived growth factor-stimulated cyclin D1 expression in mesangial cells by blocking Akt membrane translocation. <i>Molecular Pharmacology</i> , 2003 , 64, 811-22	4.3	31
19	TGF-beta induces apoptosis through Smad-mediated expression of DAP-kinase. <i>Nature Cell Biology</i> , 2002 , 4, 51-8	23.4	321
18	DAP-kinase induces apoptosis by suppressing integrin activity and disrupting matrix survival signals. <i>Journal of Cell Biology</i> , 2002 , 159, 169-79	7.3	134
17	Transcription of Epstein-Barr virus-encoded nuclear antigen 1 promoter Qp is repressed by transforming growth factor-beta via Smad4 binding element in human BL cells. <i>Virology</i> , 2000 , 277, 184-92	3.6	7
16	Etk, a Btk family tyrosine kinase, mediates cellular transformation by linking Src to STAT3 activation. <i>Molecular and Cellular Biology</i> , 2000 , 20, 2043-54	4.8	118

15	Interleukin-6 inhibits transforming growth factor-beta-induced apoptosis through the phosphatidylinositol 3-kinase/Akt and signal transducers and activators of transcription 3 pathways. <i>Journal of Biological Chemistry</i> , 1999 , 274, 23013-9	5.4	192
14	Suppression of transforming growth factor-beta-induced apoptosis through a phosphatidylinositol 3-kinase/Akt-dependent pathway. <i>Oncogene</i> , 1998 , 17, 1959-68	9.2	173
13	The type II transforming growth factor-beta receptor autophosphorylates not only on serine and threonine but also on tyrosine residues. <i>Journal of Biological Chemistry</i> , 1997 , 272, 14850-9	5.4	92
12	Relationship between adduct formation, rates of excision repair and the cytotoxic and mutagenic effects of structurally-related polycyclic aromatic carcinogens. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1997 , 376, 143-52	3.3	10
11	Identification of partners of TIF34, a component of the yeast eIF3 complex, required for cell proliferation and translation initiation. <i>EMBO Journal</i> , 1997 , 16, 6812-22	13	47
10	A WD-domain protein that is associated with and phosphorylated by the type II TGF-beta receptor. <i>Nature</i> , 1995 , 377, 548-52	50.4	183
9	Lack of correlation between degree of interference with transcription and rate of strand specific repair in the HPRT gene of diploid human fibroblasts. <i>Journal of Biological Chemistry</i> , 1995 , 270, 27222-7	5.4	14
8	Phosphorylation-dependent interaction of the cytoplasmic domains of the type I and type II transforming growth factor-beta receptors. <i>Journal of Biological Chemistry</i> , 1995 , 270, 12235-41	5.4	70
7	Inactivation of the type II receptor reveals two receptor pathways for the diverse TGF-beta activities. <i>Science</i> , 1993 , 260, 1335-8	33.3	370
6	Cloning of a type I TGF-beta receptor and its effect on TGF-beta binding to the type II receptor. <i>Science</i> , 1993 , 260, 1344-8	33.3	384
5	Determination of type I receptor specificity by the type II receptors for TGF-beta or activin. <i>Science</i> , 1993 , 262, 900-2	33.3	209
4	Preferential repair and strand-specific repair of benzo[a]pyrene diol epoxide adducts in the HPRT gene of diploid human fibroblasts. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992 , 89, 5413-7	11.5	116
3	Use of PCR amplification of cDNA to study mechanisms of human cell mutagenesis and malignant transformation. <i>Environmental and Molecular Mutagenesis</i> , 1991 , 18, 239-44	3.2	2
2	Kinds and location of mutations induced by (+/-)-7 beta,8 alpha-dihydroxy-9 alpha,10 alpha-epoxy-7,8,9,10-tetrahydrobenzo[a]pyrene in the coding region of the hypoxanthine (guanine) phosphoribosyltransferase gene in diploid human fibroblasts. <i>Carcinogenesis</i> , 1991 , 12, 71-5	4.6	81
1	Effect of excision repair by diploid human fibroblasts on the kinds and locations of mutations induced by (+/-)-7 beta,8 alpha-dihydroxy-9 alpha,10 alpha-epoxy-7,8,9,10-tetrahydrobenzo[a]pyrene in the coding region of the HPRT gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990 , 87, 8680-4	11.5	127