

Antonella Marucci

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1500515/antonella-marucci-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

29
papers

479
citations

12
h-index

21
g-index

31
ext. papers

547
ext. citations

5.4
avg, IF

2.61
L-index

#	Paper	IF	Citations
29	Role of GALNT2 on Insulin Sensitivity, Lipid Metabolism and Fat Homeostasis.. <i>International Journal of Molecular Sciences</i> , 2022 , 23,	6.3	3
28	Gain of function of Malate Dehydrogenase 2 (MDH2) and familial hyperglycemia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 ,	5.6	1
27	Morphological and molecular characterization of GALNT2-mediated adipogenesis. <i>International Journal of Obesity</i> , 2021 , 45, 1362-1366	5.5	3
26	GALNT2 as a novel modulator of adipogenesis and adipocyte insulin signaling. <i>International Journal of Obesity</i> , 2019 , 43, 2448-2457	5.5	10
25	Association of a homozygous GCK missense mutation with mild diabetes. <i>Molecular Genetics & Genomic Medicine</i> , 2019 , 7, e00728	2.3	4
24	GALNT2 effect on HDL-cholesterol and triglycerides levels in humans: Evidence of pleiotropy?. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 281-282	4.5	9
23	Suggestive evidence of a multi-cytokine resistin pathway in humans and its role on cardiovascular events in high-risk individuals. <i>Scientific Reports</i> , 2017 , 7, 44337	4.9	11
22	GALNT2 mRNA levels are associated with serum triglycerides in humans. <i>Endocrine</i> , 2016 , 53, 331-4	4	5
21	The PPAR α P12A polymorphism is not associated with all-cause mortality in patients with type 2 diabetes mellitus. <i>Endocrine</i> , 2016 , 54, 38-46	4	
20	Sex-specific effect of BMI on insulin sensitivity and TNF- α expression. <i>Acta Diabetologica</i> , 2015 , 52, 413-6	3.9	5
19	Strong evidence of sexual dimorphic effect of adiposity excess on insulin sensitivity. <i>Acta Diabetologica</i> , 2015 , 52, 991-8	3.9	4
18	Loss-of-Function Mutations in APPL1 in Familial Diabetes Mellitus. <i>American Journal of Human Genetics</i> , 2015 , 97, 177-85	11	91
17	Joint effect of insulin signaling genes on all-cause mortality. <i>Atherosclerosis</i> , 2014 , 237, 639-44	3.1	7
16	Low prevalence of HNF1A mutations after molecular screening of multiple MODY genes in 58 Italian families recruited in the pediatric or adult diabetes clinic from a single Italian hospital. <i>Diabetes Care</i> , 2014 , 37, e258-60	14.6	19
15	Serum resistin, cardiovascular disease and all-cause mortality in patients with type 2 diabetes. <i>PLoS ONE</i> , 2014 , 8, e64729	3.7	63
14	Role of obesity on all-cause mortality in whites with type 2 diabetes from Italy. <i>Acta Diabetologica</i> , 2013 , 50, 971-6	3.9	9
13	Role of somatomedin-B-like domains on ENPP1 inhibition of insulin signaling. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013 , 1833, 552-8	4.9	7

12	Role of GALNT2 in the modulation of ENPP1 expression, and insulin signaling and action: GALNT2: a novel modulator of insulin signaling. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013 , 1833, 1388-95	4.9	21
11	GALNT2 expression is reduced in patients with Type 2 diabetes: possible role of hyperglycemia. <i>PLoS ONE</i> , 2013 , 8, e70159	3.7	23
10	ENPP1 affects insulin action and secretion: evidences from in vitro studies. <i>PLoS ONE</i> , 2011 , 6, e19462	3.7	32
9	GRB10 gene and type 2 diabetes in Whites. <i>Journal of Internal Medicine</i> , 2010 , 267, 132-3	10.8	6
8	A functional variant in the gene 3Suntranslated region regulates HSP70 expression and is a potential candidate for insulin resistance-related abnormalities. <i>Journal of Internal Medicine</i> , 2010 , 267, 237-40	10.8	3
7	ENPP1 Q121 variant, increased pulse pressure and reduced insulin signaling, and nitric oxide synthase activity in endothelial cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 1678-83	9.4	22
6	The role of HSP70 on ENPP1 expression and insulin-receptor activation. <i>Journal of Molecular Medicine</i> , 2009 , 87, 139-144	5.5	13
5	Role of PC-1 and ACE genes on insulin resistance and cardiac mass in never-treated hypertensive patients. Suggestive evidence for a digenic additive modulation. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2007 , 17, 181-7	4.5	12
4	Common haplotypes at the adiponectin receptor 1 (ADIPOR1) locus are associated with increased risk of coronary artery disease in type 2 diabetes. <i>Diabetes</i> , 2006 , 55, 2763-70	0.9	37
3	The -318 C>G single-nucleotide polymorphism in GNAI2 gene promoter region impairs transcriptional activity through specific binding of Sp1 transcription factor and is associated with high blood pressure in Caucasians from Italy. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, 6115-9	12.7	17
2	Lack of evidence for the 1484insG variant at the 3SUTR of the protein tyrosine phosphatase 1B (PTP1B) gene as a genetic determinant of diabetic nephropathy development in type 1 diabetic patients. <i>Nephrology Dialysis Transplantation</i> , 2004 , 19, 2419-20	4.3	0
1	Evidence for genetic epistasis in human insulin resistance: the combined effect of PC-1 (K121Q) and PPARgamma2 (P12A) polymorphisms. <i>Journal of Molecular Medicine</i> , 2003 , 81, 718-23	5.5	40