

Mikael Eriksson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1502195/mikael-eriksson-publications-by-year.pdf>

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| | | | |
|--------------------|-------------------------|----------------|-----------------|
| 159 papers | 5,308 citations | 36 h-index | 70 g-index |
| 166 ext. papers | 7,352 ext. citations | 7.7 avg, IF | 4.77 L-index |

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 159 | Rare germline copy number variants (CNVs) and breast cancer risk.. <i>Communications Biology</i> , 2022 , 5, 65 | 6.7 | 0 |
| 158 | Fibrinogen-like protein 2 in gastrointestinal stromal tumour.. <i>Journal of Cellular and Molecular Medicine</i> , 2022 , | 5.6 | 1 |
| 157 | Pathology of Tumors Associated With Pathogenic Germline Variants in 9 Breast Cancer Susceptibility Genes.. <i>JAMA Oncology</i> , 2022 , | 13.4 | 4 |
| 156 | Interval breast cancer is associated with interferon immune response.. <i>European Journal of Cancer</i> , 2022 , 162, 194-205 | 7.5 | 1 |
| 155 | Selinexor in Advanced, Metastatic Dedifferentiated Liposarcoma: A Multinational, Randomized, Double-Blind, Placebo-Controlled Trial.. <i>Journal of Clinical Oncology</i> , 2022 , JCO2101829 | 2.2 | 1 |
| 154 | Genome-wide interaction analysis of menopausal hormone therapy use and breast cancer risk among 62,370 women.. <i>Scientific Reports</i> , 2022 , 12, 6199 | 4.9 | |
| 153 | Risk Assessment in Population-Based Breast Cancer Screening.. <i>Journal of Clinical Oncology</i> , 2022 , JCO2102827 | 10.2 | 1 |
| 152 | A risk model for digital breast tomosynthesis to predict breast cancer and guide clinical care.. <i>Science Translational Medicine</i> , 2022 , 14, eabn3971 | 17.5 | 1 |
| 151 | Breast cancer risks associated with missense variants in breast cancer susceptibility genes.. <i>Genome Medicine</i> , 2022 , 14, 51 | 14.4 | 0 |
| 150 | Prospective evaluation of a breast-cancer risk model integrating classical risk factors and polygenic risk in 15 cohorts from six countries. <i>International Journal of Epidemiology</i> , 2021 , | 7.8 | 6 |
| 149 | Ultra-rare sarcomas: A consensus paper from the Connective Tissue Oncology Society community of experts on the incidence threshold and the list of entities. <i>Cancer</i> , 2021 , 127, 2934-2942 | 6.4 | 11 |
| 148 | Characterization of Benign Breast Diseases and Association With Age, Hormonal Factors, and Family History of Breast Cancer Among Women in Sweden. <i>JAMA Network Open</i> , 2021 , 4, e2114716 | 10.4 | 7 |
| 147 | Low-Dose Tamoxifen for Mammographic Density Reduction: A Randomized Controlled Trial. <i>Journal of Clinical Oncology</i> , 2021 , 39, 1899-1908 | 2.2 | 10 |
| 146 | Mammographic microcalcifications and risk of breast cancer. <i>British Journal of Cancer</i> , 2021 , 125, 759-768 | 7.7 | 3 |
| 145 | Outcome in dedifferentiated chondrosarcoma for patients treated with multimodal therapy: Results from the EUROpean Bone Over 40 Sarcoma Study. <i>European Journal of Cancer</i> , 2021 , 151, 150-158 | 7.5 | 5 |
| 144 | Combined Associations of a Polygenic Risk Score and Classical Risk Factors With Breast Cancer Risk. <i>Journal of the National Cancer Institute</i> , 2021 , 113, 329-337 | 9.7 | 14 |
| 143 | Evaluating the role of alcohol consumption in breast and ovarian cancer susceptibility using population-based cohort studies and two-sample Mendelian randomization analyses. <i>International Journal of Cancer</i> , 2021 , 148, 1338-1350 | 7.5 | 4 |

| | | | |
|-----|---|------|-----|
| 142 | Predictors of mammographic microcalcifications. <i>International Journal of Cancer</i> , 2021 , 148, 1132-1143 | 7.5 | 3 |
| 141 | Mammography features for early markers of aggressive breast cancer subtypes and tumor characteristics: A population-based cohort study. <i>International Journal of Cancer</i> , 2021 , 148, 1351-1359 | 7.5 | 3 |
| 140 | Association between breast cancer risk and disease aggressiveness: Characterizing underlying gene expression patterns. <i>International Journal of Cancer</i> , 2021 , 148, 884-894 | 7.5 | 2 |
| 139 | Opioid Use After Intensive Care: A Nationwide Cohort Study. <i>Critical Care Medicine</i> , 2021 , 49, 462-471 | 1.4 | 4 |
| 138 | A case-only study to identify genetic modifiers of breast cancer risk for BRCA1/BRCA2 mutation carriers. <i>Nature Communications</i> , 2021 , 12, 1078 | 17.4 | 4 |
| 137 | Breast Cancer Risk Genes - Association Analysis in More than 113,000 Women. <i>New England Journal of Medicine</i> , 2021 , 384, 428-439 | 59.2 | 143 |
| 136 | Association of germline genetic variants with breast cancer-specific survival in patient subgroups defined by clinic-pathological variables related to tumor biology and type of systemic treatment. <i>Breast Cancer Research</i> , 2021 , 23, 86 | 8.3 | 1 |
| 135 | Mendelian randomisation study of smoking exposure in relation to breast cancer risk. <i>British Journal of Cancer</i> , 2021 , 125, 1135-1145 | 8.7 | 0 |
| 134 | Genetic insights into biological mechanisms governing human ovarian ageing. <i>Nature</i> , 2021 , 596, 393-397 | 30.4 | 28 |
| 133 | Postinjury Sepsis-Associations With Risk Factors, Impact on Clinical Course, and Mortality: A Retrospective Observational Study 2021 , 3, e0495 | | 1 |
| 132 | Mammographic features are associated with cardiometabolic disease risk and mortality. <i>European Heart Journal</i> , 2021 , 42, 3361-3370 | 9.5 | 1 |
| 131 | Impact of systemic adjuvant therapy and CYP2D6 activity on mammographic density in a cohort of tamoxifen-treated breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2021 , 190, 451-462 | 4.4 | 0 |
| 130 | Reply to T. Suemasu et al. <i>Journal of Clinical Oncology</i> , 2021 , 39, 2966-2968 | 2.2 | |
| 129 | Use of Low-Dose Tamoxifen to Increase Mammographic Screening Sensitivity in Premenopausal Women. <i>Cancers</i> , 2021 , 13, | 6.6 | 3 |
| 128 | Survival Outcomes Associated With 3 Years vs 1 Year of Adjuvant Imatinib for Patients With High-Risk Gastrointestinal Stromal Tumors: An Analysis of a Randomized Clinical Trial After 10-Year Follow-up. <i>JAMA Oncology</i> , 2020 , 6, 1241-1246 | 13.4 | 40 |
| 127 | Inclusion of Endogenous Plasma Dehydroepiandrosterone Sulfate and Mammographic Density in Risk Prediction Models for Breast Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 574-581 | 4.8 | 4 |
| 126 | Transcriptome-wide association study of breast cancer risk by estrogen-receptor status. <i>Genetic Epidemiology</i> , 2020 , 44, 442-468 | 2.6 | 9 |
| 125 | The association of single nucleotide polymorphisms (SNPs) with breast density and breast cancer survival: the Malmö Diet and Cancer Study. <i>Acta Radiologica</i> , 2020 , 61, 1326-1334 | 2 | 3 |

| | | | |
|-----|---|------|----|
| 124 | Three versus one year of adjuvant imatinib for high-risk gastrointestinal stromal tumor (GIST): Survival analysis of a randomized trial after 10 years of follow-up.. <i>Journal of Clinical Oncology</i> , 2020 , 38, 11503-11503 | 2.2 | 2 |
| 123 | Sense of coherence and risk of breast cancer. <i>ELife</i> , 2020 , 9, | 8.9 | 1 |
| 122 | Fine-mapping of 150 breast cancer risk regions identifies 191 likely target genes. <i>Nature Genetics</i> , 2020 , 52, 56-73 | 36.3 | 56 |
| 121 | The impact of alcohol consumption and physical activity on breast cancer: The role of breast cancer risk. <i>International Journal of Cancer</i> , 2020 , 147, 931-939 | 7.5 | 9 |
| 120 | CYP2D6 Genotype Predicts Tamoxifen Discontinuation and Prognosis in Patients With Breast Cancer. <i>Journal of Clinical Oncology</i> , 2020 , 38, 548-557 | 2.2 | 16 |
| 119 | Mammographic density change in a cohort of premenopausal women receiving tamoxifen for breast cancer prevention over 5 years. <i>Breast Cancer Research</i> , 2020 , 22, 101 | 8.3 | 5 |
| 118 | Preoperative accelerated radiotherapy combined with chemotherapy in a defined cohort of patients with high risk soft tissue sarcoma: a Scandinavian Sarcoma Group study. <i>Clinical Sarcoma Research</i> , 2020 , 10, 22 | 2.5 | 1 |
| 117 | Rare cancers of unknown etiology: lessons learned from a European multi-center case-control study. <i>European Journal of Epidemiology</i> , 2020 , 35, 937-948 | 12.1 | 0 |
| 116 | Hormonal determinants of mammographic density and density change. <i>Breast Cancer Research</i> , 2020 , 22, 95 | 8.3 | 9 |
| 115 | Hyperthyroidism is associated with breast cancer risk and mammographic and genetic risk predictors. <i>BMC Medicine</i> , 2020 , 18, 225 | 11.4 | 5 |
| 114 | Identification of Women at High Risk of Breast Cancer Who Need Supplemental Screening. <i>Radiology</i> , 2020 , 297, 327-333 | 20.5 | 11 |
| 113 | Mammographic Density Change and Risk of Breast Cancer. <i>Journal of the National Cancer Institute</i> , 2020 , 112, 391-399 | 9.7 | 14 |
| 112 | The association between breast cancer risk factors and background parenchymal enhancement at dynamic contrast-enhanced breast MRI. <i>Acta Radiologica</i> , 2020 , 61, 1600-1607 | 2 | 3 |
| 111 | Heritability of Mammographic Breast Density, Density Change, Microcalcifications, and Masses. <i>Cancer Research</i> , 2020 , 80, 1590-1600 | 10.1 | 7 |
| 110 | Two truncating variants in FANCC and breast cancer risk. <i>Scientific Reports</i> , 2019 , 9, 12524 | 4.9 | 2 |
| 109 | Localized mammographic density is associated with interval cancer and large breast cancer: a nested case-control study. <i>Breast Cancer Research</i> , 2019 , 21, 8 | 8.3 | 5 |
| 108 | Determinants of Mammographic Density Change. <i>JNCI Cancer Spectrum</i> , 2019 , 3, pkz004 | 4.6 | 17 |
| 107 | Joint association of mammographic density adjusted for age and body mass index and polygenic risk score with breast cancer risk. <i>Breast Cancer Research</i> , 2019 , 21, 68 | 8.3 | 18 |

| | | | |
|-----|---|------|-----|
| 106 | Tobacco smoking and alcohol consumption as risk factors for thymoma - A European case-control study. <i>Cancer Epidemiology</i> , 2019 , 61, 133-138 | 2.8 | 8 |
| 105 | Genome-wide association and transcriptome studies identify target genes and risk loci for breast cancer. <i>Nature Communications</i> , 2019 , 10, 1741 | 17.4 | 47 |
| 104 | Detection of potential microcalcification clusters using multivendor for-presentation digital mammograms for short-term breast cancer risk estimation. <i>Medical Physics</i> , 2019 , 46, 1938-1946 | 4.4 | 5 |
| 103 | Comparison of self-reported and register-based hospital medical data on comorbidities in women. <i>Scientific Reports</i> , 2019 , 9, 3527 | 4.9 | 7 |
| 102 | Genome-wide association study of germline variants and breast cancer-specific mortality. <i>British Journal of Cancer</i> , 2019 , 120, 647-657 | 8.7 | 28 |
| 101 | Interval breast cancer is associated with other types of tumors. <i>Nature Communications</i> , 2019 , 10, 4648 | 17.4 | 14 |
| 100 | Comparison of the sepsis-2 and sepsis-3 definitions in severely injured trauma patients. <i>Journal of Critical Care</i> , 2019 , 54, 125-129 | 4 | 3 |
| 99 | The :p.Arg658* truncating variant is associated with risk of triple-negative breast cancer. <i>Npj Breast Cancer</i> , 2019 , 5, 38 | 7.8 | 12 |
| 98 | ALT-GIST: Randomized phase II trial of imatinib alternating with regorafenib versus imatinib alone for the first-line treatment of metastatic gastrointestinal stromal tumor (GIST).. <i>Journal of Clinical Oncology</i> , 2019 , 37, 11023-11023 | 2.2 | 6 |
| 97 | Impact of gender on post- traumatic intensive care and outcomes. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2019 , 27, 115 | 3.6 | 5 |
| 96 | Prevalence of BRCA1 and BRCA2 pathogenic variants in a large, unselected breast cancer cohort. <i>International Journal of Cancer</i> , 2019 , 144, 1195-1204 | 7.5 | 18 |
| 95 | Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. <i>American Journal of Human Genetics</i> , 2019 , 104, 21-34 | 11 | 363 |
| 94 | Predicting prolonged sick leave among trauma survivors. <i>Scientific Reports</i> , 2019 , 9, 58 | 4.9 | 2 |
| 93 | Associations of obesity and circulating insulin and glucose with breast cancer risk: a Mendelian randomization analysis. <i>International Journal of Epidemiology</i> , 2019 , 48, 795-806 | 7.8 | 52 |
| 92 | The BRCA2 c.68-7T>A variant is not pathogenic: A model for clinical calibration of spliceogenicity. <i>Human Mutation</i> , 2018 , 39, 729-741 | 4.7 | 16 |
| 91 | A comprehensive tool for measuring mammographic density changes over time. <i>Breast Cancer Research and Treatment</i> , 2018 , 169, 371-379 | 4.4 | 26 |
| 90 | Affinity proteomic profiling of plasma for proteins associated to area-based mammographic breast density. <i>Breast Cancer Research</i> , 2018 , 20, 14 | 8.3 | 5 |
| 89 | Long-term prognostic implications of risk factors associated with tumor size: a case study of women regularly attending screening. <i>Breast Cancer Research</i> , 2018 , 20, 31 | 8.3 | 7 |

| | | | |
|----|---|------|-----|
| 88 | Working to improve the management of sarcoma patients across Europe: a policy checklist. <i>BMC Cancer</i> , 2018 , 18, 424 | 4.8 | 9 |
| 87 | EURO-B.O.S.S.: A European study on chemotherapy in bone-sarcoma patients aged over 40: Outcome in primary high-grade osteosarcoma. <i>Tumori</i> , 2018 , 104, 30-36 | 1.7 | 50 |
| 86 | Adjuvant chemotherapy and postoperative radiotherapy in high-risk soft tissue sarcoma patients defined by biological risk factors-A Scandinavian Sarcoma Group study (SSG XX). <i>European Journal of Cancer</i> , 2018 , 99, 78-85 | 7.5 | 20 |
| 85 | Physical activity and mammographic density in an Asian multi-ethnic cohort. <i>Cancer Causes and Control</i> , 2018 , 29, 883-894 | 2.8 | 2 |
| 84 | Inherited factors contribute to an inverse association between preeclampsia and breast cancer. <i>Breast Cancer Research</i> , 2018 , 20, 6 | 8.3 | 9 |
| 83 | Expression of cell cycle regulators and frequency of TP53 mutations in high risk gastrointestinal stromal tumors prior to adjuvant imatinib treatment. <i>PLoS ONE</i> , 2018 , 13, e0193048 | 3.7 | 11 |
| 82 | A transcriptome-wide association study of 229,000 women identifies new candidate susceptibility genes for breast cancer. <i>Nature Genetics</i> , 2018 , 50, 968-978 | 36.3 | 101 |
| 81 | Inclusion of Plasma Prolactin Levels in Current Risk Prediction Models of Premenopausal and Postmenopausal Breast Cancer. <i>JNCI Cancer Spectrum</i> , 2018 , 2, pky055 | 4.6 | 9 |
| 80 | Adjuvant Therapy and Mammographic Density Changes in Women With Breast Cancer. <i>JNCI Cancer Spectrum</i> , 2018 , 2, pky071 | 4.6 | 10 |
| 79 | Differential Burden of Rare and Common Variants on Tumor Characteristics, Survival, and Mode of Detection in Breast Cancer. <i>Cancer Research</i> , 2018 , 78, 6329-6338 | 10.1 | 13 |
| 78 | Thioredoxin a novel biomarker of post-injury sepsis. <i>Free Radical Biology and Medicine</i> , 2017 , 104, 138-143 | 3.8 | 8 |
| 77 | The Scandinavian Sarcoma Group Central Register: 6,000 patients after 25 years of monitoring of referral and treatment of extremity and trunk wall soft-tissue sarcoma. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2017 , 88, 341-347 | 4.3 | 22 |
| 76 | Assessment of Breast Cancer Risk Factors Reveals Subtype Heterogeneity. <i>Cancer Research</i> , 2017 , 77, 3708-3717 | 10.1 | 64 |
| 75 | Cohort Profile: The Karolinska Mammography Project for Risk Prediction of Breast Cancer (KARMA). <i>International Journal of Epidemiology</i> , 2017 , 46, 1740-1741g | 7.8 | 62 |
| 74 | Effect of KIT and PDGFRA Mutations on Survival in Patients With Gastrointestinal Stromal Tumors Treated With Adjuvant Imatinib: An Exploratory Analysis of a Randomized Clinical Trial. <i>JAMA Oncology</i> , 2017 , 3, 602-609 | 13.4 | 92 |
| 73 | SLUG transcription factor: a pro-survival and prognostic factor in gastrointestinal stromal tumour. <i>British Journal of Cancer</i> , 2017 , 116, 1195-1202 | 8.7 | 12 |
| 72 | A clinical model for identifying the short-term risk of breast cancer. <i>Breast Cancer Research</i> , 2017 , 19, 29 | 8.3 | 53 |
| 71 | Association analysis identifies 65 new breast cancer risk loci. <i>Nature</i> , 2017 , 551, 92-94 | 50.4 | 643 |

| | | | |
|----|--|------|-----|
| 70 | Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. <i>Nature Genetics</i> , 2017 , 49, 1767-1778 | 36.3 | 186 |
| 69 | Post-trauma morbidity, measured as sick leave, is substantial and influenced by factors unrelated to injury: a retrospective matched observational cohort study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2017 , 25, 100 | 3.6 | 5 |
| 68 | Body size in early life and risk of breast cancer. <i>Breast Cancer Research</i> , 2017 , 19, 84 | 8.3 | 9 |
| 67 | Reproductive profiles and risk of breast cancer subtypes: a multi-center case-only study. <i>Breast Cancer Research</i> , 2017 , 19, 119 | 8.3 | 26 |
| 66 | Differences in mammographic density between Asian and Caucasian populations: a comparative analysis. <i>Breast Cancer Research and Treatment</i> , 2017 , 161, 353-362 | 4.4 | 33 |
| 65 | Longitudinal fluctuation in mammographic percent density differentiates between interval and screen-detected breast cancer. <i>International Journal of Cancer</i> , 2017 , 140, 34-40 | 7.5 | 4 |
| 64 | Associations between childhood body size and seventeen adverse outcomes: analysis of 65,057 European women. <i>Scientific Reports</i> , 2017 , 7, 16917 | 4.9 | 4 |
| 63 | Body mass index and breast cancer survival: a Mendelian randomization analysis. <i>International Journal of Epidemiology</i> , 2017 , 46, 1814-1822 | 7.8 | 27 |
| 62 | - a novel candidate breast cancer susceptibility locus on 6q14.1. <i>Oncotarget</i> , 2017 , 8, 102769-102782 | 3.3 | 3 |
| 61 | Failure rate of standard rescue with leucovorin for high-dose methotrexate (HDMTX) in osteosarcoma.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 11028-11028 | 2.2 | |
| 60 | No clinical utility of KRAS variant rs61764370 for ovarian or breast cancer. <i>Gynecologic Oncology</i> , 2016 , 141, 386-401 | 4.9 | 15 |
| 59 | Comparison of MAPIE versus MAP in patients with a poor response to preoperative chemotherapy for newly diagnosed high-grade osteosarcoma (EURAMOS-1): an open-label, international, randomised controlled trial. <i>Lancet Oncology, The</i> , 2016 , 17, 1396-1408 | 21.7 | 253 |
| 58 | Common diseases as determinants of menopausal age. <i>Human Reproduction</i> , 2016 , 31, 2856-2864 | 5.7 | 20 |
| 57 | No evidence that protein truncating variants in BRIP1 are associated with breast cancer risk: implications for gene panel testing. <i>Journal of Medical Genetics</i> , 2016 , 53, 298-309 | 5.8 | 83 |
| 56 | Genetic variation in the immunosuppression pathway genes and breast cancer susceptibility: a pooled analysis of 42,510 cases and 40,577 controls from the Breast Cancer Association Consortium. <i>Human Genetics</i> , 2016 , 135, 137-54 | 6.3 | 6 |
| 55 | Associations of Breast Cancer Risk Prediction Tools With Tumor Characteristics and Metastasis. <i>Journal of Clinical Oncology</i> , 2016 , 34, 251-8 | 2.2 | 27 |
| 54 | Adjuvant Imatinib for High-Risk GI Stromal Tumor: Analysis of a Randomized Trial. <i>Journal of Clinical Oncology</i> , 2016 , 34, 244-50 | 2.2 | 131 |
| 53 | HSP90 inhibition blocks ERBB3 and RET phosphorylation in myxoid/round cell liposarcoma and causes massive cell death in vitro and in vivo. <i>Oncotarget</i> , 2016 , 7, 433-45 | 3.3 | 11 |

| | | | |
|----|--|------|-----|
| 52 | Association of breast cancer risk with genetic variants showing differential allelic expression: Identification of a novel breast cancer susceptibility locus at 4q21. <i>Oncotarget</i> , 2016 , 7, 80140-80163 | 3.3 | 21 |
| 51 | PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS. <i>Journal of Medical Genetics</i> , 2016 , 53, 800-811 | 5.8 | 121 |
| 50 | Needle biopsy through the abdominal wall for the diagnosis of gastrointestinal stromal tumour - Does it increase the risk for tumour cell seeding and recurrence?. <i>European Journal of Cancer</i> , 2016 , 59, 128-133 | 7.5 | 33 |
| 49 | Evidence that the 5p12 Variant rs10941679 Confers Susceptibility to Estrogen-Receptor-Positive Breast Cancer through FGF10 and MRPS30 Regulation. <i>American Journal of Human Genetics</i> , 2016 , 99, 903-911 | 11 | 43 |
| 48 | 15-year follow-up of the Second Nordic Mantle Cell Lymphoma trial (MCL2): prolonged remissions without survival plateau. <i>British Journal of Haematology</i> , 2016 , 175, 410-418 | 4.5 | 118 |
| 47 | Worse quality of life in young and recently diagnosed breast cancer survivors compared with female survivors of other cancers: A cross-sectional study. <i>International Journal of Cancer</i> , 2016 , 139, 2415-25 | 7.5 | 18 |
| 46 | Inherited variants in the inner centromere protein (INCENP) gene of the chromosomal passenger complex contribute to the susceptibility of ER-negative breast cancer. <i>Carcinogenesis</i> , 2015 , 36, 256-71 | 4.6 | 12 |
| 45 | Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. <i>Nature Genetics</i> , 2015 , 47, 373-80 | 36.3 | 406 |
| 44 | Predictors of Discontinuation of Adjuvant Hormone Therapy in Patients With Breast Cancer. <i>Journal of Clinical Oncology</i> , 2015 , 33, 2262-9 | 2.2 | 56 |
| 43 | Identification of novel genetic markers of breast cancer survival. <i>Journal of the National Cancer Institute</i> , 2015 , 107, | 9.7 | 38 |
| 42 | Socio-economic status and co-morbidity as risk factors for trauma. <i>European Journal of Epidemiology</i> , 2015 , 30, 151-7 | 12.1 | 43 |
| 41 | Background risk of breast cancer and the association between physical activity and mammographic density. <i>Breast Cancer Research</i> , 2015 , 17, 50 | 8.3 | 16 |
| 40 | Height and Breast Cancer Risk: Evidence From Prospective Studies and Mendelian Randomization. <i>Journal of the National Cancer Institute</i> , 2015 , 107, | 9.7 | 74 |
| 39 | Radiotherapy for GIST progressing during or after tyrosine kinase inhibitor therapy: A prospective study. <i>Radiotherapy and Oncology</i> , 2015 , 116, 233-8 | 5.3 | 25 |
| 38 | Identification and characterization of novel associations in the CASP8/ALS2CR12 region on chromosome 2 with breast cancer risk. <i>Human Molecular Genetics</i> , 2015 , 24, 285-98 | 5.6 | 35 |
| 37 | Common germline polymorphisms associated with breast cancer-specific survival. <i>Breast Cancer Research</i> , 2015 , 17, 58 | 8.3 | 24 |
| 36 | Identification of two novel mammographic density loci at 6Q25.1. <i>Breast Cancer Research</i> , 2015 , 17, 75 | 8.3 | 18 |
| 35 | Referral patterns, treatment and outcome of high-grade malignant bone sarcoma in Scandinavia--SSG Central Register 25 years' experience. <i>Journal of Surgical Oncology</i> , 2015 , 112, 853-60 | 2.8 | 10 |

| | | | |
|----|--|------|-----|
| 34 | Acute kidney injury following severe trauma: Risk factors and long-term outcome. <i>Journal of Trauma and Acute Care Surgery</i> , 2015 , 79, 407-12 | 3.3 | 61 |
| 33 | The genetic basis of quality of life in healthy Swedish women: a candidate gene approach. <i>PLoS ONE</i> , 2015 , 10, e0118292 | 3.7 | 6 |
| 32 | Prediction of breast cancer risk based on profiling with common genetic variants. <i>Journal of the National Cancer Institute</i> , 2015 , 107, | 9.7 | 324 |
| 31 | Fine-mapping identifies two additional breast cancer susceptibility loci at 9q31.2. <i>Human Molecular Genetics</i> , 2015 , 24, 2966-84 | 5.6 | 36 |
| 30 | Risk factors and tumor characteristics of interval cancers by mammographic density. <i>Journal of Clinical Oncology</i> , 2015 , 33, 1030-7 | 2.2 | 82 |
| 29 | Fine-scale mapping of the 5q11.2 breast cancer locus reveals at least three independent risk variants regulating MAP3K1. <i>American Journal of Human Genetics</i> , 2015 , 96, 5-20 | 11 | 59 |
| 28 | Successful change of treatment strategy in elderly patients with primary central nervous system lymphoma by de-escalating induction and introducing temozolomide maintenance: results from a phase II study by the Nordic Lymphoma Group. <i>Haematologica</i> , 2015 , 100, 534-40 | 6.6 | 59 |
| 27 | Three vs. 1 year of adjuvant imatinib (IM) for operable high-risk GIST: The second planned analysis of the randomized SSGXVIII/AIO trial.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 10505-10505 | 2.2 | 4 |
| 26 | Event-free survival and overall survival in 2,253 patients with osteosarcoma registered to EURAMOS-1.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 10512-10512 | 2.2 | 2 |
| 25 | In Search for the Genetic Basis of Quality of Life in Healthy Swedish Women--A GWAS Study Using the iCOGS Custom Genotyping Array. <i>PLoS ONE</i> , 2015 , 10, e0140563 | 3.7 | 1 |
| 24 | Evidence that breast cancer risk at the 2q35 locus is mediated through IGFBP5 regulation. <i>Nature Communications</i> , 2014 , 4, 4999 | 17.4 | 87 |
| 23 | Deregulation of COMMD1 is associated with poor prognosis in diffuse large B-cell lymphoma. <i>PLoS ONE</i> , 2014 , 9, e91031 | 3.7 | 17 |
| 22 | MicroRNA related polymorphisms and breast cancer risk. <i>PLoS ONE</i> , 2014 , 9, e109973 | 3.7 | 37 |
| 21 | Area and volumetric density estimation in processed full-field digital mammograms for risk assessment of breast cancer. <i>PLoS ONE</i> , 2014 , 9, e110690 | 3.7 | 21 |
| 20 | Automated measurement of volumetric mammographic density: a tool for widespread breast cancer risk assessment. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2014 , 23, 1764-72 | 4 | 52 |
| 19 | Volumetric mammographic density: heritability and association with breast cancer susceptibility loci. <i>Journal of the National Cancer Institute</i> , 2014 , 106, | 9.7 | 15 |
| 18 | Common non-synonymous SNPs associated with breast cancer susceptibility: findings from the Breast Cancer Association Consortium. <i>Human Molecular Genetics</i> , 2014 , 23, 6096-111 | 5.6 | 48 |
| 17 | Plasma Affinity Proteomic Immunoprofiling As a Novel Prognostic Tool in High Risk Diffuse Large B-Cell Lymphoma Patients: A Nordic Lymphoma Group Study. <i>Blood</i> , 2014 , 124, 1618-1618 | 2.2 | |

| | | | |
|----|---|-----|----|
| 16 | MAP plus maintenance pegylated interferon α 2b (MAPIfn) versus MAP alone in patients with resectable high-grade osteosarcoma and good histologic response to preoperative MAP: First results of the EURAMOS-1 good response randomization.. <i>Journal of Clinical Oncology</i> , 2013 , 31, LBA10504-LBA10504 | 2.2 | 14 |
| 15 | MAP plus maintenance pegylated interferon α 2b (MAP-IFN) versus MAP alone in patients (pts) with resectable high-grade osteosarcoma and good histologic response to preoperative MAP: First results of the EURAMOS-1 good response randomization.. <i>Journal of Clinical Oncology</i> , 2013 , 31, LBA10504-LBA10504 | 2.2 | 2 |
| 14 | Error in a study of the outcome of mantle cell lymphoma: Nordic MCL2 Trial Update: 6-year follow-up after intensive immunochemotherapy for untreated mantle cell lymphoma followed by BEAM or BEACOPP autologous stem-cell support: still very long survival but late relapses do occur. <i>British Journal of Haematology</i> , 2012 , 158, 815-816 | 4.5 | 1 |
| 13 | Epidemiological Studies on Cancer and Exposure to Dioxins and Related Compounds 2012 , 303-358 | | 1 |
| 12 | EURAMOS-1 study: Recruitment, characteristics, and initial treatment of more than 2,000 patients (pts) with high-grade osteosarcoma.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 10081-10081 | 2.2 | 2 |
| 11 | Use of exon-based transcriptome profiling to identify novel signaling pathways and survival-associated genes in diffuse large B-cell lymphoma.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 8074-8074 | 2.2 | 2 |
| 10 | Age-Adjusted Combined Immunochemotherapy without Radiotherapy in Newly Diagnosed PCNSL □ A Phase II Trial of the Nordic Lymphoma Group. <i>Blood</i> , 2011 , 118, 1602-1602 | 2.2 | |
| 9 | Occupation and occupational exposure to endocrine disrupting chemicals in male breast cancer: a case-control study in Europe. <i>Occupational and Environmental Medicine</i> , 2010 , 67, 837-44 | 2.1 | 46 |
| 8 | R-CHOEP-14 □ Followed by Systemic CNS Prophylaxis for Diffuse Large B-Cell Lymphoma/Follicular Lymphoma Grade 3 with Age Adjusted IPI Score 2□: Final Results of a Nordic Lymphoma Group Phase 2 Study Including 156 Patients Aged 18-65 Years.. <i>Blood</i> , 2010 , 116, 2805-2805 | 2.2 | 2 |
| 7 | R-CHOEP-14 X 6 Followed by Systemic CNS Prophylaxis for Diffuse Large B-Cell Lymphoma (DLBCL)/Follicular Lymphoma (FL) Grade 3 with Age Adjusted IPI Score 2□: Preliminary Results of a Nordic Lymphoma Group (NLG) Phase 2 Study Including 160 Patients Aged 18- 64 Years. <i>Blood</i> , 2008 , 112, 3604-3604 | 2.2 | |
| 6 | Mantle Cell Lymphoma Can Be Cured by Intensive Immunochemotherapy with In-Vivo Purged Stem-Cell Support; Final Report of the Nordic Lymphoma Group MCL2 Study.. <i>Blood</i> , 2007 , 110, LB1-LB1 | 2.2 | |
| 5 | European multi-centre case-control study on risk factors for rare cancers of unknown aetiology. <i>European Journal of Cancer</i> , 2005 , 41, 601-12 | 7.5 | 27 |
| 4 | Nordic Mantle Cell Lymphoma (MCL) Project: Preemptive Rituximab Treatment of Molecular Relapse Following Autotransplant Can Reinduce Molecular Remission and Prolonged Disease-Free Survival.. <i>Blood</i> , 2005 , 106, 2429-2429 | 2.2 | 1 |
| 3 | Angioid streaks are part of a familial syndrome of dyserythropoietic anaemia (CDA III). <i>British Journal of Haematology</i> , 1997 , 98, 845-9 | 4.5 | 27 |
| 2 | Prospective Evaluation of a Breast Cancer Risk Model Integrating Classical Risk Factors and Polygenic Risk in 15 Cohorts from Six Countries | | 2 |
| 1 | Fine-mapping of 150 breast cancer risk regions identifies 178 high confidence target genes | | 2 |