

# Martin Stevens

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/1525143/martin-stevens-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

139  
papers

2,496  
citations

26  
h-index

43  
g-index

147  
ext. papers

2,953  
ext. citations

3.3  
avg, IF

4.97  
L-index

#	Paper	IF	Citations
139	Gentle thawing of the frozen shoulder: a prospective study of supervised neglect versus intensive physical therapy in seventy-seven patients with frozen shoulder syndrome followed up for two years. <i>Journal of Shoulder and Elbow Surgery</i> , <b>2004</b> , 13, 499-502	4.3	206
138	Conservative or surgical treatment for subacromial impingement syndrome? A systematic review. <i>Journal of Shoulder and Elbow Surgery</i> , <b>2009</b> , 18, 652-60	4.3	126
137	Incidence, prevalence, and consultation rates of shoulder complaints in general practice. <i>Scandinavian Journal of Rheumatology</i> , <b>2012</b> , 41, 150-5	1.9	119
136	Reliability and validity of the short questionnaire to assess health-enhancing physical activity (SQUASH) in patients after total hip arthroplasty. <i>BMC Musculoskeletal Disorders</i> , <b>2008</b> , 9, 141	2.8	99
135	Grip strength is strongly associated with height, weight and gender in childhood: a cross sectional study of 2241 children and adolescents providing reference values. <i>Journal of Physiotherapy</i> , <b>2013</b> , 59, 255-61	2.9	78
134	The validity of the sit-and-reach test and the modified sit-and-reach test in middle-aged to older men and women. <i>Research Quarterly for Exercise and Sport</i> , <b>2003</b> , 74, 331-6	1.9	67
133	Preoperative or postoperative self-efficacy: which is a better predictor of outcome after total hip or knee arthroplasty?. <i>Patient Education and Counseling</i> , <b>2007</b> , 66, 92-9	3.1	63
132	Effects of remote feedback in home-based physical activity interventions for older adults: a systematic review. <i>Patient Education and Counseling</i> , <b>2013</b> , 91, 14-24	3.1	61
131	Sensitivity of sensor-based sit-to-stand peak power to the effects of training leg strength, leg power and balance in older adults. <i>Gait and Posture</i> , <b>2014</b> , 39, 303-7	2.6	56
130	Minimally invasive and computer-navigated total hip arthroplasty: a qualitative and systematic review of the literature. <i>BMC Musculoskeletal Disorders</i> , <b>2010</b> , 11, 92	2.8	49
129	Adherence to and effectiveness of an individually tailored home-based exercise program for frail older adults, driven by mobility monitoring: design of a prospective cohort study. <i>BMC Public Health</i> , <b>2014</b> , 14, 570	4.1	48
128	Recovery of gait after short-stay total hip arthroplasty. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2007</b> , 88, 361-7	2.8	46
127	The development of a comprehensive multidisciplinary care pathway for patients with a hip fracture: design and results of a clinical trial. <i>BMC Musculoskeletal Disorders</i> , <b>2014</b> , 15, 188	2.8	43
126	Reliability of the Groningen Fitness Test for the Elderly. <i>Journal of Aging and Physical Activity</i> , <b>2001</b> , 9, 194-212	1.6	40
125	The association between comorbidity and length of hospital stay and costs in total hip arthroplasty patients: a systematic review. <i>Journal of Arthroplasty</i> , <b>2014</b> , 29, 1009-14	4.4	39
124	Habitual physical activity after total knee replacement. <i>Physical Therapy</i> , <b>2012</b> , 92, 1109-16	3.3	37
123	Physical functioning before and after total hip arthroplasty: perception and performance. <i>Physical Therapy</i> , <b>2008</b> , 88, 712-9	3.3	35

122	Patient satisfaction at and after discharge. Effect of a time lag. <i>Patient Education and Counseling</i> , <b>2006</b> , 60, 241-5	3.1	35
121	Effect of exercise and weight loss in people who have hip osteoarthritis and are overweight or obese: a prospective cohort study. <i>Physical Therapy</i> , <b>2013</b> , 93, 137-46	3.3	32
120	Similar Superior Patient-Reported Outcome Measures for Anterior and Posterolateral Approaches After Total Hip Arthroplasty: Postoperative Patient-Reported Outcome Measure Improvement After 3 months in 12,774 Primary Total Hip Arthroplasties Using the Anterior, Anterolateral, Straight Lateral, or Posterolateral Approach. <i>Journal of Arthroplasty</i> , <b>2018</b> , 33, 1786-1793	4.4	31
119	Groningen Active Living Model (GALM): stimulating physical activity in sedentary older adults. <i>Preventive Medicine</i> , <b>1999</b> , 29, 267-76	4.3	31
118	Cross-cultural adaptation of the Dutch Short Musculoskeletal Function Assessment questionnaire (SMFA-NL): internal consistency, validity, repeatability and responsiveness. <i>Injury</i> , <b>2012</b> , 43, 726-33	2.5	29
117	Six-month effects of the Groningen active living model (GALM) on physical activity, health and fitness outcomes in sedentary and underactive older adults aged 55-65. <i>Patient Education and Counseling</i> , <b>2006</b> , 62, 132-41	3.1	29
116	Reliability and validity of the Dutch version of the Foot and Ankle Outcome Score (FAOS). <i>BMC Musculoskeletal Disorders</i> , <b>2013</b> , 14, 183	2.8	26
115	Comparison of gait in patients following a computer-navigated minimally invasive anterior approach and a conventional posterolateral approach for total hip arthroplasty: a randomized controlled trial. <i>Journal of Orthopaedic Research</i> , <b>2013</b> , 31, 288-94	3.8	26
114	Reliability and validity of the Dutch version of the International Physical Activity Questionnaire in patients after total hip arthroplasty or total knee arthroplasty. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2013</b> , 43, 650-9	4.2	26
113	Reliability and concurrent validity of the Dutch hip and knee replacement expectations surveys. <i>BMC Musculoskeletal Disorders</i> , <b>2010</b> , 11, 242	2.8	25
112	Habitual physical activity behavior of patients after primary total hip arthroplasty. <i>Physical Therapy</i> , <b>2008</b> , 88, 1039-48	3.3	25
111	Sensor-based monitoring of sit-to-stand performance is indicative of objective and self-reported aspects of functional status in older adults. <i>Gait and Posture</i> , <b>2015</b> , 41, 935-40	2.6	24
110	The therapeutic validity and effectiveness of physiotherapeutic exercise following total hip arthroplasty for osteoarthritis: A systematic review. <i>PLoS ONE</i> , <b>2018</b> , 13, e0194517	3.7	24
109	Subjects with hip osteoarthritis show distinctive patterns of trunk movements during gait-a body-fixed-sensor based analysis. <i>Journal of NeuroEngineering and Rehabilitation</i> , <b>2012</b> , 9, 3	5.3	24
108	Patients with shoulder complaints in general practice: consumption of medical care. <i>Rheumatology</i> , <b>2011</b> , 50, 389-95	3.9	23
107	Groningen Active Living Model (GALM): stimulating physical activity in sedentary older adults; validation of the behavioral change model. <i>Preventive Medicine</i> , <b>2003</b> , 37, 561-70	4.3	23
106	Physical activity behavior of patients 1 year after primary total hip arthroplasty: a prospective multicenter cohort study. <i>Physical Therapy</i> , <b>2011</b> , 91, 373-80	3.3	22
105	A Home-Based Exercise Program Driven by Tablet Application and Mobility Monitoring for Frail Older Adults: Feasibility and Practical Implications. <i>Preventing Chronic Disease</i> , <b>2017</b> , 14, E12	3.7	21

104	Local Treatment with Adjuvant Therapy for Central Atypical Cartilaginous Tumors in the Long Bones: Analysis of Outcome and Complications in One Hundred and Eight Patients with a Minimum Follow-up of Two Years. <i>Journal of Bone and Joint Surgery - Series A</i> , <b>2016</b> , 98, 303-13	5.6	21
103	Test-retest reliability of sensor-based sit-to-stand measures in young and older adults. <i>Gait and Posture</i> , <b>2014</b> , 40, 220-4	2.6	21
102	Poorer survival after a primary implant during revision total knee arthroplasty. <i>International Orthopaedics</i> , <b>2013</b> , 37, 415-9	3.8	21
101	The Influence of Hand Preference on Grip Strength in Children and Adolescents; A Cross-Sectional Study of 2284 Children and Adolescents. <i>PLoS ONE</i> , <b>2015</b> , 10, e0143476	3.7	21
100	Neuropathic-like symptoms and the association with joint-specific function and quality of life in patients with hip and knee osteoarthritis. <i>PLoS ONE</i> , <b>2018</b> , 13, e0199165	3.7	20
99	Computer-assisted surgery in orthopedic oncology. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2014</b> , 85, 663-9	4.3	20
98	The influence of overweight/obesity on patient-perceived physical functioning and health-related quality of life after primary total hip arthroplasty. <i>Obesity Surgery</i> , <b>2012</b> , 22, 523-9	3.7	19
97	Changes in body weight after total hip arthroplasty: short-term and long-term effects. <i>Physical Therapy</i> , <b>2012</b> , 92, 680-7	3.3	19
96	Physical activity behavior after total hip arthroplasty (THA): a prediction based on patient characteristics. <i>Patient Education and Counseling</i> , <b>2007</b> , 69, 196-9	3.1	19
95	Biceps Autograft Augmentation for Rotator Cuff Repair: A Systematic Review. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , <b>2018</b> , 34, 1297-1305	5.4	18
94	Effectiveness of computer-navigated minimally invasive total hip surgery compared to conventional total hip arthroplasty: design of a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , <b>2007</b> , 8, 4	2.8	18
93	The Modified painDETECT Questionnaire for Patients with Hip or Knee Osteoarthritis: Translation into Dutch, Cross-Cultural Adaptation and Reliability Assessment. <i>PLoS ONE</i> , <b>2015</b> , 10, e0146117	3.7	17
92	Twelve-month effects of the Groningen active living model (GALM) on physical activity, health and fitness outcomes in sedentary and underactive older adults aged 55-65. <i>Patient Education and Counseling</i> , <b>2007</b> , 66, 167-76	3.1	17
91	Groningen Orthopaedic Social Support Scale: validity and reliability. <i>Journal of Advanced Nursing</i> , <b>2004</b> , 47, 57-63	3.1	17
90	Groningen active living model (GALM): stimulating physical activity in sedentary older adults; first results. <i>Preventive Medicine</i> , <b>2000</b> , 31, 547-53	4.3	17
89	The effect of bearing type on the outcome of total hip arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2018</b> , 89, 163-169	4.3	17
88	Validity and Reliability of the Dutch Version of the International Hip Outcome Tool (iHOT-12NL) in Patients With Disorders of the Hip. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2015</b> , 45, 1026-34, A1-2	4.2	16
87	Physical activity participation among patients after total hip and knee arthroplasty. <i>Clinics in Geriatric Medicine</i> , <b>2012</b> , 28, 509-20	3.8	16

86	After total knee arthroplasty, many people are not active enough to maintain their health and fitness: an observational study. <i>Journal of Physiotherapy</i> , <b>2012</b> , 58, 113-6	2.9	15
85	Patient Characteristics Influence Revision Rate of Total Hip Arthroplasty: American Society of Anesthesiologists Score and Body Mass Index Were the Strongest Predictors for Short-Term Revision After Primary Total Hip Arthroplasty. <i>Journal of Arthroplasty</i> , <b>2020</b> , 35, 188-192.e2	4.4	15
84	Does imageless computer-assisted TKA lead to improved rotational alignment or fewer outliers? A systematic review. <i>Clinical Orthopaedics and Related Research</i> , <b>2014</b> , 472, 3124-33	2.2	14
83	Compensatory trunk movements in patients with hip osteoarthritis: accuracy and reproducibility of a body-fixed sensor-based assessment. <i>American Journal of Physical Medicine and Rehabilitation</i> , <b>2011</b> , 90, 681-7	2.6	14
82	The Groningen Orthopedic Exit Strategy (GOES): a home-based support program for total hip and knee arthroplasty patients after shortened hospital stay. <i>Patient Education and Counseling</i> , <b>2004</b> , 54, 95-9	3.1	14
81	Feasibility and Patient Experience of a Home-Based Rehabilitation Program Driven by a Tablet App and Mobility Monitoring for Patients After a Total Hip Arthroplasty. <i>JMIR MHealth and UHealth</i> , <b>2019</b> , 7, e10342	5.5	14
80	LEAK study: design of a nationwide randomised controlled trial to find the best way to treat wound leakage after primary hip and knee arthroplasty. <i>BMJ Open</i> , <b>2017</b> , 7, e018673	3	13
79	Assessment of prosthesis alignment after revision total knee arthroplasty using EOS 2D and 3D imaging: a reliability study. <i>PLoS ONE</i> , <b>2014</b> , 9, e104613	3.7	13
78	A Dutch translation of the Self-Efficacy for Rehabilitation Outcome Scale (SER): a first impression on reliability and validity. <i>Patient Education and Counseling</i> , <b>2005</b> , 58, 121-6	3.1	13
77	The Groningen Enjoyment Questionnaire: a measure of enjoyment in leisure-time physical activity. <i>Perceptual and Motor Skills</i> , <b>2000</b> , 90, 601-4	2.2	13
76	Accuracy and concurrent validity of a sensor-based analysis of sit-to-stand movements in older adults. <i>Gait and Posture</i> , <b>2016</b> , 45, 198-203	2.6	12
75	Tibial component with and without stem extension in a trabecular metal cone construct. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2017</b> , 25, 3644-3652	5.5	12
74	Groningen orthopaedic exit strategy: Validation of a support program after total hip or knee arthroplasty. <i>Patient Education and Counseling</i> , <b>2007</b> , 65, 171-9	3.1	12
73	Clinical outcomes and predictors of patient satisfaction in displaced midshaft clavicle fractures in adults: Results from a retrospective multicentre study. <i>Injury</i> , <b>2017</b> , 48, 2788-2792	2.5	11
72	Sport advice given by Dutch orthopaedic surgeons to patients after a total hip arthroplasty or total knee arthroplasty. <i>PLoS ONE</i> , <b>2018</b> , 13, e0202494	3.7	11
71	Evaluation of the effect of a comprehensive multidisciplinary care pathway for hip fractures: design of a controlled study. <i>BMC Musculoskeletal Disorders</i> , <b>2013</b> , 14, 291	2.8	10
70	A Dutch version of the Social Support for Exercise Behaviors Scale. <i>Perceptual and Motor Skills</i> , <b>2000</b> , 90, 771-4	2.2	10
69	Preoperative physical activity level has no relationship to the degree of recovery one year after primary total hip or knee arthroplasty: a cohort study. <i>PLoS ONE</i> , <b>2014</b> , 9, e115559	3.7	10

68	Duloxetine in OsteoArthritis (DOA) study: study protocol of a pragmatic open-label randomised controlled trial assessing the effect of preoperative pain treatment on postoperative outcome after total hip or knee arthroplasty. <i>BMJ Open</i> , <b>2016</b> , 6, e010343	3	10
67	Recruiting expertise: how surgical trainees engage supervisors for learning in the operating room. <i>Medical Education</i> , <b>2019</b> , 53, 616-627	3.7	9
66	Effectiveness of rehabilitation after a total hip arthroplasty: a protocol for an observational study for the comparison of usual care in the Netherlands versus Germany. <i>BMJ Open</i> , <b>2017</b> , 7, e016020	3	9
65	The effects of exercise and weight loss in overweight patients with hip osteoarthritis: design of a prospective cohort study. <i>BMC Musculoskeletal Disorders</i> , <b>2009</b> , 10, 24	2.8	9
64	Effectiveness of a Home-Based Rehabilitation Program After Total Hip Arthroplasty Driven by a Tablet App and Remote Coaching: Nonrandomized Controlled Trial Combining a Single-Arm Intervention Cohort With Historical Controls. <i>JMIR Rehabilitation and Assistive Technologies</i> , <b>2020</b> , 7, e14139	3.2	9
63	Intralesional treatment versus wide resection for central low-grade chondrosarcoma of the long bones. <i>The Cochrane Library</i> , <b>2019</b> , 3, CD010778	5.2	9
62	Evaluation of accuracy and precision of CT-guidance in Radiofrequency Ablation for osteoid osteoma in 86 patients. <i>PLoS ONE</i> , <b>2017</b> , 12, e0169171	3.7	8
61	Validation and User Evaluation of a Sensor-Based Method for Detecting Mobility-Related Activities in Older Adults. <i>PLoS ONE</i> , <b>2015</b> , 10, e0137668	3.7	8
60	Time spent in primary care for hip osteoarthritis patients once the diagnosis is set: a prospective observational study. <i>BMC Family Practice</i> , <b>2011</b> , 12, 48	2.6	8
59	Predictive value of the Western Ontario and McMaster Universities Osteoarthritis Index for the amount of physical activity after total hip arthroplasty. <i>Physical Therapy</i> , <b>2008</b> , 88, 211-8	3.3	8
58	A new interdisciplinary treatment strategy versus usual medical care for the treatment of subacromial impingement syndrome: a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , <b>2007</b> , 8, 15	2.8	8
57	The Validity of a New Low-Dose Stereoradiography System to Perform 2D and 3D Knee Prosthetic Alignment Measurements. <i>PLoS ONE</i> , <b>2016</b> , 11, e0146187	3.7	8
56	Managing persistent wound leakage after total knee and hip arthroplasty. Results of a nationwide survey among Dutch orthopaedic surgeons. <i>Journal of Bone and Joint Infection</i> , <b>2017</b> , 2, 202-207	2.7	7
55	Medication Use is a Better Predictor of Length of Hospital Stay in Total Hip Arthroplasty Than the American Society of Anesthetists (ASA) Score. <i>Journal of Arthroplasty</i> , <b>2017</b> , 32, 24-27	4.4	7
54	Do CAS measurements correlate with EOS 3D alignment measurements in primary TKA?. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2017</b> , 25, 2894-2903	5.5	6
53	Activity Impairment and Work Productivity Loss After Total Knee Arthroplasty: A Prospective Study. <i>Journal of Arthroplasty</i> , <b>2019</b> , 34, 2637-2645	4.4	6
52	Appropriate care for orthopedic patients: effect of implementation of the Clinical Practice Guideline for Diagnosis and Treatment of Subacromial Pain Syndrome in the Netherlands. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2019</b> , 90, 191-195	4.3	6
51	The Groningen Active Living Model, an example of successful recruitment of sedentary and underactive older adults. <i>Preventive Medicine</i> , <b>2008</b> , 47, 398-401	4.3	6

50	Differences in Physical-Fitness Test Scores between Actively and Passively Recruited Older Adults: Consequences for Norm-Based Classification. <i>Journal of Aging and Physical Activity</i> , <b>2002</b> , 10, 143-159	1.6	6
49	Validity of the Dutch modified painDETECT questionnaire for patients with hip or knee osteoarthritis. <i>Disability and Rehabilitation</i> , <b>2019</b> , 41, 941-947	2.4	6
48	No association between preoperative physical activity level and time to return to work in patients after total hip or knee arthroplasty: A prospective cohort study. <i>PLoS ONE</i> , <b>2019</b> , 14, e0221932	3.7	5
47	Preoperative characteristics of working-age patients undergoing total knee arthroplasty. <i>PLoS ONE</i> , <b>2017</b> , 12, e0183550	3.7	5
46	The influence of computer-assisted surgery on rotational, coronal and sagittal alignment in revision total knee arthroplasty. <i>BMC Musculoskeletal Disorders</i> , <b>2014</b> , 15, 94	2.8	5
45	Two-year recovery courses of physical and mental impairments, activity limitations, and participation restrictions after total knee arthroplasty among working-age patients. <i>Disability and Rehabilitation</i> , <b>2020</b> , 1-10	2.4	5
44	The Validity of the Circumduction Test in Elderly Men and Women. <i>Journal of Aging and Physical Activity</i> , <b>2003</b> , 11, 433-444	1.6	4
43	Background and Intensity of the GALM Physical Activity Program. <i>Journal of Physical Activity and Health</i> , <b>2005</b> , 2, 51-62	2.5	4
42	Validity and reliability of the Dutch version of the Copenhagen Hip And Groin Outcome Score (HAGOS-NL) in patients with hip pathology. <i>PLoS ONE</i> , <b>2017</b> , 12, e0186064	3.7	4
41	Implementing Individually Tailored Prescription of Physical Activity in Routine Clinical Care: Protocol of the Physicians Implement Exercise = Medicine (PIE=M) Development and Implementation Project. <i>JMIR Research Protocols</i> , <b>2020</b> , 9, e19397	2	4
40	The evolution of hand function during remodelling in nonreduced angulated paediatric forearm fractures: a prospective cohort study. <i>Journal of Pediatric Orthopaedics Part B</i> , <b>2020</b> , 29, 172-178	1.4	4
39	Which patients improve most after total hip arthroplasty? Influence of patient characteristics on patient-reported outcome measures of 22,357 total hip arthroplasties in the Dutch Arthroplasty Register. <i>HIP International</i> , <b>2021</b> , 31, 593-602	1.7	4
38	Actual and perceived nursing workload and the complexity of patients with total hip arthroplasty. <i>Applied Nursing Research</i> , <b>2018</b> , 39, 195-199	1.8	4
37	Remote timed up and go evaluation from activities of daily living reveals changing mobility after surgery. <i>Physiological Measurement</i> , <b>2019</b> , 40, 035004	2.9	3
36	Recovery of strength after reduced pediatric fractures of the forearm, wrist or hand; A prospective study. <i>PLoS ONE</i> , <b>2020</b> , 15, e0230862	3.7	3
35	Minimally invasive total hip and knee arthroplasty-implications for the elderly patient. <i>Clinics in Geriatric Medicine</i> , <b>2012</b> , 28, 447-58	3.8	3
34	People who undergo revision arthroplasty report more limitations but no decrease in physical activity compared with primary total hip arthroplasty: an observational study. <i>Australian Journal of Physiotherapy</i> , <b>2009</b> , 55, 185-9		3
33	A Dutch Translation of a Questionnaire Assessing Self-Efficacy in Leisure-Time Physical Activity. <i>Journal of Aging and Physical Activity</i> , <b>2001</b> , 9, 223-232	1.6	3

32	Development of an Exercise Program for the Frail Elderly. <i>Journal of Aging and Physical Activity</i> , <b>2001</b> , 9, 452-465	1.6	3
31	Compensatory Movement Patterns Are Based on Abnormal Activity of the Biceps Brachii and Posterior Deltoid Muscles in Patients with Symptomatic Rotator Cuff Tears. <i>Clinical Orthopaedics and Related Research</i> , <b>2021</b> , 479, 378-388	2.2	3
30	Radiofrequency ablation in the treatment of atypical cartilaginous tumours in the long bones: lessons learned from our experience. <i>Skeletal Radiology</i> , <b>2019</b> , 48, 881-887	2.7	3
29	Equity-Specific Effects of Interventions to Promote Physical Activity among Middle-Aged and Older Adults: Development of a Collaborative Equity-Specific Re-Analysis Strategy. <i>International Journal of Environmental Research and Public Health</i> , <b>2019</b> , 16,	4.6	2
28	The Groningen Enjoyment Questionnaire: A Measure of Enjoyment in Leisure-Time Physical Activity		2
27	A Dutch Version of the Social Support for Exercise Behaviors Scale		2
26	The Supervisor <sup>B</sup> Toolkit: Strategies of Supervisors to Entrust and Regulate Autonomy of Residents in the Operating Room. <i>Annals of Surgery</i> , <b>2020</b> , 275,	7.8	2
25	To mix or not to mix? Medicolegal implications of mixed components in total hip arthroplasty. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2020</b> , 91, 624-626	4.3	2
24	Effectiveness of an individually tailored home-based exercise programme for pre-frail older adults, driven by a tablet application and mobility monitoring: a pilot study. <i>European Review of Aging and Physical Activity</i> , <b>2021</b> , 18, 10	6.5	2
23	Strict adherence to evidence-based protocol in choice of implants and surgical technique leads to fewer hip fracture reoperations. <i>PLoS ONE</i> , <b>2019</b> , 14, e0210239	3.7	1
22	Long-term results of the iBP elbow prosthesis: beware of destructive metallosis!. <i>BMC Musculoskeletal Disorders</i> , <b>2019</b> , 20, 415	2.8	1
21	Adaptation, validity and reliability of the modified painDETECT questionnaire for patients with subacromial pain syndrome. <i>PLoS ONE</i> , <b>2019</b> , 14, e0211880	3.7	1
20	Barriers and facilitators perceived by healthcare professionals for implementing lifestyle interventions in patients with osteoarthritis: a scoping review.. <i>BMJ Open</i> , <b>2022</b> , 12, e056831	3	1
19	"What affects the implementation of lifestyle interventions in patients with osteoarthritis? A multidisciplinary focus group study among healthcare professionals". <i>Disability and Rehabilitation</i> , <b>2021</b> , 1-11	2.4	1
18	Effect of preoperative duloxetine treatment on postoperative chronic residual pain after total hip or knee arthroplasty: a randomised controlled trial. <i>BMJ Open</i> , <b>2021</b> , 11, e052944	3	1
17	Exploring patient satisfaction after operative and nonoperative treatment for midshaft clavicle fractures: a focus group analysis. <i>BMC Musculoskeletal Disorders</i> , <b>2020</b> , 21, 560	2.8	1
16	Residents Think in the "Now" and Supervisors Think Ahead in the Operating Room. A Survey Study About Task Perception of Residents and Supervising Surgeons. <i>Journal of Surgical Education</i> , <b>2021</b> , 78, 104-112	3.4	1
15	Fingerprints of Teaching Interactions: Capturing and Quantifying How Supervisor Regulate Autonomy of Residents in the Operating Room. <i>Journal of Surgical Education</i> , <b>2021</b> , 78, 1197-1208	3.4	1



14	Psychosocial Working Conditions Play an Important Role in the Return-to-Work Process After Total Knee and Hip Arthroplasty. <i>Journal of Occupational Rehabilitation</i> , <b>2021</b> , 1	3.6	1
13	Equity-specific effects of interventions to promote physical activity among middle-aged and older adults: results from applying a novel equity-specific re-analysis strategy. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , <b>2021</b> , 18, 65	8.4	0
12	A comprehensive multidisciplinary care pathway for hip fractures better outcome than usual care?. <i>Injury</i> , <b>2021</b> , 52, 1819-1825	2.5	0
11	Being active with a total hip or knee prosthesis: a systematic review into physical activity and sports recommendations and interventions to improve physical activity behavior.. <i>European Review of Aging and Physical Activity</i> , <b>2022</b> , 19, 7	6.5	0
10	Facilitators and barriers for the implementation of exercise as medicine in routine clinical care in Dutch university medical centres: a mixed methodology study on clinicians' perceptions.. <i>BMJ Open</i> , <b>2022</b> , 12, e052920	3	0
9	Amount and type of physical activity and sports from one year forward after hip or knee arthroplasty-A systematic review.. <i>PLoS ONE</i> , <b>2021</b> , 16, e0261784	3.7	0
8	Influence of social support on return to work after total hip or total knee arthroplasty: a prospective multicentre cohort study. <i>BMJ Open</i> , <b>2022</b> , 12, e059225	3	0
7	Decrease in heart rate after longitudinal participation in the Groningen Active Living Model (GALM) recreational sports programme. <i>Journal of Sports Sciences</i> , <b>2009</b> , 27, 975-83	3.6	
6	Test-retest reliability of a body-fixed sensor gait analysis protocol for evaluating trunk movements and gait parameters in patients with hip osteoarthritis. <i>Gait and Posture</i> , <b>2009</b> , 30, S124	2.6	
5	Attentional and visual demands for sprint performance in non-fatigued and fatigued conditions: reliability of a repeated sprint test. <i>BMC Musculoskeletal Disorders</i> , <b>2010</b> , 11, 84	2.8	
4	Artrose, sport en bewegen <b>2016</b> , 83-88		
3	11 Deelname aan een sport- en spelprogramma ter verbetering van de cardiorespiratoire fitheid bij ouderen <b>2011</b> , 163-180		
2	Prospective cohort study comparing a triceps-sparing and triceps-detaching approach in total elbow arthroplasty: a protocol. <i>BMJ Open</i> , <b>2021</b> , 11, e046098	3	
1	Validity and reliability of the German translation of the Diabetes Foot Self-Care Behavior Scale (DFSBS-D). <i>PLoS ONE</i> , <b>2022</b> , 17, e0269395	3.7	