

Run-Zhou Ni

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1526819/run-zhou-ni-publications-by-year.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

88

papers

1,583

citations

22

h-index

33

g-index

88

ext. papers

1,829

ext. citations

3.6

avg, IF

4.17

L-index

#	Paper	IF	Citations
88	MKP-4 suppresses hepatocarcinogenesis by targeting ERK1/2 pathway. <i>Cancer Cell International</i> , 2019 , 19, 61	6.4	7
87	Correlation between S100A11 and the TGF- β /SMAD4 pathway and its effects on the proliferation and apoptosis of pancreatic cancer cell line PANC-1. <i>Molecular and Cellular Biochemistry</i> , 2019 , 450, 53-64	4.2	4
86	CDK14 involvement in proliferation migration and invasion of esophageal cancer. <i>Annals of Translational Medicine</i> , 2019 , 7, 681	3.2	7
85	Actin related protein 3 (ARP3) promotes apoptosis of intestinal epithelial cells in ulcerative colitis. <i>Pathology Research and Practice</i> , 2019 , 215, 235-242	3.4	4
84	S100A11 promotes human pancreatic cancer PANC-1 cell proliferation and is involved in the PI3K/AKT signaling pathway. <i>Oncology Letters</i> , 2018 , 15, 175-182	2.6	18
83	Runt-related transcription factor 2 (RUNX2) inhibits apoptosis of intestinal epithelial cells in Crohn's disease. <i>Pathology Research and Practice</i> , 2018 , 214, 245-252	3.4	3
82	α -AR regulates the expression of AKR1B1 in human pancreatic cancer cells and promotes their proliferation via the ERK1/2 pathway. <i>Molecular Biology Reports</i> , 2018 , 45, 1863-1871	2.8	12
81	High NUSAP1 expression predicts poor prognosis in colon cancer. <i>Pathology Research and Practice</i> , 2018 , 214, 968-973	3.4	22
80	Modification of p27 with O-linked N-acetylglucosamine regulates cell proliferation in hepatocellular carcinoma. <i>Molecular Carcinogenesis</i> , 2017 , 56, 258-271	5	9
79	Long non-coding RNA SNHG20 predicts a poor prognosis for HCC and promotes cell invasion by regulating the epithelial-to-mesenchymal transition. <i>Biomedicine and Pharmacotherapy</i> , 2017 , 89, 857-863	7.5	70
78	Chromosome region maintenance-1 (CRM1) regulates apoptosis of intestinal epithelial cells via p27kip1 in Crohn's disease. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2017 , 41, 445-458	2.4	2
77	High Expression of BCCIP Can Promote Proliferation of Esophageal Squamous Cell Carcinoma. <i>Digestive Diseases and Sciences</i> , 2017 , 62, 387-395	4	3
76	PRDM5 promotes the apoptosis of epithelial cells induced by IFN- γ during Crohn's disease. <i>Pathology Research and Practice</i> , 2017 , 213, 666-673	3.4	7
75	Sex comb on midleg like-2 is a novel specific marker for the diagnosis of gastroenteropancreatic neuroendocrine tumors. <i>Experimental and Therapeutic Medicine</i> , 2017 , 14, 1749-1755	2.1	1
74	Upregulated TRIM32 correlates with enhanced cell proliferation and poor prognosis in hepatocellular carcinoma. <i>Molecular and Cellular Biochemistry</i> , 2016 , 421, 127-37	4.2	22
73	Hyper-O-GlcNAcylation of YB-1 affects Ser102 phosphorylation and promotes cell proliferation in hepatocellular carcinoma. <i>Experimental Cell Research</i> , 2016 , 349, 230-238	4.2	24
72	Expression pattern of BCCIP in hepatocellular carcinoma is correlated with poor prognosis and enhanced cell proliferation. <i>Tumor Biology</i> , 2016 , 37, 16305	2.9	6

71	Overexpression of DLX2 is associated with poor prognosis and sorafenib resistance in hepatocellular carcinoma. <i>Experimental and Molecular Pathology</i> , 2016 , 101, 58-65	4.4	16
70	Low expression of PIDD is associated with cell proliferation and apoptosis in hepatocellular carcinoma. <i>Tumor Biology</i> , 2016 , 37, 10447-57	2.9	2
69	The DNA damage repair protein Ku70 regulates tumor cell and hepatic carcinogenesis by interacting with FOXO4. <i>Pathology Research and Practice</i> , 2016 , 212, 153-61	3.4	10
68	Far upstream element-binding protein 1 (FUBP1) is a potential c-Myc regulator in esophageal squamous cell carcinoma (ESCC) and its expression promotes ESCC progression. <i>Tumor Biology</i> , 2016 , 37, 4115-26	2.9	18
67	Expression and Clinical Role of Cdc5L as a Novel Cell Cycle Protein in Hepatocellular Carcinoma. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 795-805	4	15
66	Downregulated Expression of PTPN9 Contributes to Human Hepatocellular Carcinoma Growth and Progression. <i>Pathology and Oncology Research</i> , 2016 , 22, 555-65	2.6	11
65	Enhanced expression of early mitotic inhibitor-1 predicts a poor prognosis in esophageal squamous cell carcinoma patients. <i>Oncology Letters</i> , 2016 , 12, 114-120	2.6	6
64	GAB2 promotes cell proliferation by activating the ERK signaling pathway in hepatocellular carcinoma. <i>Tumor Biology</i> , 2016 , 37, 11763-11773	2.9	20
63	Decreased Expression of EHD2 Promotes Tumor Metastasis and Indicates Poor Prognosis in Hepatocellular Carcinoma. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 2554-67	4	12
62	The suppressor of cytokine signaling SOCS1 promotes apoptosis of intestinal epithelial cells via p53 signaling in Crohn's disease. <i>Experimental and Molecular Pathology</i> , 2016 , 101, 1-11	4.4	15
61	Breaking Hepatitis B Virus Tolerance and Inducing Protective Immunity Based on Mimicking T Cell-Independent Antigen. <i>Viral Immunology</i> , 2016 , 29, 502-509	1.7	
60	PKC iota promotes cellular proliferation by accelerated G1/S transition via interaction with CDK7 in esophageal squamous cell carcinoma. <i>Tumor Biology</i> , 2016 , 37, 13799-13809	2.9	9
59	High CHMP4B expression is associated with accelerated cell proliferation and resistance to doxorubicin in hepatocellular carcinoma. <i>Tumor Biology</i> , 2015 , 36, 2569-81	2.9	10
58	Downregulation of FOXP2 promoter human hepatocellular carcinoma cell invasion. <i>Tumor Biology</i> , 2015 , 36, 9611-9	2.9	24
57	Sam68 promotes cellular proliferation and predicts poor prognosis in esophageal squamous cell carcinoma. <i>Tumor Biology</i> , 2015 , 36, 8735-45	2.9	6
56	Upregulated HOXC8 Expression Is Associated with Poor Prognosis and Oxaliplatin Resistance in Hepatocellular Carcinoma. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 3351-63	4	15
55	High expression of vacuolar protein sorting 4B (VPS4B) is associated with accelerated cell proliferation and poor prognosis in human hepatocellular carcinoma. <i>Pathology Research and Practice</i> , 2015 , 211, 240-7	3.4	13
54	Expression and clinical role of NF45 as a novel cell cycle protein in esophageal squamous cell carcinoma (ESCC). <i>Tumor Biology</i> , 2015 , 36, 747-56	2.9	32

53	Interaction with CCNH/CDK7 facilitates CtBP2 promoting esophageal squamous cell carcinoma (ESCC) metastasis via upregulating epithelial-mesenchymal transition (EMT) progression. <i>Tumor Biology</i> , 2015 , 36, 6701-14	2.9	19
52	Epithelial-specific ETS-1 (ESE1/ELF3) regulates apoptosis of intestinal epithelial cells in ulcerative colitis via accelerating NF- κ B activation. <i>Immunologic Research</i> , 2015 , 62, 198-212	4.3	18
51	Overexpression of SYF2 correlates with enhanced cell growth and poor prognosis in human hepatocellular carcinoma. <i>Molecular and Cellular Biochemistry</i> , 2015 , 410, 1-9	4.2	14
50	Vacuolar protein sorting 4B regulates apoptosis of intestinal epithelial cells via p38 MAPK in Crohn's disease. <i>Experimental and Molecular Pathology</i> , 2015 , 98, 55-64	4.4	13
49	Pyruvate kinase M2 regulates apoptosis of intestinal epithelial cells in Crohn's disease. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 393-404	4	23
48	The RNA-binding protein Sam68 regulates tumor cell viability and hepatic carcinogenesis by inhibiting the transcriptional activity of FOXOs. <i>Journal of Molecular Histology</i> , 2015 , 46, 485-97	3.3	9
47	Transplantation of mouse embryonic stem cell-derived oligodendrocytes in the murine model of globoid cell leukodystrophy. <i>Stem Cell Research and Therapy</i> , 2015 , 6, 30	8.3	16
46	Increased expression of glycinamide ribonucleotide transformylase is associated with a poor prognosis in hepatocellular carcinoma, and it promotes liver cancer cell proliferation. <i>Human Pathology</i> , 2014 , 45, 1370-8	3.7	15
45	Identification of Gem as a new candidate prognostic marker in hepatocellular carcinoma. <i>Pathology Research and Practice</i> , 2014 , 210, 719-25	3.4	5
44	Chronic pouchitis is associated with pouch polyp formation in patients with underlying ulcerative colitis. <i>Journal of Crohns and Colitis</i> , 2014 , 8, 363-9	1.5	10
43	High expression of peroxiredoxin 4 affects the survival time of colorectal cancer patients, but is not an independent unfavorable prognostic factor. <i>Molecular and Clinical Oncology</i> , 2014 , 2, 767-772	1.6	17
42	Upregulation of SYF2 in esophageal squamous cell carcinoma promotes tumor cell proliferation and predicts poor prognosis. <i>Tumor Biology</i> , 2014 , 35, 10275-85	2.9	5
41	S100 family signaling network and related proteins in pancreatic cancer (Review). <i>International Journal of Molecular Medicine</i> , 2014 , 33, 769-76	4.4	48
40	The expression levels and prognostic value of high temperature required A2 (HtrA2) in NSCLC. <i>Pathology Research and Practice</i> , 2014 , 210, 939-43	3.4	10
39	The downregulation of ErbB3 binding protein 1 (EBP1) is associated with poor prognosis and enhanced cell proliferation in hepatocellular carcinoma. <i>Molecular and Cellular Biochemistry</i> , 2014 , 396, 175-85	4.2	17
38	Effect of combined mutations in the enhancer II and basal core promoter of hepatitis B virus on development of hepatocellular carcinoma in Qidong, China. <i>Hepatology Research</i> , 2014 , 44, 1186-95	5.1	13
37	Upregulated expression of CAP1 is associated with tumor migration and metastasis in hepatocellular carcinoma. <i>Pathology Research and Practice</i> , 2014 , 210, 169-75	3.4	25
36	Expression of SGTA correlates with prognosis and tumor cell proliferation in human hepatocellular carcinoma. <i>Pathology and Oncology Research</i> , 2014 , 20, 51-60	2.6	14

35	Polycomb group oncogene RING1 is over-expressed in non-small cell lung cancer. <i>Pathology and Oncology Research</i> , 2014 , 20, 549-56	2.6	8
34	Expression of FOXJ1 in hepatocellular carcinoma: correlation with patientsSprognosis and tumor cell proliferation. <i>Molecular Carcinogenesis</i> , 2013 , 52, 647-59	5	33
33	Vacuolar protein sorting 4B, an ATPase protein positively regulates the progression of NSCLC via promoting cell division. <i>Molecular and Cellular Biochemistry</i> , 2013 , 381, 163-71	4.2	20
32	High SKIP expression is correlated with poor prognosis and cell proliferation of hepatocellular carcinoma. <i>Medical Oncology</i> , 2013 , 30, 537	3.7	8
31	Expression and clinical role of small glutamine-rich tetratricopeptide repeat (TPR)-containing protein alpha (SGTA) as a novel cell cycle protein in NSCLC. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 1539-49	4.9	15
30	Early mitotic inhibitor-1, an anaphase-promoting complex/cyclosome inhibitor, can control tumor cell proliferation in hepatocellular carcinoma: correlation with Skp2 stability and degradation of p27(Kip1). <i>Human Pathology</i> , 2013 , 44, 365-73	3.7	25
29	Effects of hepatitis B e-antigen on recurrence of hepatitis B-related hepatocellular carcinoma after curative resection: A meta-analysis. <i>Hepatology Research</i> , 2013 , 43, 347-54	5.1	5
28	Epithelial membrane protein 3 is frequently shown as promoter methylation and functions as a tumor suppressor gene in non-small cell lung cancer. <i>Experimental and Molecular Pathology</i> , 2013 , 95, 313-8	4.4	23
27	Decreased expression and prognostic role of mitogen-activated protein kinase phosphatase 4 in hepatocellular carcinoma. <i>Journal of Gastrointestinal Surgery</i> , 2013 , 17, 756-65	3.3	14
26	Increased Eubulin1b expression indicates poor prognosis and resistance to chemotherapy in hepatocellular carcinoma. <i>Digestive Diseases and Sciences</i> , 2013 , 58, 2713-20	4	34
25	Glyoxylate reductase/hydroxypyruvate reductase: a novel prognostic marker for hepatocellular carcinoma patients after curative resection. <i>Pathobiology</i> , 2013 , 80, 155-62	3.6	3
24	CtBP2 contributes to malignant development of human esophageal squamous cell carcinoma by regulation of p16INK4A. <i>Journal of Cellular Biochemistry</i> , 2013 , 114, 1343-54	4.7	25
23	Overexpressed nuclear BAG-1 in human hepatocellular carcinoma is associated with poor prognosis and resistance to doxorubicin. <i>Journal of Cellular Biochemistry</i> , 2013 , 114, 2120-30	4.7	17
22	Combined analysis of serum Eglutamyl transferase isoenzyme II, E-fucosidase and Efetoprotein detected using a commercial kit in the diagnosis of hepatocellular carcinoma. <i>Experimental and Therapeutic Medicine</i> , 2013 , 5, 89-94	2.1	17
21	Serum GP73 is complementary to AFP and GGT-II for the diagnosis of hepatocellular carcinoma. <i>Oncology Letters</i> , 2013 , 6, 1152-1158	2.6	25
20	Clinical significance of indefinite for dysplasia on pouch biopsy in patients with underlying inflammatory bowel disease. <i>Journal of Gastrointestinal Surgery</i> , 2012 , 16, 562-71	3.3	13
19	High expression of S100A11 in pancreatic adenocarcinoma is an unfavorable prognostic marker. <i>Medical Oncology</i> , 2012 , 29, 1886-91	3.7	38
18	Ectopic expression of guanylyl cyclase C and endogenous ligand guanylin correlates significantly with Helicobacter pylori infection in gastric carcinogenesis. <i>Medical Oncology</i> , 2012 , 29, 1748-57	3.7	2

17	Pre-colectomy appendectomy and risk for Crohn's disease in patients with ileal pouch-anal anastomosis. <i>Journal of Gastrointestinal Surgery</i> , 2012 , 16, 1370-8	3.3	6
16	Clinical and biological significance of never in mitosis gene A-related kinase 6 (NEK6) expression in hepatic cell cancer. <i>Pathology and Oncology Research</i> , 2012 , 18, 201-7	2.6	20
15	The expressions and clinical significances of tissue and serum galectin-3 in pancreatic carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2012 , 138, 1035-43	4.9	49
14	Expression of Pirh2, a p27(Kip1) ubiquitin ligase, in hepatocellular carcinoma: correlation with p27(Kip1) and cell proliferation. <i>Human Pathology</i> , 2011 , 42, 507-15	3.7	24
13	The presence of IGHG1 in human pancreatic carcinomas is associated with immune evasion mechanisms. <i>Pancreas</i> , 2011 , 40, 753-61	2.6	30
12	Transmural inflammation is not pathognomonic for Crohn's disease of the pouch. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2011 , 25, 3509-17	5.2	20
11	Diagnosis and management of dysplasia and cancer of the ileal pouch in patients with underlying inflammatory bowel disease. <i>Cancer</i> , 2011 , 117, 3081-92	6.4	45
10	Ileal pouch for everyone, even when we are not sure of the diagnosis before or at colectomy?. <i>Inflammatory Bowel Diseases</i> , 2010 , 16, 716-8	4.5	
9	Comparative proteomic analysis of differentially expressed proteins in human pancreatic cancer tissue. <i>Hepatobiliary and Pancreatic Diseases International</i> , 2009 , 8, 193-200	2.1	27
8	Correlation between thymidylate synthase genotype and susceptibility to gastric carcinoma. <i>Chinese Journal of Clinical Oncology</i> , 2008 , 5, 448-452		2
7	Clinical value of hepatoma-specific alpha-fetoprotein in the diagnosis of hepatocellular carcinoma. <i>Chinese Journal of Clinical Oncology</i> , 2006 , 3, 153-157		
6	Anti-hepatoma effect of arsenic trioxide on experimental liver cancer induced by 2-acetamidofluorene in rats. <i>World Journal of Gastroenterology</i> , 2005 , 11, 5938-43	5.6	18
5	Glycylproline dipeptidyl aminopeptidase isoenzyme in diagnosis of primary hepatocellular carcinoma. <i>World Journal of Gastroenterology</i> , 2003 , 9, 710-3	5.6	8
4	Cell growth inhibition by a novel vitamin K is associated with induction of protein tyrosine phosphorylation. <i>Journal of Biological Chemistry</i> , 1998 , 273, 9906-11	5.4	42
3	Molecular cloning of two types of cDNA encoding subunit RC6-I of rat proteasomes. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1995 , 1264, 45-52		15
2	Long-term maintenance of functional rat hepatocytes in primary culture by additions of pyruvate and various hormones. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1995 , 1243, 329-35	4	14
1	Fas-mediated apoptosis in primary cultured mouse hepatocytes. <i>Experimental Cell Research</i> , 1994 , 215, 332-7	4.2	192