

Noriyuki Matsuda, æ³/4ç”°æ²ä¹<

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1532973/publications.pdf>

Version: 2024-02-01

50
papers

7,825
citations

136740

32
h-index

182168

51
g-index

52
all docs

52
docs citations

52
times ranked

10469
citing authors

#	ARTICLE	IF	CITATIONS
1	Molecular mechanisms and physiological functions of mitophagy. <i>EMBO Journal</i> , 2021, 40, e104705.	3.5	553
2	Cleaved PGAM5 dephosphorylates nuclear serine/arginine-rich proteins during mitophagy. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2021, 1868, 119045.	1.9	2
3	Loss of peptide: <i>N</i> -glycanase causes proteasome dysfunction mediated by a sugar-recognizing ubiquitin ligase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	23
4	Mammalian BCAS3 and C16orf70 associate with the phagophore assembly site in response to selective and non-selective autophagy. <i>Autophagy</i> , 2021, 17, 2011-2036.	4.3	6
5	Unfolding is the driving force for mitochondrial import and degradation of the Parkinson's disease-related protein DJ-1. <i>Journal of Cell Science</i> , 2021, 134, .	1.2	3
6	Two sides of a coin: Physiological significance and molecular mechanisms for damage-induced mitochondrial localization of PINK1 and Parkin. <i>Neuroscience Research</i> , 2020, 159, 16-24.	1.0	8
7	Critical role of mitochondrial ubiquitination and the OPTN-ATG9A axis in mitophagy. <i>Journal of Cell Biology</i> , 2020, 219, .	2.3	114
8	Parkin recruitment to impaired mitochondria for nonselective ubiquitylation is facilitated by MITOL. <i>Journal of Biological Chemistry</i> , 2019, 294, 10300-10314.	1.6	79
9	Parkin-mediated ubiquitylation redistributes MITOL/March5 from mitochondria to peroxisomes. <i>EMBO Reports</i> , 2019, 20, e47728.	2.0	35
10	Cleaved PGAM5 is released from mitochondria depending on proteasome-mediated rupture of the outer mitochondrial membrane during mitophagy. <i>Journal of Biochemistry</i> , 2019, 165, 19-25.	0.9	19
11	Endosomal Rab cycles regulate Parkin-mediated mitophagy. <i>ELife</i> , 2018, 7, .	2.8	113
12	Discovery and Optimization of Inhibitors of the Parkinson's Disease Associated Protein DJ-1. <i>ACS Chemical Biology</i> , 2018, 13, 2783-2793.	1.6	27
13	Structural insights into ubiquitin phosphorylation by PINK1. <i>Scientific Reports</i> , 2018, 8, 10382.	1.6	35
14	Parkinson's disease-related DJ-1 functions in thiol quality control against aldehyde attack in vitro. <i>Scientific Reports</i> , 2017, 7, 12816.	1.6	41
15	Structural basis for specific cleavage of Lys6-linked polyubiquitin chains by USP30. <i>Nature Structural and Molecular Biology</i> , 2017, 24, 911-919.	3.6	61
16	Ubiquitination of exposed glycoproteins by SCF ^{FBXO27} directs damaged lysosomes for autophagy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 8574-8579.	3.3	96
17	The ubiquitin signal and autophagy: an orchestrated dance leading to mitochondrial degradation. <i>EMBO Reports</i> , 2016, 17, 300-316.	2.0	197
18	Unexpected mitochondrial matrix localization of Parkinson's disease-related DJ-1 mutants but not wild-type DJ-1. <i>Genes To Cells</i> , 2016, 21, 772-788.	0.5	21

#	ARTICLE	IF	CITATIONS
19	Constitutive Activation of PINK1 Protein Leads to Proteasome-mediated and Non-apoptotic Cell Death Independently of Mitochondrial Autophagy. <i>Journal of Biological Chemistry</i> , 2016, 291, 16162-16174.	1.6	23
20	Phospho-ubiquitin: upending the PINK1/Parkin ubiquitin cascade. <i>Journal of Biochemistry</i> , 2016, 159, 379-385.	0.9	53
21	Unconventional PINK1 localization mechanism to the outer membrane of depolarized mitochondria drives Parkin recruitment. <i>Journal of Cell Science</i> , 2015, 128, 964-78.	1.2	103
22	Molecular mechanisms underlying PINK1 and Parkin catalyzed ubiquitylation of substrates on damaged mitochondria. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2015, 1853, 2791-2796.	1.9	35
23	The PARK2/Parkin receptor on damaged mitochondria revisited—uncovering the role of phosphorylated ubiquitin chains. <i>Autophagy</i> , 2015, 11, 1700-1701.	4.3	6
24	Phosphorylated ubiquitin chain is the genuine Parkin receptor. <i>Journal of Cell Biology</i> , 2015, 209, 111-128.	2.3	217
25	Tagged tags engage disposal. <i>Nature</i> , 2015, 524, 294-295.	13.7	6
26	Site-specific Interaction Mapping of Phosphorylated Ubiquitin to Uncover Parkin Activation. <i>Journal of Biological Chemistry</i> , 2015, 290, 25199-25211.	1.6	50
27	Ubiquitin is phosphorylated by PINK1 to activate parkin. <i>Nature</i> , 2014, 510, 162-166.	13.7	1,185
28	Proteostasis and neurodegeneration: The roles of proteasomal degradation and autophagy. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2014, 1843, 197-204.	1.9	153
29	Parkin-catalyzed Ubiquitin-Ester Transfer Is Triggered by PINK1-dependent Phosphorylation. <i>Journal of Biological Chemistry</i> , 2013, 288, 22019-22032.	1.6	173
30	A Dimeric PINK1-containing Complex on Depolarized Mitochondria Stimulates Parkin Recruitment. <i>Journal of Biological Chemistry</i> , 2013, 288, 36372-36384.	1.6	168
31	Different dynamic movements of wild-type and pathogenic VCPs and their cofactors to damaged mitochondria in a Parkin-mediated mitochondrial quality control system. <i>Genes To Cells</i> , 2013, 18, 1131-1143.	0.5	35
32	The principal PINK1 and Parkin cellular events triggered in response to dissipation of mitochondrial membrane potential occur in primary neurons. <i>Genes To Cells</i> , 2013, 18, 672-681.	0.5	38
33	PINK1 autophosphorylation upon membrane potential dissipation is essential for Parkin recruitment to damaged mitochondria. <i>Nature Communications</i> , 2012, 3, 1016.	5.8	465
34	Mitochondrial hexokinase HKI is a novel substrate of the Parkin ubiquitin ligase. <i>Biochemical and Biophysical Research Communications</i> , 2012, 428, 197-202.	1.0	65
35	Parkin Mediates Apparent E2-Independent Monoubiquitination In Vitro and Contains an Intrinsic Activity That Catalyzes Polyubiquitination. <i>PLoS ONE</i> , 2011, 6, e19720.	1.1	40
36	p62/SQSTM1 cooperates with Parkin for perinuclear clustering of depolarized mitochondria. <i>Genes To Cells</i> , 2010, 15, 887-900.	0.5	345

#	ARTICLE	IF	CITATIONS
37	PINK1 stabilized by mitochondrial depolarization recruits Parkin to damaged mitochondria and activates latent Parkin for mitophagy. <i>Journal of Cell Biology</i> , 2010, 189, 211-221.	2.3	1,600
38	Uncovering the roles of PINK1 and Parkin in mitophagy. <i>Autophagy</i> , 2010, 6, 952-954.	4.3	41
39	Does Impairment of the Ubiquitin-Proteasome System or the Autophagy-Lysosome Pathway Predispose Individuals to Neurodegenerative Disorders such as Parkinson's Disease?. <i>Journal of Alzheimer's Disease</i> , 2010, 19, 1-9.	1.2	89
40	MG53 nucleates assembly of cell membrane repair machinery. <i>Nature Cell Biology</i> , 2009, 11, 56-64.	4.6	396
41	Direct interactions between NEDD8 and ubiquitin E2 conjugating enzymes upregulate cullin-based E3 ligase activity. <i>Nature Structural and Molecular Biology</i> , 2007, 14, 167-168.	3.6	105
42	Diverse Effects of Pathogenic Mutations of Parkin That Catalyze Multiple Monoubiquitylation in Vitro. <i>Journal of Biological Chemistry</i> , 2006, 281, 3204-3209.	1.6	166
43	UV-Induced Ubiquitylation of XPC Protein Mediated by UV-DDB-Ubiquitin Ligase Complex. <i>Cell</i> , 2005, 121, 387-400.	13.5	517
44	DDB2, the xeroderma pigmentosum group E gene product, is directly ubiquitylated by Cullin 4A-based ubiquitin ligase complex. <i>DNA Repair</i> , 2005, 4, 537-545.	1.3	65
45	A palmitoylated RING finger ubiquitin ligase and its homologue in the brain membranes. <i>Journal of Neurochemistry</i> , 2003, 86, 749-762.	2.1	25
46	Ubiquitin Ligase Activities of Bombyx mori Nucleopolyhedrovirus RING Finger Proteins. <i>Journal of Virology</i> , 2003, 77, 923-930.	1.5	69
47	EL5, a rice N-acetylchitoooligosaccharide elicitor-responsive RING-H2 finger protein, is a ubiquitin ligase which functions in vitro in co-operation with an elicitor-responsive ubiquitin-conjugating enzyme, OsUBC5b. <i>Plant Journal</i> , 2002, 30, 447-455.	2.8	98
48	Modes of interaction between the Arabidopsis Rab protein, Ara4, and its putative regulator molecules revealed by a yeast expression system. <i>Plant Journal</i> , 2000, 21, 341-349.	2.8	21
49	Overexpression of PRA2, a Rab/Yipt-family Small GTPase from Pea <i>Pisum sativum</i> , Aggravates the Growth Defect of Yeast ypt Mutants.. <i>Cell Structure and Function</i> , 2000, 25, 11-20.	0.5	9
50	RMA1 an Arabidopsis thaliana Gene Whose cDNA Suppresses the Yeast <i>secl5</i> Mutation, Encodes a Novel Protein with a RING Finger Motif and a Membrane Anchor. <i>Plant and Cell Physiology</i> , 1998, 39, 545-554.	1.5	27