

# Satoru Tsuneto

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1542549/publications.pdf>

Version: 2024-02-01

65  
papers

794  
citations

759233

12  
h-index

552781

26  
g-index

67  
all docs

67  
docs citations

67  
times ranked

743  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Does Urinary Catheterization Affect the Quality of Death in Patients with Advanced Cancer? A Secondary Analysis of a Multicenter Prospective Cohort Study. <i>Journal of Palliative Medicine</i> , 2022, 25, 205-212.  | 1.1 | 2         |
| 2  | Assessment of Changes in Symptoms Is Feasible and Prognostic in the Last Weeks of Life: An International Multicenter Cohort Study. <i>Journal of Palliative Medicine</i> , 2022, 25, 388-395.  | 1.1 | 4         |
| 3  | Comparison of the accuracy of clinicians' prediction of survival and Palliative Prognostic Score: an East Asian cross-cultural study. <i>Supportive Care in Cancer</i> , 2022, 30, 2367-2374.  | 2.2 | 5         |
| 4  | Family experience of palliative sedation therapy: proportional vs. continuous deep sedation. <i>Supportive Care in Cancer</i> , 2022, 30, 3903-3915.   | 2.2 | 3         |
| 5  | Association between the amount of artificial hydration and quality of dying among terminally ill patients with cancer: The East Asian Collaborative Cross-Cultural Study to Elucidate the Dying Process. <i>Cancer</i> , 2022, 128, 1699-1708.   | 4.1 | 2         |
| 6  | Defining "Continuous Deep Sedation" Using Treatment Protocol: A Proposal Article. <i>Palliative Medicine Reports</i> , 2022, 3, 8-15.  | 0.9 | 1         |
| 7  | Association between temporary discharge from the inpatient palliative care unit and achievement of good death in end-of-life cancer patients: A nationwide survey of bereaved family members. <i>Japan Journal of Nursing Science</i> , 2022, 19, e12474.  | 1.3 | 1         |
| 8  | Association between experiences of advanced cancer patients at the end of life and depression in their bereaved caregivers. <i>Psycho-Oncology</i> , 2022, 31, 1243-1252.  | 2.3 | 5         |
| 9  | Prevalence and severity of symptoms and signs in patients with advanced cancer in the last days of life: the East Asian collaborative cross-cultural study to elucidate the dying process (EASED). <i>Supportive Care in Cancer</i> , 2022, 30, 5499-5508.   | 2.2 | 6         |
| 10 | Association of the RASS Score with Intensity of Symptoms, Discomfort, and Communication Capacity in Terminally Ill Cancer Patients Receiving Palliative Sedation: Is RASS an Appropriate Outcome Measure?. <i>Palliative Medicine Reports</i> , 2022, 3, 47-54.  | 0.9 | 3         |
| 11 | The impact of death rattle on bereaved families: not the sound itself, but the resonance with their feelings. <i>Japanese Journal of Clinical Oncology</i> , 2022, 52, 774-778.  | 1.3 | 1         |
| 12 | Communication and Behavior of Palliative Care Physicians of Patients With Cancer Near End of Life in Three East Asian Countries. <i>Journal of Pain and Symptom Management</i> , 2021, 61, 315-322.e1.   | 1.2 | 36        |
| 13 | Predicting models of depression or complicated grief among bereaved family members of patients with cancer. <i>Psycho-Oncology</i> , 2021, 30, 1151-1159.  | 2.3 | 4         |
| 14 | Complexity of desire for hastened death in terminally ill cancer patients: A cluster analysis. <i>Palliative and Supportive Care</i> , 2021, 19, 646-655.  | 1.0 | 6         |
| 15 | Are cancer patients living alone more or less likely to achieve a good death? Two cross-sectional surveys of bereaved families. <i>Journal of Advanced Nursing</i> , 2021, 77, 3745-3758.  | 3.3 | 4         |
| 16 | The Association of Family Functioning With Possible Major Depressive Disorders and Complicated Grief Among Bereaved Family Members of Patients With Cancer: Results From the J-HOPE4 Study, a Nationwide Cross-Sectional Follow-Up Survey in Japan. <i>Journal of Pain and Symptom Management</i> , 2021, 62, 1154-1164. | 1.2 | 3         |
| 17 | Factors related to spiritual well-being in the last days of life in three East Asian countries: An international multicenter prospective cohort study. <i>Palliative Medicine</i> , 2021, 35, 1564-1577.   | 3.1 | 3         |
| 18 | Are family relationships associated with family conflict in advanced cancer patients?. <i>Psycho-Oncology</i> , 2021, , .  | 2.3 | 1         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Differences between dedicated and not dedicated hospice physicians in symptoms and signs improvement among advanced cancer patients. <i>Medicine (United States)</i> , 2021, 100, e26915.   | 1.0 | 2         |
| 20 | Physician's Communication in Code Status Discussions for Terminally Ill Cancer Patients in Inpatient Hospice/Palliative Care Units in Japan: A Nationwide Post-Bereavement Survey. <i>Journal of Pain and Symptom Management</i> , 2021, 62, e120-e129. | 1.2 | 2         |
| 21 | Development of a Scoring System to Determine Proportional Appropriateness of Continuous Deep Sedation: A Concept-of-Proof Study. <i>Journal of Palliative Medicine</i> , 2021, 24, 1539-1544.   | 1.1 | 0         |
| 22 | Care Associated With Satisfaction of Bereaved Family Members of Terminally Ill Cancer Patients With Dyspnea: A Cross-sectional Nationwide Survey. <i>Journal of Pain and Symptom Management</i> , 2021, 62, 796-804.                                    | 1.2 | 2         |
| 23 | Differences among institutions in the prevalence of and indications for urinary catheterization of advanced cancer patients at palliative care units: A multicenter prospective cohort study (EASED). <i>Palliative Medicine</i> , 2021, 35, 799-805.   | 3.1 | 3         |
| 24 | The Bereaved Families' Preferences for Individualized Goals of Care for Terminal Dyspnea: What Is an Acceptable Balance between Dyspnea Intensity and Communication Capacity?. <i>Palliative Medicine Reports</i> , 2020, 1, 42-49.                     | 0.9 | 7         |
| 25 | Continuous deep sedation at the end of life in children with cancer: experience at a single center in Japan. <i>Pediatric Hematology and Oncology</i> , 2020, 37, 365-374.  | 0.8 | 6         |
| 26 | Insomnia and changes in alcohol consumption: Relation between possible complicated grief and depression among bereaved family caregivers. <i>Journal of Affective Disorders</i> , 2020, 275, 1-6.   | 4.1 | 9         |
| 27 | Achievement of a good death among young adult patients with cancer: analyses of combined data from three nationwide surveys among bereaved family members. <i>Supportive Care in Cancer</i> , 2019, 27, 1519-1527.                                      | 2.2 | 8         |
| 28 | Changes in opinions on palliative sedation of palliative care specialists over 16 years and their effects on clinical practice. <i>Supportive Care in Cancer</i> , 2019, 27, 2211-2219.   | 2.2 | 10        |
| 29 | Difference in Opinions About Continuous Deep Sedation Among Cancer Patients, Bereaved Families, and Physicians. <i>Journal of Pain and Symptom Management</i> , 2019, 57, e5-e9.  | 1.2 | 0         |
| 30 | Japanese physicians' experiences of terminally ill patients voluntarily stopping eating and drinking: a national survey. <i>BMJ Supportive and Palliative Care</i> , 2019, 9, 143-145.  | 1.6 | 16        |
| 31 | Communication Disparity Between the Bereaved and Others: What Hurts Them and What Is Unhelpful? A Nationwide Study of the Cancer Bereaved. <i>Journal of Pain and Symptom Management</i> , 2018, 55, 1061-1067.e1.                                      | 1.2 | 5         |
| 32 | A double-blind, randomized comparative study to investigate the morphine to hydromorphone conversion ratio in Japanese cancer patients. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 442-449.   | 1.3 | 8         |
| 33 | Families' Sense of Abandonment When Patients Are Referred to Hospice. <i>Oncologist</i> , 2018, 23, 1109-1115.  | 3.7 | 6         |
| 34 | End-of-Life Care and Discussions in Japanese Geriatric Health Service Facilities: A Nationwide Survey of Managing Directors' Viewpoints. <i>American Journal of Hospice and Palliative Medicine</i> , 2018, 35, 83-91.                                  | 1.4 | 7         |
| 35 | End-of-life experiences of family caregivers of deceased patients with cancer: A nationwide survey. <i>Psycho-Oncology</i> , 2018, 27, 272-278.   | 2.3 | 6         |
| 36 | Which Research Questions Are Important for the Bereaved Families of Palliative Care Cancer Patients? A Nationwide Survey. <i>Journal of Pain and Symptom Management</i> , 2018, 55, 379-386.  | 1.2 | 5         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Are Bereaved Family Members Satisfied With Information Provision About Palliative Care Units in Japan?. <i>American Journal of Hospice and Palliative Medicine</i> , 2018, 35, 275-283.   | 1.4 | 2         |
| 38 | “What I Did for My Loved One Is More Important than Whether We Talked About Death”: A Nationwide Survey of Bereaved Family Members. <i>Journal of Palliative Medicine</i> , 2018, 21, 335-341.  | 1.1 | 15        |
| 39 | Efficacy of immediate-release oxycodone for dyspnoea in cancer patient: cancer dyspnoea relief (CDR) trial. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 1070-1075.   | 1.3 | 14        |
| 40 | A randomized, double-blind, non-inferiority study of hydromorphone hydrochloride immediate-release tablets versus oxycodone hydrochloride immediate-release powder for cancer pain: efficacy and safety in Japanese cancer patients. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 542-547.  | 1.3 | 7         |
| 41 | Regional Differences in Quality of Care at Palliative Care Units (PCUs), and Complicated Grief and Depression of Bereaved Family Members: Results from a Nationwide Bereavement Survey in Japan. <i>Palliative Care Research</i> , 2018, 13, 235-243.   | 0.0 | 1         |
| 42 | Medical Staff’s Support for Family Members Who Verbally Communicate Feelings to Patients in Palliative Care Units: A Survey of Bereaved Family Members. <i>Palliative Care Research</i> , 2018, 13, 263-271.  | 0.0 | 0         |
| 43 | Perspectives and attitudes toward the integration of oncology and palliative care in Japan: Qualitative analysis of a nationwide survey.. <i>Journal of Clinical Oncology</i> , 2018, 36, 96-96.  | 1.6 | 2         |
| 44 | Authors' Reply to Rady and Verheijde. <i>Journal of Pain and Symptom Management</i> , 2017, 53, e12-e13.  | 1.2 | 0         |
| 45 | Effects of End-of-Life Discussions on the Mental Health of Bereaved Family Members and Quality of Patient Death and Care. <i>Journal of Pain and Symptom Management</i> , 2017, 54, 17-26.e1.   | 1.2 | 71        |
| 46 | The relationship between cancer patients' place of death and bereaved caregivers' mental health status. <i>Psycho-Oncology</i> , 2017, 26, 1959-1964.   | 2.3 | 24        |
| 47 | Authors' Reply to Twycross. <i>Journal of Pain and Symptom Management</i> , 2017, 53, e15-e16.  | 1.2 | 0         |
| 48 | Talking About Death With Terminally-Ill Cancer Patients: What Contributes to the Regret of Bereaved Family Members?. <i>Journal of Pain and Symptom Management</i> , 2017, 54, 853-860.e1.  | 1.2 | 20        |
| 49 | Psychological and psychiatric symptoms of terminally ill patients with cancer and their family caregivers in the home-care setting: A nation-wide survey from the perspective of bereaved family members in Japan. <i>Journal of Psychosomatic Research</i> , 2017, 103, 127-132.   | 2.6 | 16        |
| 50 | Continuous Deep Sedation: A Proposal for Performing More Rigorous Empirical Research. <i>Journal of Pain and Symptom Management</i> , 2017, 53, 146-152.  | 1.2 | 44        |
| 51 | A randomized, double-blind study of hydromorphone hydrochloride extended-release tablets versus oxycodone hydrochloride extended-release tablets for cancer pain: efficacy and safety in Japanese cancer patients (EXHEAL: a Phase III study of EXtended-release HydromorphonE for cAncer pain reLief). <i>Journal of Pain Research</i> , 2017, Volume 10, 1953-1962. | 2.0 | 7         |
| 52 | Symptom Prevalence, Severity, and Distress Assessed by Memorial Symptom Assessment Scale (MSAS) in the General Population in Japan. <i>Palliative Care Research</i> , 2017, 12, 761-770.  | 0.0 | 0         |
| 53 | Exploratory Qualitative Study of Regret Stemming from Ending Terminal Treatment and Psychological Coping among the Bereaved Family Members of Cancer Patients: What Does the Family Regret about Terminal Treatment Choices and Why?. <i>Palliative Care Research</i> , 2017, 12, 753-760.  | 0.0 | 2         |
| 54 | Why Patients in Institutions with Religious Background Achieve “Good Death” ? : Findings from National Bereavement Survey of 127 Hospice and Palliative Care Units. <i>Palliative Care Research</i> , 2017, 12, 211-220.  | 0.0 | 2         |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 55 | A Second Time Nationwide Survey of Quality of End-of-Life Cancer Care in General Hospitals, Inpatient Palliative Care Units, and Clinics in Japan: The J-HOPE 2 Study. <i>Palliative Care Research</i> , 2016, 11, 254-264.                               | 0.0  | 9         |
| 56 | Family caregivers require mental health specialists for end-of-life psychosocial problems at home: a nationwide survey in Japan. <i>Psycho-Oncology</i> , 2016, 25, 641-647.  | 2.3  | 7         |
| 57 | End-of-Life Medical Treatments in the Last Two Weeks of Life in Palliative Care Units in Japan, 2005-2006: A Nationwide Retrospective Cohort Survey. <i>Journal of Palliative Medicine</i> , 2016, 19, 1188-1196.   | 1.1  | 9         |
| 58 | Uniform definition of continuous-deep sedation. <i>Lancet Oncology</i> , The, 2016, 17, e222.   | 10.7 | 11        |
| 59 | Shock Index and Decreased Level of Consciousness as Terminal Cancer Patients' Survival Time Predictors: A Retrospective Cohort Study. <i>Journal of Pain and Symptom Management</i> , 2016, 51, 220-231.e2.   | 1.2  | 8         |
| 60 | Why People Accept Opioids: Role of General Attitudes Toward Drugs, Experience as a Bereaved Family, Information From Medical Professionals, and Personal Beliefs Regarding a Good Death. <i>Journal of Pain and Symptom Management</i> , 2015, 49, 45-54. | 1.2  | 6         |
| 61 | Independent Validation of the Modified Prognosis Palliative Care Study Predictor Models in Three Palliative Care Settings. <i>Journal of Pain and Symptom Management</i> , 2015, 49, 853-860.   | 1.2  | 35        |
| 62 | Palliative Care Physicians' Attitudes Toward Patient Autonomy and a Good Death in East Asian Countries. <i>Journal of Pain and Symptom Management</i> , 2015, 50, 190-199.e1.   | 1.2  | 67        |
| 63 | Family evaluation about caring the body of deceased patient by nurses in Japanese inpatient hospices. <i>Palliative Care Research</i> , 2015, 10, 101-107.  | 0.0  | 1         |
| 64 | Definition of Sedation for Symptom Relief. <i>Journal of Pain and Symptom Management</i> , 2002, 24, 447-453.   | 1.2  | 142       |
| 65 | Proposed definitions for terminal sedation. <i>Lancet</i> , The, 2001, 358, 335-336.  | 13.7 | 80        |