

# Ling Qi

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1546945/ling-qi-publications-by-year.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

84  
papers

5,255  
citations

40  
h-index

72  
g-index

87  
ext. papers

6,392  
ext. citations

10.9  
avg, IF

5.6  
L-index

#	Paper	IF	Citations
84	Endoplasmic reticulum-associated degradation is required for nephrin maturation and kidney glomerular filtration function. <i>Journal of Clinical Investigation</i> , <b>2021</b> , 131,	15.9	3
83	Transcription- and phosphorylation-dependent control of a functional interplay between XBP1s and PINK1 governs mitophagy and potentially impacts Parkinson disease pathophysiology. <i>Autophagy</i> , <b>2021</b> , 1-23	10.2	2
82	Lipoprotein Lipase and Its Regulators: An Unfolding Story. <i>Trends in Endocrinology and Metabolism</i> , <b>2021</b> , 32, 48-61	8.8	21
81	Normal and defective pathways in biogenesis and maintenance of the insulin storage pool. <i>Journal of Clinical Investigation</i> , <b>2021</b> , 131,	15.9	10
80	ER-associated degradation preserves hematopoietic stem cell quiescence and self-renewal by restricting mTOR activity. <i>Blood</i> , <b>2020</b> , 136, 2975-2986	2.2	12
79	Endoplasmic Reticulum Protein Quality Control in $\beta$ Cells. <i>Seminars in Cell and Developmental Biology</i> , <b>2020</b> , 103, 59-67	7.5	9
78	The obesity-induced adipokine sST2 exacerbates adipose T and ILC2 depletion and promotes insulin resistance. <i>Science Advances</i> , <b>2020</b> , 6, eaay6191	14.3	25
77	Endoplasmic reticulum-associated degradation regulates mitochondrial dynamics in brown adipocytes. <i>Science</i> , <b>2020</b> , 368, 54-60	33.3	39
76	Selective EMC subunits act as molecular tethers of intracellular organelles exploited during viral entry. <i>Nature Communications</i> , <b>2020</b> , 11, 1127	17.4	8
75	Sel1L-Hrd1 ER-associated degradation maintains $\beta$ cell identity via TGF- $\beta$ signaling. <i>Journal of Clinical Investigation</i> , <b>2020</b> , 130, 3499-3510	15.9	14
74	NOTCH3 is non-enzymatically fragmented in inherited cerebral small-vessel disease. <i>Journal of Biological Chemistry</i> , <b>2020</b> , 295, 1960-1972	5.4	6
73	Protein quality control through endoplasmic reticulum-associated degradation maintains haematopoietic stem cell identity and niche interactions. <i>Nature Cell Biology</i> , <b>2020</b> , 22, 1162-1169	23.4	6
72	Toll-like receptors TLR2 and TLR4 block the replication of pancreatic $\beta$ cells in diet-induced obesity. <i>Nature Immunology</i> , <b>2019</b> , 20, 677-686	19.1	30
71	Intrinsic Structural Features of the Human IRE1 $\alpha$ Transmembrane Domain Sense Membrane Lipid Saturation. <i>Cell Reports</i> , <b>2019</b> , 27, 307-320.e5	10.6	20
70	Landscape of Intercellular Crosstalk in Healthy and NASH Liver Revealed by Single-Cell Secretome Gene Analysis. <i>Molecular Cell</i> , <b>2019</b> , 75, 644-660.e5	17.6	218
69	Medullary thymic epithelial NF- $\kappa$ B-inducing kinase (NIK)/IKK $\beta$ pathway shapes autoimmunity and liver and lung homeostasis in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2019</b> , 116, 19090-19097	11.5	15
68	ER-associated degradation in health and disease - from substrate to organism. <i>Journal of Cell Science</i> , <b>2019</b> , 132,	5.3	22

67	Requirement for translocon-associated protein (TRAP) in insulin biogenesis. <i>Science Advances</i> , <b>2019</b> , 5, eaax0292	14.3	10
66	Misfolded proinsulin in the endoplasmic reticulum during development of beta cell failure in diabetes. <i>Annals of the New York Academy of Sciences</i> , <b>2018</b> , 1418, 5-19	6.5	29
65	Designing a retrievable and scalable cell encapsulation device for potential treatment of type 1 diabetes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2018</b> , 115, E263-E272	11.5	97
64	IRE1 governs cytoskeleton remodelling and cell migration through a direct interaction with filamin A. <i>Nature Cell Biology</i> , <b>2018</b> , 20, 942-953	23.4	63
63	Hypothalamic ER-associated degradation regulates POMC maturation, feeding, and age-associated obesity. <i>Journal of Clinical Investigation</i> , <b>2018</b> , 128, 1125-1140	15.9	26
62	Hepatic Sel1L-Hrd1 ER-associated degradation (ERAD) manages FGF21 levels and systemic metabolism via CREBH. <i>EMBO Journal</i> , <b>2018</b> , 37,	13	27
61	Coordinate regulation of mutant NPC1 degradation by selective ER autophagy and MARCH6-dependent ERAD. <i>Nature Communications</i> , <b>2018</b> , 9, 3671	17.4	49
60	Quality Control in the Endoplasmic Reticulum: Crosstalk between ERAD and UPR pathways. <i>Trends in Biochemical Sciences</i> , <b>2018</b> , 43, 593-605	10.3	163
59	PERK and XBP1 differentially regulate CXCL10 and CCL2 production. <i>Experimental Eye Research</i> , <b>2017</b> , 155, 1-14	3.7	11
58	New Insights into the Physiological Role of Endoplasmic Reticulum-Associated Degradation. <i>Trends in Cell Biology</i> , <b>2017</b> , 27, 430-440	18.3	102
57	Hypoxia-Inducible Lipid Droplet-Associated Is Not a Direct Physiological Regulator of Lipolysis in Adipose Tissue. <i>Endocrinology</i> , <b>2017</b> , 158, 1231-1251	4.8	17
56	Feeding mice fat promotes foam cell formation in mesenteric lymph nodes without leading to ascites. <i>Journal of Lipid Research</i> , <b>2017</b> , 58, 1100-1113	6.3	13
55	ER-associated degradation is required for vasopressin prohormone processing and systemic water homeostasis. <i>Journal of Clinical Investigation</i> , <b>2017</b> , 127, 3897-3912	15.9	37
54	The Sel1L-Hrd1 Endoplasmic Reticulum-Associated Degradation Complex Manages a Key Checkpoint in B Cell Development. <i>Cell Reports</i> , <b>2016</b> , 16, 2630-2640	10.6	27
53	Epithelial Sel1L is required for the maintenance of intestinal homeostasis. <i>Molecular Biology of the Cell</i> , <b>2016</b> , 27, 483-90	3.5	23
52	A CRISPR-Based Screen Identifies Genes Essential for West-Nile-Virus-Induced Cell Death. <i>Cell Reports</i> , <b>2015</b> , 12, 673-83	10.6	150
51	Fish Oil-Rich Diet Promotes Hematopoiesis and Alters Hematopoietic Niche. <i>Endocrinology</i> , <b>2015</b> , 156, 2821-30	4.8	22
50	Endoplasmic reticulum quality control in cancer: Friend or foe. <i>Seminars in Cancer Biology</i> , <b>2015</b> , 33, 25-33	32.7	40

49	IRE1 $\alpha$ is an endogenous substrate of endoplasmic-reticulum-associated degradation. <i>Nature Cell Biology</i> , <b>2015</b> , 17, 1546-55	23.4	115
48	Developing robust, hydrogel-based, nanofiber-enabled encapsulation devices (NEEDs) for cell therapies. <i>Biomaterials</i> , <b>2015</b> , 37, 40-8	15.6	69
47	High-Resolution Metabolomics with Acyl-CoA Profiling Reveals Widespread Remodeling in Response to Diet. <i>Molecular and Cellular Proteomics</i> , <b>2015</b> , 14, 1489-500	7.6	68
46	Sel1L is indispensable for mammalian endoplasmic reticulum-associated degradation, endoplasmic reticulum homeostasis, and survival. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2014</b> , 111, E582-91	11.5	101
45	Diet-induced alterations in gut microflora contribute to lethal pulmonary damage in TLR2/TLR4-deficient mice. <i>Cell Reports</i> , <b>2014</b> , 8, 137-49	10.6	35
44	Chronic intake of high fish oil diet induces myeloid-derived suppressor cells to promote tumor growth. <i>Cancer Immunology, Immunotherapy</i> , <b>2014</b> , 63, 663-73	7.4	17
43	Overexpression of TRB3 in muscle alters muscle fiber type and improves exercise capacity in mice. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , <b>2014</b> , 306, R925-33	3.2	17
42	The ER-associated degradation adaptor protein Sel1L regulates LPL secretion and lipid metabolism. <i>Cell Metabolism</i> , <b>2014</b> , 20, 458-70	24.6	62
41	Tipping the balance in metabolic regulation: regulating regulatory T cells by costimulation. <i>Diabetes</i> , <b>2014</b> , 63, 1179-81	0.9	2
40	Hypoxia-inducible lipid droplet-associated (HILPDA) is a novel peroxisome proliferator-activated receptor (PPAR) target involved in hepatic triglyceride secretion. <i>Journal of Biological Chemistry</i> , <b>2014</b> , 289, 19279-93	5.4	42
39	Adipocyte spliced form of X-box-binding protein 1 promotes adiponectin multimerization and systemic glucose homeostasis. <i>Diabetes</i> , <b>2014</b> , 63, 867-79	0.9	24
38	Phenformin activates the unfolded protein response in an AMP-activated protein kinase (AMPK)-dependent manner. <i>Journal of Biological Chemistry</i> , <b>2013</b> , 288, 13631-8	5.4	19
37	ER-stress-associated functional link between Parkin and DJ-1 via a transcriptional cascade involving the tumor suppressor p53 and the spliced X-box binding protein XBP-1. <i>Journal of Cell Science</i> , <b>2013</b> , 126, 2124-33	5.3	57
36	Direct control of hepatic glucose production by interleukin-13 in mice. <i>Journal of Clinical Investigation</i> , <b>2013</b> , 123, 261-71	15.9	92
35	The transcriptional co-regulator HCF-1 is required for INS-1 $\beta$ cell glucose-stimulated insulin secretion. <i>PLoS ONE</i> , <b>2013</b> , 8, e78841	3.7	3
34	A Conserved Structural Determinant Located at the Interdomain Region of Mammalian IRE1 $\alpha$ . <i>FASEB Journal</i> , <b>2013</b> , 27, 794.18	0.9	
33	Nonmuscle myosin IIB links cytoskeleton to IRE1 $\beta$ signaling during ER stress. <i>Developmental Cell</i> , <b>2012</b> , 23, 1141-52	10.2	45
32	The full capacity of AICAR to reduce obesity-induced inflammation and insulin resistance requires myeloid SIRT1. <i>PLoS ONE</i> , <b>2012</b> , 7, e49935	3.7	40

31	Mechanisms of inflammatory responses in obese adipose tissue. <i>Annual Review of Nutrition</i> , <b>2012</b> , 32, 261-86	9.9	207
30	The ATP-P2X7 signaling axis is dispensable for obesity-associated inflammasome activation in adipose tissue. <i>Diabetes</i> , <b>2012</b> , 61, 1471-8	0.9	58
29	Activation of natural killer T cells promotes M2 Macrophage polarization in adipose tissue and improves systemic glucose tolerance via interleukin-4 (IL-4)/STAT6 protein signaling axis in obesity. <i>Journal of Biological Chemistry</i> , <b>2012</b> , 287, 13561-71	5.4	155
28	Short term high fat diet challenge promotes alternative macrophage polarization in adipose tissue via natural killer T cells and interleukin-4. <i>Journal of Biological Chemistry</i> , <b>2012</b> , 287, 24378-86	5.4	107
27	Gr-1+ CD11b+ myeloid-derived suppressor cells suppress inflammation and promote insulin sensitivity in obesity. <i>Journal of Biological Chemistry</i> , <b>2011</b> , 286, 23591-9	5.4	111
26	Stressed out about obesity: IRE1/XBP1 in metabolic disorders. <i>Trends in Endocrinology and Metabolism</i> , <b>2011</b> , 22, 374-81	8.8	63
25	Detecting and quantitating physiological endoplasmic reticulum stress. <i>Methods in Enzymology</i> , <b>2011</b> , 490, 137-46	1.7	29
24	A conserved structural determinant located at the interdomain region of mammalian inositol-requiring enzyme 1alpha. <i>Journal of Biological Chemistry</i> , <b>2011</b> , 286, 30859-30866	5.4	34
23	Haploid insufficiency of suppressor enhancer Lin12 1-like (SEL1L) protein predisposes mice to high fat diet-induced hyperglycemia. <i>Journal of Biological Chemistry</i> , <b>2011</b> , 286, 22275-82	5.4	9
22	PKA phosphorylation couples hepatic inositol-requiring enzyme 1alpha to glucagon signaling in glucose metabolism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 15852-7	11.5	64
21	Deficiency of suppressor enhancer Lin12 1 like (SEL1L) in mice leads to systemic endoplasmic reticulum stress and embryonic lethality. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 13694-703	5.4	57
20	XBP-1 couples endoplasmic reticulum stress to augmented IFN-beta induction via a cis-acting enhancer in macrophages. <i>Journal of Immunology</i> , <b>2010</b> , 185, 2324-30	5.3	91
19	The roles of ATF3, an adaptive-response gene, in high-fat-diet-induced diabetes and pancreatic beta-cell dysfunction. <i>Molecular Endocrinology</i> , <b>2010</b> , 24, 1423-33		58
18	Angptl4 protects against severe proinflammatory effects of saturated fat by inhibiting fatty acid uptake into mesenteric lymph node macrophages. <i>Cell Metabolism</i> , <b>2010</b> , 12, 580-92	24.6	178
17	Emerging roles for XBP1, a sUPeR transcription factor. <i>Gene Expression</i> , <b>2010</b> , 15, 13-25	3.4	78
16	SUMO modification regulates the transcriptional activity of XBP1. <i>Biochemical Journal</i> , <b>2010</b> , 429, 95-102	3.8	55
15	The pseudokinase tribbles homolog 3 interacts with ATF4 to negatively regulate insulin exocytosis in human and mouse beta cells. <i>Journal of Clinical Investigation</i> , <b>2010</b> , 120, 2876-88	15.9	92
14	A Phos-tag-based approach reveals the extent of physiological endoplasmic reticulum stress. <i>PLoS ONE</i> , <b>2010</b> , 5, e11621	3.7	52

13	Adipocyte CREB promotes insulin resistance in obesity. <i>Cell Metabolism</i> , <b>2009</b> , 9, 277-86	24.6	139
12	The IRE1alpha-XBP1 pathway of the unfolded protein response is required for adipogenesis. <i>Cell Metabolism</i> , <b>2009</b> , 9, 556-64	24.6	203
11	TRB3 links the E3 ubiquitin ligase COP1 to lipid metabolism. <i>Science</i> , <b>2006</b> , 312, 1763-6	33.3	240
10	Telomere fusion to chromosome breaks reduces oncogenic translocations and tumour formation. <i>Nature Cell Biology</i> , <b>2005</b> , 7, 706-11	23.4	23
9	The CREB coactivator TORC2 is a key regulator of fasting glucose metabolism. <i>Nature</i> , <b>2005</b> , 437, 1109-14	10.4	783
8	Invariant chain and the MHC class II cytoplasmic domains regulate localization of MHC class II molecules to lipid rafts in tumor cell-based vaccines. <i>Journal of Immunology</i> , <b>2004</b> , 172, 907-14	5.3	7
7	Short telomeres and ataxia-telangiectasia mutated deficiency cooperatively increase telomere dysfunction and suppress tumorigenesis. <i>Cancer Research</i> , <b>2003</b> , 63, 8188-96	10.1	52
6	Immunologic Targets for the Gene Therapy of Cancer <b>2002</b> , 127-142		
5	H2-O inhibits presentation of bacterial superantigens, but not endogenous self antigens. <i>Journal of Immunology</i> , <b>2001</b> , 167, 1371-8	5.3	14
4	MHC class II presentation of endogenous tumor antigen by cellular vaccines depends on the endocytic pathway but not H2-M. <i>Traffic</i> , <b>2000</b> , 1, 152-60	5.7	21
3	Tumor cells present MHC class II-restricted nuclear and mitochondrial antigens and are the predominant antigen presenting cells in vivo. <i>Journal of Immunology</i> , <b>2000</b> , 165, 5451-61	5.3	58
2	Cell-based vaccines for the stimulation of immunity to metastatic cancers. <i>Immunological Reviews</i> , <b>1999</b> , 170, 101-14	11.3	42
1	Endoplasmic reticulum associated degradation preserves hematopoietic stem cell quiescence and self-renewal by restricting mTOR activity		1