

Peter Angel

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1549573/peter-angel-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

108
papers

11,446
citations

47
h-index

106
g-index

111
ext. papers

12,405
ext. citations

7.8
avg, IF

5.59
L-index

#	Paper	IF	Citations
108	Podoplanin is required for tumor cell invasion in cutaneous squamous cell carcinoma. <i>Experimental Dermatology</i> , 2021 , 30, 1619-1630	4	0
107	A Set of Cell Lines Derived from a Genetic Murine Glioblastoma Model Recapitulates Molecular and Morphological Characteristics of Human Tumors. <i>Cancers</i> , 2021 , 13,	6.6	1
106	JUNB suppresses distant metastasis by influencing the initial metastatic stage. <i>Clinical and Experimental Metastasis</i> , 2021 , 38, 411-423	4.7	1
105	Glioblastoma epigenome profiling identifies SOX10 as a master regulator of molecular tumour subtype. <i>Nature Communications</i> , 2020 , 11, 6434	17.4	7
104	Modeling glioblastoma invasion using human brain organoids and single-cell transcriptomics. <i>Neuro-Oncology</i> , 2020 , 22, 1138-1149	1	37
103	Regulatory T cells sense effector T-cell activation through synchronized JunB expression. <i>FEBS Letters</i> , 2019 , 593, 1020-1029	3.8	6
102	Intratumoral platelet aggregate formation in a murine preclinical glioma model depends on podoplanin expression on tumor cells. <i>Blood Advances</i> , 2019 , 3, 1092-1102	7.8	15
101	Podoplanin Positive Myeloid Cells Promote Glioma Development by Immune Suppression. <i>Frontiers in Oncology</i> , 2019 , 9, 187	5.3	6
100	GPD1 Specifically Marks Dormant Glioma Stem Cells with a Distinct Metabolic Profile. <i>Cell Stem Cell</i> , 2019 , 25, 241-257.e8	18	33
99	Podoplanin expression is a prognostic biomarker but may be dispensable for the malignancy of glioblastoma. <i>Neuro-Oncology</i> , 2019 , 21, 326-336	1	10
98	An advanced glioma cell invasion assay based on organotypic brain slice cultures. <i>BMC Cancer</i> , 2018 , 18, 103	4.8	34
97	Homeostatic nuclear RAGE-ATM interaction is essential for efficient DNA repair. <i>Nucleic Acids Research</i> , 2017 , 45, 10595-10613	20.1	37
96	TGF- β 1 and TGF- β 2 abundance in liver diseases of mice and men. <i>Oncotarget</i> , 2016 , 7, 19499-518	3.3	40
95	High S100A8 and S100A12 protein expression is a favorable prognostic factor for survival of oropharyngeal squamous cell carcinoma. <i>International Journal of Cancer</i> , 2015 , 136, 2037-46	7.5	28
94	A pro-tumorigenic function of S100A8/A9 in carcinogen-induced hepatocellular carcinoma. <i>Cancer Letters</i> , 2015 , 369, 396-404	9.9	19
93	Chronic liver inflammation and hepatocellular carcinogenesis are independent of S100A9. <i>International Journal of Cancer</i> , 2015 , 136, 2458-63	7.5	4
92	Epithelial deletion of podoplanin is dispensable for re-epithelialization of skin wounds. <i>Experimental Dermatology</i> , 2015 , 24, 785-7	4	8

91	Junb controls lymphatic vascular development in zebrafish via miR-182. <i>Scientific Reports</i> , 2015 , 5, 15007	4.9	17
90	Neutralization of the CD95 ligand by APG101 inhibits invasion of glioma cells in vitro. <i>Anti-Cancer Drugs</i> , 2015 , 26, 716-27	2.4	19
89	Human and mouse VEGFA-amplified hepatocellular carcinomas are highly sensitive to sorafenib treatment. <i>Cancer Discovery</i> , 2014 , 4, 730-43	24.4	137
88	Effects of selective MMP-13 inhibition in squamous cell carcinoma depend on estrogen. <i>International Journal of Cancer</i> , 2014 , 135, 2749-59	7.5	3
87	Efficient keratinocyte differentiation strictly depends on JNK-induced soluble factors in fibroblasts. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 1332-1341	4.3	28
86	Loss of stromal JUNB does not affect tumor growth and angiogenesis. <i>International Journal of Cancer</i> , 2014 , 134, 1511-6	7.5	7
85	Keratinocyte-specific deletion of the receptor RAGE modulates the kinetics of skin inflammation in vivo. <i>Journal of Investigative Dermatology</i> , 2013 , 133, 2400-2406	4.3	22
84	Inflammation-mediated skin tumorigenesis induced by epidermal c-Fos. <i>Genes and Development</i> , 2013 , 27, 1959-73	12.6	43
83	Collagenase-3 (MMP-13) deficiency protects C57BL/6 mice from antibody-induced arthritis. <i>Arthritis Research and Therapy</i> , 2013 , 15, R222	5.7	26
82	Receptor for advanced glycation endproducts (RAGE) is a key regulator of oval cell activation and inflammation-associated liver carcinogenesis in mice. <i>Hepatology</i> , 2013 , 58, 363-73	11.2	66
81	Procollagen I-expressing renin cell precursors. <i>American Journal of Physiology - Renal Physiology</i> , 2013 , 305, F355-61	4.3	7
80	Stathmin regulates keratinocyte proliferation and migration during cutaneous regeneration. <i>PLoS ONE</i> , 2013 , 8, e75075	3.7	12
79	The receptor for advanced glycation end products is dispensable in a mouse model of oral and esophageal carcinogenesis. <i>Histology and Histopathology</i> , 2013 , 28, 1585-94	1.4	7
78	Opposing function of MYBBP1A in proliferation and migration of head and neck squamous cell carcinoma cells. <i>BMC Cancer</i> , 2012 , 12, 72	4.8	15
77	Hepatocyte-specific S100a8 and S100a9 transgene expression in mice causes Cxcl1 induction and systemic neutrophil enrichment. <i>Cell Communication and Signaling</i> , 2012 , 10, 40	7.5	17
76	Expression of podoplanin in human astrocytic brain tumors is controlled by the PI3K-AKT-AP-1 signaling pathway and promoter methylation. <i>Neuro-Oncology</i> , 2012 , 14, 426-39	1	44
75	KIAA1797/FOCAD encodes a novel focal adhesion protein with tumour suppressor function in gliomas. <i>Brain</i> , 2012 , 135, 1027-41	11.2	37
74	Enhanced StefinA and Sprr2 expression during papilloma formation in HPV8 transgenic mice. <i>Journal of Dermatological Science</i> , 2011 , 62, 84-90	4.3	12

73	Accelerated aging phenotype in mice with conditional deficiency for mitochondrial superoxide dismutase in the connective tissue. <i>Aging Cell</i> , 2011 , 10, 239-54	9.9	73
72	Expression and function of the kallikrein-related peptidase 6 in the human melanoma microenvironment. <i>Journal of Investigative Dermatology</i> , 2011 , 131, 2281-8	4.3	36
71	The Transcription Factor AP-1 in Squamous Cell Carcinogenesis: Lessons from Mouse Models of Skin Carcinogenesis 2011 , 185-199		1
70	Control of hair follicle cell fate by underlying mesenchyme through a CSL-Wnt5a-FoxN1 regulatory axis. <i>Genes and Development</i> , 2010 , 24, 1519-32	12.6	75
69	Impaired skin regeneration and remodeling after cutaneous injury and chemically induced hyperplasia in taps-transgenic mice. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 1922-30	4.3	14
68	MMP13 as a stromal mediator in controlling persistent angiogenesis in skin carcinoma. <i>Carcinogenesis</i> , 2010 , 31, 1175-84	4.6	101
67	Identification of the Rage-dependent gene regulatory network in a mouse model of skin inflammation. <i>BMC Genomics</i> , 2010 , 11, 537	4.5	17
66	Loss of matrix metalloproteinase-13 attenuates murine radiation-induced pulmonary fibrosis. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 77, 582-90	4	36
65	Junb regulates arterial contraction capacity, cellular contractility, and motility via its target Myl9 in mice. <i>Journal of Clinical Investigation</i> , 2010 , 120, 2307-18	15.9	26
64	Overexpression of far upstream element binding proteins: a mechanism regulating proliferation and migration in liver cancer cells. <i>Hepatology</i> , 2009 , 50, 1130-9	11.2	79
63	S100A8 and S100A9 are novel nuclear factor kappa B target genes during malignant progression of murine and human liver carcinogenesis. <i>Hepatology</i> , 2009 , 50, 1251-62	11.2	108
62	The receptor RAGE: Bridging inflammation and cancer. <i>Cell Communication and Signaling</i> , 2009 , 7, 12	7.5	143
61	AP-1-controlled hepatocyte growth factor activation promotes keratinocyte migration via CEACAM1 and urokinase plasminogen activator/urokinase plasminogen receptor. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 1140-8	4.3	16
60	Stromal expression of MMP-13 is required for melanoma invasion and metastasis. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 2686-93	4.3	86
59	The transcription factor Fos: a Janus-type regulator in health and disease. <i>Histology and Histopathology</i> , 2009 , 24, 1451-61	1.4	85
58	Dual Role of S100A8 and S100A9 in Inflammation-Associated Cancer. <i>Anti-Inflammatory and Anti-Allergy Agents in Medicinal Chemistry</i> , 2009 , 8, 329-336	2	4
57	Gene network dynamics controlling keratinocyte migration. <i>Molecular Systems Biology</i> , 2008 , 4, 199	12.2	44
56	Podoplanin is a novel fos target gene in skin carcinogenesis. <i>Cancer Research</i> , 2008 , 68, 6877-83	10.1	60

55	RAGE signaling sustains inflammation and promotes tumor development. <i>Journal of Experimental Medicine</i> , 2008 , 205, 275-85	16.6	306
54	CEBPbeta, JunD and c-Jun contribute to the transcriptional activation of the metastasis-associated C4.4A gene. <i>International Journal of Cancer</i> , 2007 , 120, 2135-47	7.5	12
53	Critical role for NF-kappaB-induced JunB in VEGF regulation and tumor angiogenesis. <i>EMBO Journal</i> , 2007 , 26, 710-9	13	102
52	Conditional deletion of insulin-like growth factor-I in collagen type 1alpha2-expressing cells results in postnatal lethality and a dramatic reduction in bone accretion. <i>Endocrinology</i> , 2007 , 148, 5706-15	4.8	81
51	Kallikrein 6 induces E-cadherin shedding and promotes cell proliferation, migration, and invasion. <i>Cancer Research</i> , 2007 , 67, 8198-206	10.1	108
50	JunB is required for IgE-mediated degranulation and cytokine release of mast cells. <i>Journal of Immunology</i> , 2007 , 179, 6873-80	5.3	16
49	S100A8 and S100A9 in inflammation and cancer. <i>Biochemical Pharmacology</i> , 2006 , 72, 1622-31	6	505
48	p44 mitogen-activated protein kinase (extracellular signal-regulated kinase 1)-dependent signaling contributes to epithelial skin carcinogenesis. <i>Cancer Research</i> , 2006 , 66, 2700-7	10.1	69
47	Cutting edge: the AP-1 subunit JunB determines NK cell-mediated target cell killing by regulation of the NKG2D-ligand RAE-1epsilon. <i>Journal of Immunology</i> , 2006 , 176, 7-11	5.3	39
46	JunB is required for endothelial cell morphogenesis by regulating core-binding factor beta. <i>Journal of Cell Biology</i> , 2006 , 175, 981-91	7.3	32
45	A novel aspartic proteinase-like gene expressed in stratified epithelia and squamous cell carcinoma of the skin. <i>American Journal of Pathology</i> , 2006 , 168, 1354-64	5.8	15
44	Epidermal development and wound healing in matrix metalloproteinase 13-deficient mice. <i>Journal of Investigative Dermatology</i> , 2006 , 126, 486-96	4.3	72
43	Delayed wound healing and epidermal hyperproliferation in mice lacking JunB in the skin. <i>Journal of Investigative Dermatology</i> , 2006 , 126, 902-11	4.3	53
42	c-Jun and JunB are essential for hypoglycemia-mediated VEGF induction. <i>Annals of the New York Academy of Sciences</i> , 2006 , 1091, 310-8	6.5	20
41	Up-regulation of insulin-like growth factor axis components in human primary prostate cancer correlates with tumor grade. <i>Human Pathology</i> , 2005 , 36, 1186-96	3.7	87
40	c-Fos-dependent induction of the small ras-related GTPase Rab11a in skin carcinogenesis. <i>American Journal of Pathology</i> , 2005 , 167, 243-53	5.8	39
39	Psoriasis-like skin disease and arthritis caused by inducible epidermal deletion of Jun proteins. <i>Nature</i> , 2005 , 437, 369-75	50.4	463
38	High invasive melanoma cells induce matrix metalloproteinase-1 synthesis in fibroblasts by interleukin-1alpha and basic fibroblast growth factor-mediated mechanisms. <i>Journal of Investigative Dermatology</i> , 2005 , 124, 638-43	4.3	56

37	Two-hybrid fluorescence cross-correlation spectroscopy detects protein-protein interactions in vivo. <i>ChemPhysChem</i> , 2005 , 6, 984-90	3.2	79
36	Preeclampsia: increased expression of soluble ADAM 12. <i>Journal of Molecular Medicine</i> , 2005 , 83, 887-965.5		41
35	Calcium-binding proteins S100A8 and S100A9 as novel diagnostic markers in human prostate cancer. <i>Clinical Cancer Research</i> , 2005 , 11, 5146-52	12.9	202
34	Increased keratinocyte proliferation by JUN-dependent expression of PTN and SDF-1 in fibroblasts. <i>Journal of Cell Science</i> , 2005 , 118, 1981-9	5.3	84
33	Hyaluronan-oligosaccharide-induced transcription of metalloproteases. <i>Journal of Cell Science</i> , 2004 , 117, 359-67	5.3	132
32	Mice lacking JunB are osteopenic due to cell-autonomous osteoblast and osteoclast defects. <i>Journal of Cell Biology</i> , 2004 , 164, 613-23	7.3	166
31	Identification of novel AP-1 target genes in fibroblasts regulated during cutaneous wound healing. <i>Oncogene</i> , 2004 , 23, 7005-17	9.2	50
30	Cre recombinase-mediated gene targeting of mesenchymal cells. <i>Genesis</i> , 2004 , 38, 139-44	1.9	40
29	AP-1 subunits: quarrel and harmony among siblings. <i>Journal of Cell Science</i> , 2004 , 117, 5965-73	5.3	923
28	Altered endochondral bone development in matrix metalloproteinase 13-deficient mice. <i>Development (Cambridge)</i> , 2004 , 131, 5883-95	6.6	450
27	Early activation and induction of apoptosis in T cells is independent of c-Fos. <i>Annals of the New York Academy of Sciences</i> , 2003 , 1010, 225-31	6.5	6
26	Profile of gene expression induced by the tumour promoter TPA in murine epithelial cells. <i>International Journal of Cancer</i> , 2003 , 104, 699-708	7.5	51
25	Increase of AKT/PKB expression correlates with gleason pattern in human prostate cancer. <i>International Journal of Cancer</i> , 2003 , 107, 676-80	7.5	103
24	An unexpected role for FosB in activation-induced cell death of T cells. <i>Oncogene</i> , 2003 , 22, 1333-9	9.2	28
23	TAF7 (TAFII55) plays a role in the transcription activation by c-Jun. <i>Journal of Biological Chemistry</i> , 2003 , 278, 21510-6	5.4	20
22	Defective endochondral ossification in mice with strongly compromised expression of JunB. <i>Journal of Cell Science</i> , 2003 , 116, 4587-96	5.3	34
21	Function of AP-1 target genes in mesenchymal-epithelial cross-talk in skin. <i>Biochemical Pharmacology</i> , 2002 , 64, 949-56	6	65
20	Calgranulins S100A8 and S100A9 are negatively regulated by glucocorticoids in a c-Fos-dependent manner and overexpressed throughout skin carcinogenesis. <i>Oncogene</i> , 2002 , 21, 4266-76	9.2	106

19	Th2 cell-specific cytokine expression and allergen-induced airway inflammation depend on JunB. <i>EMBO Journal</i> , 2002 , 21, 6321-9	13	109
18	Cell cycle promoting activity of JunB through cyclin A activation. <i>Journal of Biological Chemistry</i> , 2002 , 277, 35961-8	5.4	69
17	Parathyroid hormone inhibits c-Jun N-terminal kinase activity in rat osteoblastic cells by a protein kinase A-dependent pathway. <i>Endocrinology</i> , 2002 , 143, 1880-8	4.8	23
16	Organotypic cocultures with genetically modified mouse fibroblasts as a tool to dissect molecular mechanisms regulating keratinocyte growth and differentiation. <i>Journal of Investigative Dermatology</i> , 2001 , 116, 816-20	4.3	97
15	Keratinocyte-specific onset of serine protease BSSP expression in experimental carcinogenesis. <i>Journal of Investigative Dermatology</i> , 2001 , 117, 634-40	4.3	22
14	Induction of the AP-1 members c-Jun and JunB by TGF-beta/Smad suppresses early Smad-driven gene activation. <i>Oncogene</i> , 2001 , 20, 2205-11	9.2	85
13	Function and regulation of AP-1 subunits in skin physiology and pathology. <i>Oncogene</i> , 2001 , 20, 2413-23	9.2	343
12	Expression of collagenase-3 (MMP-13) in c-fos-induced osteosarcomas and chondrosarcomas is restricted to a subset of cells of the osteo-/chondrogenic lineage. <i>Differentiation</i> , 2001 , 69, 49-57	3.5	9
11	AP-1 and Cbfa/runt physically interact and regulate parathyroid hormone-dependent MMP13 expression in osteoblasts through a new osteoblast-specific element 2/AP-1 composite element. <i>Journal of Biological Chemistry</i> , 2001 , 276, 20029-38	5.4	151
10	Expression of human collagenase I (MMP-1) and TIMP-1 in a baculovirus-based expression system. <i>Methods in Molecular Biology</i> , 2001 , 151, 207-18	1.4	
9	The collagen receptor DDR2 regulates proliferation and its elimination leads to dwarfism. <i>EMBO Reports</i> , 2001 , 2, 446-52	6.5	209
8	p53 and c-Jun functionally synergize in the regulation of the DNA repair gene hMSH2 in response to UV. <i>Journal of Biological Chemistry</i> , 2000 , 275, 37469-73	5.4	70
7	A novel AP-1 element in the CD95 ligand promoter is required for induction of apoptosis in hepatocellular carcinoma cells upon treatment with anticancer drugs. <i>Molecular and Cellular Biology</i> , 2000 , 20, 7826-37	4.8	114
6	c-Jun and JunB antagonistically control cytokine-regulated mesenchymal-epidermal interaction in skin. <i>Cell</i> , 2000 , 103, 745-55	56.2	347
5	The DNA binding-independent function of the glucocorticoid receptor mediates repression of AP-1-dependent genes in skin. <i>Journal of Cell Biology</i> , 1999 , 147, 1365-70	7.3	170
4	JunB is essential for mammalian placentation. <i>EMBO Journal</i> , 1999 , 18, 934-48	13	206
3	Phorbol ester-inducible genes contain a common cis element recognized by a TPA-modulated trans-acting factor. <i>Cell</i> , 1987 , 49, 729-39	56.2	2948
2	A novel neural stem cell-derived immunocompetent mouse model of glioblastoma for preclinical studies		2

1 Modeling glioblastoma invasion using human brain organoids and single-cell transcriptomics

1