

# Sharada Mokkapati

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/155711/publications.pdf>

Version: 2024-02-01

16  
papers

656  
citations

840776

11  
h-index

940533

16  
g-index

16  
all docs

16  
docs citations

16  
times ranked

1192  
citing authors

#	ARTICLE	IF	CITATIONS
1	Compound Genetic Ablation of Nidogen 1 and 2 Causes Basement Membrane Defects and Perinatal Lethality in Mice. <i>Molecular and Cellular Biology</i> , 2005, 25, 6846-6856.	2.3	201
2	Nidogensâ€™ Extracellular matrix linker molecules. <i>Microscopy Research and Technique</i> , 2008, 71, 387-395.	2.2	99
3	Î²-Catenin Activation in a Novel Liver Progenitor Cell Type Is Sufficient to Cause Hepatocellular Carcinoma and Hepatoblastoma. <i>Cancer Research</i> , 2014, 74, 4515-4525.	0.9	82
4	Epidermal Transglutaminase (TGase 3) Is Required for Proper Hair Development, but Not the Formation of the Epidermal Barrier. <i>PLoS ONE</i> , 2012, 7, e34252.	2.5	46
5	Basement Membranes in Skin Are Differently Affected by Lack of Nidogen 1 and 2. <i>Journal of Investigative Dermatology</i> , 2008, 128, 2259-2267.	0.7	45
6	Lack of Nidogen-1 and -2 Prevents Basement Membrane Assembly in Skin-Organotypic Coculture. <i>Journal of Investigative Dermatology</i> , 2007, 127, 545-554.	0.7	41
7	Pim kinase isoforms: devils defending cancer cells from therapeutic and immune attacks. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2016, 21, 1203-1213.	4.9	31
8	Nephron Progenitor But Not Stromal Progenitor Cells Give Rise to Wilms Tumors in Mouse Models with Î²-Catenin Activation or Wt1 Ablation and Igf2 Upregulation. <i>Neoplasia</i> , 2016, 18, 71-81.	5.3	26
9	Absence of the Basement Membrane Component Nidogen 2, but Not of Nidogen 1, Results in Increased Lung Metastasis in Mice. <i>Journal of Histochemistry and Cytochemistry</i> , 2012, 60, 280-289.	2.5	22
10	The development of interferon-based gene therapy for BCG unresponsive bladder cancer: from bench to bedside. <i>World Journal of Urology</i> , 2019, 37, 2041-2049.	2.2	21
11	Basement Membrane Deposition of Nidogen 1 but Not Nidogen 2 Requires the Nidogen Binding Module of the Laminin Î³1 Chain. <i>Journal of Biological Chemistry</i> , 2011, 286, 1911-1918.	3.4	14
12	Inhibition of urothelial carcinoma through targeted type I interferon-mediated immune activation. <i>Oncolmmunology</i> , 2019, 8, e1577125.	4.6	10
13	Antiadenovirus Antibodies Predict Response Durability to Nadofaragene Firadenovec Therapy in BCG-unresponsive Nonâ€™ muscle-invasive Bladder Cancer: Secondary Analysis of a Phase 3 Clinical Trial. <i>European Urology</i> , 2022, 81, 223-228.	1.9	8
14	TCF21 Promotes Luminal-Like Differentiation and Suppresses Metastasis in Bladder Cancer. <i>Molecular Cancer Research</i> , 2020, 18, 811-821.	3.4	4
15	Effects of thiazolidinedione in patients with active bladder cancer. <i>BJU International</i> , 2018, 121, 244-251.	2.5	3
16	Lentiviral interferon: A novel method for gene therapy in bladder cancer. <i>Molecular Therapy - Oncolytics</i> , 2022, 26, 141-157.	4.4	3