MarÃ-a Jose LÃ³pez-Armada

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/155743/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Mitochondrial Dysfunction and Oxidative Stress in Rheumatoid Arthritis. Antioxidants, 2022, 11, 1151.	2.2	22
2	Autophagy Activation by Resveratrol Reduces Severity of Experimental Rheumatoid Arthritis. Molecular Nutrition and Food Research, 2021, 65, e2000377.	1.5	13
3	Mitochondrial Dysfunction Plays a Relevant Role in Pathophysiology of Peritoneal Membrane Damage Induced by Peritoneal Dialysis. Antioxidants, 2021, 10, 447.	2.2	7
4	Role of mitochondrial dysfunction on rheumatic diseases. Biochemical Pharmacology, 2019, 165, 181-195.	2.0	30
5	The mitochondrial inhibitor oligomycin induces an inflammatory response in the rat knee joint. BMC Musculoskeletal Disorders, 2017, 18, 254.	0.8	21
6	Resveratrol lowers synovial hyperplasia, inflammatory markers and oxidative damage in an acute antigen-induced arthritis model. Rheumatology, 2016, 55, 1889-1900.	0.9	45
7	A Pathogenetic Role for Endothelin-1 in Peritoneal Dialysis-Associated Fibrosis. Journal of the American Society of Nephrology: JASN, 2015, 26, 173-182.	3.0	31
8	Mitochondrial dysfunction promotes and aggravates the inflammatory response in normal human synoviocytes. Rheumatology, 2014, 53, 1332-1343.	0.9	61
9	Mitochondrial dysfunction and the inflammatory response. Mitochondrion, 2013, 13, 106-118.	1.6	372
10	Nitric Oxide and the Respiratory Chain in Synovial Cells and Chondrocytes. , 2013, , 49-66.		1
11	Mitochondrial dysfunction increases inflammatory responsiveness to cytokines in normal human chondrocytes. Arthritis and Rheumatism, 2012, 64, 2927-2936.	6.7	130
12	Effect of nitric oxide on mitochondrial activity of human synovial cells. BMC Musculoskeletal Disorders, 2011, 12, 42.	0.8	50
13	Clinical significance of high levels of soluble tumour necrosis factor-Â receptor-2 produced by alternative splicing in rheumatoid arthritis: a longitudinal prospective cohort study. Rheumatology, 2011, 50, 721-728.	0.9	17
14	Proteomic analysis by twoâ€dimensional electrophoresis to identify the normal human chondrocyte proteome stimulated by tumor necrosis factor α and interleukinâ€1β. Arthritis and Rheumatism, 2010, 62, 802-814.	6.7	31
15	Proteomic analysis of human osteoarthritic chondrocytes reveals protein changes in stress and glycolysis. Proteomics, 2008, 8, 495-507.	1.3	108
16	Mitochondrial dysfunction activates cyclooxygenase 2 expression in cultured normal human chondrocytes. Arthritis and Rheumatism, 2008, 58, 2409-2419.	6.7	86
17	Differential effects of tumor necrosis factor-α and interleukin-1β on cell death in human articular chondrocytes. Osteoarthritis and Cartilage, 2008, 16, 715-722.	0.6	78
18	Anti-apoptotic effect of transforming growth factor-β1 on human articular chondrocytes: role of protein phosphatase 2A. Osteoarthritis and Cartilage, 2008, 16, 1370-1378.	0.6	29

#	Article	IF	CITATIONS
19	Decreased metalloproteinase production as a response to mechanical pressure in human cartilage: a mechanism for homeostatic regulation. Arthritis Research and Therapy, 2006, 8, R149.	1.6	36
20	Mitochondrial proteomic characterization of human normal articular chondrocytes. Osteoarthritis and Cartilage, 2006, 14, 507-518.	0.6	54
21	Cytokines, tumor necrosis factor-α and interleukin-1β, differentially regulate apoptosis in osteoarthritis cultured human chondrocytes. Osteoarthritis and Cartilage, 2006, 14, 660-669.	0.6	163
22	Mitochondrial activity is modulated by TNFÎ \pm and IL-1β in normal human chondrocyte cells. Osteoarthritis and Cartilage, 2006, 14, 1011-1022.	0.6	121
23	Proteomic characterization of human normal articular chondrocytes: A novel tool for the study of osteoarthritis and other rheumatic diseases. Proteomics, 2005, 5, 3048-3059.	1.3	106
24	Phosphatase-1 and -2A inhibition modulates apoptosis in human osteoarthritis chondrocytes independently of nitric oxide production. Annals of the Rheumatic Diseases, 2005, 64, 1079-1082.	0.5	11
25	Effect of nitric oxide on mitochondrial respiratory activity of human articular chondrocytes. Annals of the Rheumatic Diseases, 2004, 64, 388-395.	0.5	122
26	EP2/EP4 signalling inhibits monocyte chemoattractant protein-1 production induced by interleukin 1Â in synovial fibroblasts. Annals of the Rheumatic Diseases, 2004, 63, 1197-1204.	0.5	40
27	Xeno-implantation of pig chondrocytes into rabbit to treat localized articular cartilage defects: an animal model. Wound Repair and Regeneration, 2004, 12, 337-345.	1.5	41
28	Pig chondrocyte xenoimplants for human chondral defect repair: an in vitro model. Wound Repair and Regeneration, 2004, 12, 444-452.	1.5	22
29	Mitochondrial dysfunction in osteoarthritis. Mitochondrion, 2004, 4, 715-728.	1.6	153
30	Mitochondrial respiratory activity is altered in osteoarthritic human articular chondrocytes. Arthritis and Rheumatism, 2003, 48, 700-708.	6.7	195
31	Fibrin generated in the synovial fluid activates intimal cells from their apical surface: a sequential morphological study in antigen-induced arthritis. British Journal of Rheumatology, 2003, 42, 19-25.	2.5	28
32	Modulation of cell recruitment by anti-inflammatory agents in antigen-induced arthritis. Annals of the Rheumatic Diseases, 2002, 61, 1027-1030.	0.5	32
33	Cyclosporin A prevents the histologic damage of antigen arthritis without inducing fibrosis. Arthritis and Rheumatism, 2000, 43, 311.	6.7	13
34	The 80-kD fibronectin fragment increases the production of fibronectin and tumour necrosis factor-alpha (TNF-α) in cultured mesangial cells. Clinical and Experimental Immunology, 1997, 107, 398-403.	1.1	17
35	Antifibroproliferative effect of tenidap in chronic antigen-induced arthritis. Arthritis and Rheumatism, 1997, 40, 2147-2156.	6.7	8
36	Anti-Fas antibodies induce cytolysis and apoptosis in cultured human mesangial cells. Kidney International, 1996, 49, 1064-1070.	2.6	55

#	Article	IF	CITATIONS
37	Glomerular up-regulation of EIIIA and V120 fibronectin isoforms in proliferative immune complex nephritis. Kidney International, 1996, 50, 908-919.	2.6	13
38	An orally active ETA/ETB receptor antagonist ameliorates proteinuria and glomerular lesions in rats with proliferative nephritis. Kidney International, 1996, 50, 962-972.	2.6	67
39	Effects and Interactions of Endothelin-1 and Angiotensin II on Matrix Protein Expression and Synthesis and Mesangial Cell Growth. Hypertension, 1996, 27, 885-892.	1.3	138