
## Karin Leandersson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1559039/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. BMC Cancer, 2012, 12, 306.	2.6	531
2	WNT5A induces release of exosomes containing pro-angiogenic and immunosuppressive factors from malignant melanoma cells. Molecular Cancer, 2014, 13, 88.	19.2	213
3	On the origin of myeloid-derived suppressor cells. Oncotarget, 2017, 8, 3649-3665.	1.8	156
4	Wnt-5a/Ca 2+ -Induced NFAT Activity Is Counteracted by Wnt-5a/Yes-Cdc42-Casein Kinase 1α Signaling in Human Mammary Epithelial Cells. Molecular and Cellular Biology, 2006, 26, 6024-6036.	2.3	144
5	Cancer-associated fibroblast-secreted CXCL16 attracts monocytes to promote stroma activation in triple-negative breast cancers. Nature Communications, 2016, 7, 13050.	12.8	135
6	A high frequency of MDSCs in sepsis patients, with the granulocytic subtype dominating in gram-positive cases. Journal of Leukocyte Biology, 2014, 96, 685-693.	3.3	128
7	Systemic Monocytic-MDSCs Are Generated from Monocytes and Correlate with Disease Progression in Breast Cancer Patients. PLoS ONE, 2015, 10, e0127028.	2.5	116
8	Infiltration of CD3+ and CD68+ cells in bladder cancer is subtype specific and affects the outcome of patients with muscle-invasive tumors11Grant support: The Swedish Cancer Society, the Swedish research council, the Nilsson Cancer foundation, the BioCARE Strategic Cancer Research program, the Lund Medical Faculty, and FoU Landstinget Kronoberg and Södra Regionvårdnänden Urologic Oncology: Seminars and Original Investigations, 2014, 32, 791-797.	1.6	106
9	Wht5a Induces a Tolerogenic Phenotype of Macrophages in Sepsis and Breast Cancer Patients. Journal of Immunology, 2012, 188, 5448-5458.	0.8	100
10	Expression and signaling activity of Wnt-5a/discoidin domain receptor-1 and Syk plays distinct but decisive roles in breast cancer patient survival. Clinical Cancer Research, 2005, 11, 520-8.	7.0	89
11	Wnt-5a mRNA translation is suppressed by the Elav-like protein HuR in human breast epithelial cells. Nucleic Acids Research, 2006, 34, 3988-3999.	14.5	86
12	The Generation and Identity of Human Myeloid-Derived Suppressor Cells. Frontiers in Oncology, 2020, 10, 109.	2.8	77
13	Cartilage oligomeric matrix protein contributes to the development and metastasis of breast cancer. Oncogene, 2016, 35, 5585-5596.	5.9	74
14	Prognostic stromal gene signatures in breast cancer. Breast Cancer Research, 2015, 17, 23.	5.0	67
15	The clinical impact of tumourâ€infiltrating lymphocytes in colorectal cancer differs by anatomical subsite: A cohort study. International Journal of Cancer, 2017, 141, 1654-1666.	5.1	65
16	Expression of programmed cell death protein 1 (PD-1) and its ligand PD-L1 in colorectal cancer: Relationship with sidedness and prognosis. OncoImmunology, 2018, 7, e1465165.	4.6	59
17	The integrative clinical impact of tumor-infiltrating T lymphocytes and NK cells in relation to B lymphocyte and plasma cell density in esophageal and gastric adenocarcinoma. Oncotarget, 2017, 8, 72108-72126.	1.8	53
18	The STAT3 inhibitor galiellalactone inhibits the generation of MDSCâ€like monocytes by prostate cancer cells and decreases immunosuppressive and tumorigenic factors. Prostate, 2019, 79, 1611-1621.	2.3	47

KARIN LEANDERSSON

#	Article	IF	CITATIONS
19	Clinical relevance of systemic monocytic-MDSCs in patients with metastatic breast cancer. Cancer Immunology, Immunotherapy, 2020, 69, 435-448.	4.2	44
20	Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer. Breast Cancer Research, 2015, 17, 130.	5.0	41
21	Quantitative, qualitative and spatial analysis of lymphocyte infiltration in periampullary and pancreatic adenocarcinoma. International Journal of Cancer, 2020, 146, 3461-3473.	5.1	39
22	Expression of PD-L1 and PD-1 in Chemoradiotherapy-NaÃ⁻ve Esophageal and Gastric Adenocarcinoma: Relationship With Mismatch Repair Status and Survival. Frontiers in Oncology, 2019, 9, 136.	2.8	36
23	S100A9 expressed in ERâ^'PgRâ^' breast cancers induces inflammatory cytokines and is associated with an impaired overall survival. British Journal of Cancer, 2015, 113, 1234-1243.	6.4	35
24	Docetaxel promotes the generation of anti-tumorigenic human macrophages. Experimental Cell Research, 2018, 362, 525-531.	2.6	34
25	Clinical impact of T cells, B cells and the PD-1/PD-L1 pathway in muscle invasive bladder cancer: a comparative study of transurethral resection and cystectomy specimens. Oncolmmunology, 2019, 8, e1644108.	4.6	34
26	The clinical importance of tumour-infiltrating macrophages and dendritic cells in periampullary adenocarcinoma differs by morphological subtype. Journal of Translational Medicine, 2017, 15, 152.	4.4	33
27	The Prognostic Impact of NK/NKT Cell Density in Periampullary Adenocarcinoma Differs by Morphological Type and Adjuvant Treatment. PLoS ONE, 2016, 11, e0156497.	2.5	32
28	Papillary renal cell carcinoma-derived chemerin, IL-8, and CXCL16 promote monocyte recruitment and differentiation into foam-cell macrophages. Laboratory Investigation, 2017, 97, 1296-1305.	3.7	28
29	Prognostic implications of the expression levels of different immunoglobulin heavy chain-encoding RNAs in early breast cancer. Npj Breast Cancer, 2020, 6, 28.	5.2	25
30	Wnt5a is a TLR2/4-ligand that induces tolerance in human myeloid cells. Communications Biology, 2019, 2, 176.	4.4	24
31	Infiltration of γâ¢Î´T cells, IL-17+ T cells and FoxP3+ T cells in human breast cancer. Cancer Biomarkers, 2018, 20, 395-409.	1.7	22
32	Wnt5a Inhibits Human Monocyteâ€Derived Myeloid Dendritic Cell Generation. Scandinavian Journal of Immunology, 2013, 78, 194-204.	2.7	21
33	Impact of systemic therapy on circulating leukocyte populations in patients with metastatic breast cancer. Scientific Reports, 2019, 9, 13451.	3.3	21
34	Tumor-Associated CD68+, CD163+, and MARCO+ Macrophages as Prognostic Biomarkers in Patients With Treatment-Naìve Gastroesophageal Adenocarcinoma. Frontiers in Oncology, 2020, 10, 534761.	2.8	20
35	Complement inhibitor factor H expressed by breast cancer cells differentiates CD14 <sup>+</sup> human monocytes into immunosuppressive macrophages. Oncolmmunology, 2020, 9, 1731135.	4.6	20
36	Dual mechanisms of action of the RNA-binding protein human antigen R explains its regulatory effect on melanoma cell migration. Translational Research, 2016, 172, 45-60.	5.0	19

KARIN LEANDERSSON

#	Article	IF	CITATIONS
37	Topographical Distribution and Spatial Interactions of Innate and Semi-Innate Immune Cells in Pancreatic and Other Periampullary Adenocarcinoma. Frontiers in Immunology, 2020, 11, 558169.	4.8	18
38	Infiltration of NK and plasma cells is associated with a distinct immune subset in nonâ€small cell lung cancer. Journal of Pathology, 2021, 255, 243-256.	4.5	17
39	Heterogeneity among septic shock patients in a set of immunoregulatory markers. European Journal of Clinical Microbiology and Infectious Diseases, 2014, 33, 313-324.	2.9	15
40	T cells, B cells, and PD-L1 expression in esophageal and gastric adenocarcinoma before and after neoadjuvant chemotherapy: relationship with histopathological response and survival. Oncolmmunology, 2021, 10, 1921443.	4.6	14
41	Human G-MDSCs are neutrophils at distinct maturation stages promoting tumor growth in breast cancer. Life Science Alliance, 2020, 3, e202000893.	2.8	14
42	The Immune Landscape of Colorectal Cancer. Cancers, 2021, 13, 5545.	3.7	14
43	Localization and Regulation of Polymeric Ig Receptor in Healthy and Diseased Human Kidney. American Journal of Pathology, 2019, 189, 1933-1944.	3.8	10
44	Co-localization of CD169 <sup>+</sup> macrophages and cancer cells in lymph node metastases of breast cancer patients is linked to improved prognosis and PDL1 expression. OncoImmunology, 2020, 9, 1848067.	4.6	9
45	Chemotherapy, host response and molecular dynamics in periampullary cancer: the CHAMP study. BMC Cancer, 2020, 20, 308.	2.6	9
46	High Infiltration of CD68 <sup>+</sup> /CD163 <sup>â^</sup> Macrophages Is an Adverse Prognostic Factor after Neoadjuvant Chemotherapy in Esophageal and Gastric Adenocarcinoma. Journal of Innate Immunity, 2022, 14, 615-628.	3.8	8
47	Peripheral Blood Mononuclear Cell Populations Correlate with Outcome in Patients with Metastatic Breast Cancer. Cells, 2022, 11, 1639.	4.1	8
48	Inflammatory macrophage derived TNFα downregulates estrogen receptor α via FOXO3a inactivation in human breast cancer cells. Experimental Cell Research, 2020, 390, 111932.	2.6	7
49	T cells developing in fetal thymus of T-cell receptor α-chain transgenic mice colonize γδT-cell-specific epithelial niches but lack long-term reconstituting potential. Immunology, 2006, 119, 134-142.	4.4	6
50	Pre-diagnostic anthropometry, sex, and risk of colorectal cancer according to tumor immune cell composition. Oncolmmunology, 2019, 8, e1664275.	4.6	5
51	Deletion of Nemo-like Kinase in T Cells Reduces Single-Positive CD8+ Thymocyte Population. Journal of Immunology, 2020, 205, 1830-1841.	0.8	4
52	Branching Copy-Number Evolution and Parallel Immune Profiles across the Regional Tumor Space of Resected Pancreatic Cancer. Molecular Cancer Research, 2022, 20, 749-761.	3.4	3
53	The prognostic impact of tumor-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite Journal of Clinical Oncology, 2017, 35, 47-47.	1.6	1
54	Prognostic significance of professional antigen presenting cells according to morphological subtype of periampullary adenocarcinoma Journal of Clinical Oncology, 2017, 35, 121-121.	1.6	0