

Karin Leandersson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1559039/publications.pdf>

Version: 2024-02-01

54
papers

2,945
citations

230014

27
h-index

198040

52
g-index

55
all docs

55
docs citations

55
times ranked

6622
citing authors

#	ARTICLE	IF	CITATIONS
1	Branching Copy-Number Evolution and Parallel Immune Profiles across the Regional Tumor Space of Resected Pancreatic Cancer. <i>Molecular Cancer Research</i> , 2022, 20, 749-761.	1.5	3
2	High Infiltration of CD68 ⁺ /CD163 ⁺ Macrophages Is an Adverse Prognostic Factor after Neoadjuvant Chemotherapy in Esophageal and Gastric Adenocarcinoma. <i>Journal of Innate Immunity</i> , 2022, 14, 615-628.	1.8	8
3	Peripheral Blood Mononuclear Cell Populations Correlate with Outcome in Patients with Metastatic Breast Cancer. <i>Cells</i> , 2022, 11, 1639.	1.8	8
4	T cells, B cells, and PD-L1 expression in esophageal and gastric adenocarcinoma before and after neoadjuvant chemotherapy: relationship with histopathological response and survival. <i>Oncolmmunology</i> , 2021, 10, 1921443.	2.1	14
5	Infiltration of NK and plasma cells is associated with a distinct immune subset in non-small cell lung cancer. <i>Journal of Pathology</i> , 2021, 255, 243-256.	2.1	17
6	The Immune Landscape of Colorectal Cancer. <i>Cancers</i> , 2021, 13, 5545.	1.7	14
7	Clinical relevance of systemic monocytic-MDSCs in patients with metastatic breast cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 435-448.	2.0	44
8	Prognostic implications of the expression levels of different immunoglobulin heavy chain-encoding RNAs in early breast cancer. <i>Npj Breast Cancer</i> , 2020, 6, 28.	2.3	25
9	Deletion of Nemo-like Kinase in T Cells Reduces Single-Positive CD8 ⁺ Thymocyte Population. <i>Journal of Immunology</i> , 2020, 205, 1830-1841.	0.4	4
10	Co-localization of CD169 ⁺ macrophages and cancer cells in lymph node metastases of breast cancer patients is linked to improved prognosis and PDL1 expression. <i>Oncolmmunology</i> , 2020, 9, 1848067.	2.1	9
11	Topographical Distribution and Spatial Interactions of Innate and Semi-Innate Immune Cells in Pancreatic and Other Periampullary Adenocarcinoma. <i>Frontiers in Immunology</i> , 2020, 11, 558169.	2.2	18
12	Tumor-Associated CD68 ⁺ , CD163 ⁺ , and MARCO ⁺ Macrophages as Prognostic Biomarkers in Patients With Treatment-Naïve Gastroesophageal Adenocarcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 534761.	1.3	20
13	Inflammatory macrophage derived TNF α downregulates estrogen receptor α via FOXO3a inactivation in human breast cancer cells. <i>Experimental Cell Research</i> , 2020, 390, 111932.	1.2	7
14	The Generation and Identity of Human Myeloid-Derived Suppressor Cells. <i>Frontiers in Oncology</i> , 2020, 10, 109.	1.3	77
15	Quantitative, qualitative and spatial analysis of lymphocyte infiltration in periampullary and pancreatic adenocarcinoma. <i>International Journal of Cancer</i> , 2020, 146, 3461-3473.	2.3	39
16	Chemotherapy, host response and molecular dynamics in periampullary cancer: the CHAMP study. <i>BMC Cancer</i> , 2020, 20, 308.	1.1	9
17	Complement inhibitor factor H expressed by breast cancer cells differentiates CD14 ⁺ human monocytes into immunosuppressive macrophages. <i>Oncolmmunology</i> , 2020, 9, 1731135.	2.1	20
18	Human G-MDSCs are neutrophils at distinct maturation stages promoting tumor growth in breast cancer. <i>Life Science Alliance</i> , 2020, 3, e202000893.	1.3	14

#	ARTICLE	IF	CITATIONS
19	The STAT3 inhibitor galiellalactone inhibits the generation of MDSC-like monocytes by prostate cancer cells and decreases immunosuppressive and tumorigenic factors. <i>Prostate</i> , 2019, 79, 1611-1621.	1.2	47
20	Clinical impact of T cells, B cells and the PD-1/PD-L1 pathway in muscle invasive bladder cancer: a comparative study of transurethral resection and cystectomy specimens. <i>Oncolmmunology</i> , 2019, 8, e1644108.	2.1	34
21	Impact of systemic therapy on circulating leukocyte populations in patients with metastatic breast cancer. <i>Scientific Reports</i> , 2019, 9, 13451.	1.6	21
22	Localization and Regulation of Polymeric Ig Receptor in Healthy and Diseased Human Kidney. <i>American Journal of Pathology</i> , 2019, 189, 1933-1944.	1.9	10
23	Wnt5a is a TLR2/4-ligand that induces tolerance in human myeloid cells. <i>Communications Biology</i> , 2019, 2, 176.	2.0	24
24	Expression of PD-L1 and PD-1 in Chemoradiotherapy-Naïve Esophageal and Gastric Adenocarcinoma: Relationship With Mismatch Repair Status and Survival. <i>Frontiers in Oncology</i> , 2019, 9, 136.	1.3	36
25	Pre-diagnostic anthropometry, sex, and risk of colorectal cancer according to tumor immune cell composition. <i>Oncolmmunology</i> , 2019, 8, e1664275.	2.1	5
26	Docetaxel promotes the generation of anti-tumorigenic human macrophages. <i>Experimental Cell Research</i> , 2018, 362, 525-531.	1.2	34
27	Infiltration of CD4+ T cells, IL-17+ T cells and FoxP3+ T cells in human breast cancer. <i>Cancer Biomarkers</i> , 2018, 20, 395-409.	0.8	22
28	Expression of programmed cell death protein 1 (PD-1) and its ligand PD-L1 in colorectal cancer: Relationship with sidedness and prognosis. <i>Oncolmmunology</i> , 2018, 7, e1465165.	2.1	59
29	The clinical impact of tumour-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite: A cohort study. <i>International Journal of Cancer</i> , 2017, 141, 1654-1666.	2.3	65
30	Papillary renal cell carcinoma-derived chemerin, IL-8, and CXCL16 promote monocyte recruitment and differentiation into foam-cell macrophages. <i>Laboratory Investigation</i> , 2017, 97, 1296-1305.	1.7	28
31	The clinical importance of tumour-infiltrating macrophages and dendritic cells in periampullary adenocarcinoma differs by morphological subtype. <i>Journal of Translational Medicine</i> , 2017, 15, 152.	1.8	33
32	On the origin of myeloid-derived suppressor cells. <i>Oncotarget</i> , 2017, 8, 3649-3665.	0.8	156
33	The prognostic impact of tumor-infiltrating lymphocytes in colorectal cancer differs by anatomical subsite.. <i>Journal of Clinical Oncology</i> , 2017, 35, 47-47.	0.8	1
34	The integrative clinical impact of tumor-infiltrating T lymphocytes and NK cells in relation to B lymphocyte and plasma cell density in esophageal and gastric adenocarcinoma. <i>Oncotarget</i> , 2017, 8, 72108-72126.	0.8	53
35	Prognostic significance of professional antigen presenting cells according to morphological subtype of periampullary adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2017, 35, 121-121.	0.8	0
36	Cartilage oligomeric matrix protein contributes to the development and metastasis of breast cancer. <i>Oncogene</i> , 2016, 35, 5585-5596.	2.6	74

#	ARTICLE	IF	CITATIONS
37	Cancer-associated fibroblast-secreted CXCL16 attracts monocytes to promote stroma activation in triple-negative breast cancers. <i>Nature Communications</i> , 2016, 7, 13050.	5.8	135
38	Dual mechanisms of action of the RNA-binding protein human antigen R explains its regulatory effect on melanoma cell migration. <i>Translational Research</i> , 2016, 172, 45-60.	2.2	19
39	The Prognostic Impact of NK/NKT Cell Density in Periampullary Adenocarcinoma Differs by Morphological Type and Adjuvant Treatment. <i>PLoS ONE</i> , 2016, 11, e0156497.	1.1	32
40	Systemic Monocytic-MDSCs Are Generated from Monocytes and Correlate with Disease Progression in Breast Cancer Patients. <i>PLoS ONE</i> , 2015, 10, e0127028.	1.1	116
41	Prognostic stromal gene signatures in breast cancer. <i>Breast Cancer Research</i> , 2015, 17, 23.	2.2	67
42	S100A9 expressed in ER ⁺ /PgR ⁺ breast cancers induces inflammatory cytokines and is associated with an impaired overall survival. <i>British Journal of Cancer</i> , 2015, 113, 1234-1243.	2.9	35
43	Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer. <i>Breast Cancer Research</i> , 2015, 17, 130.	2.2	41
44	Heterogeneity among septic shock patients in a set of immunoregulatory markers. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2014, 33, 313-324.	1.3	15
45	WNT5A induces release of exosomes containing pro-angiogenic and immunosuppressive factors from malignant melanoma cells. <i>Molecular Cancer</i> , 2014, 13, 88.	7.9	213
46	Infiltration of CD3 ⁺ and CD68 ⁺ cells in bladder cancer is subtype specific and affects the outcome of patients with muscle-invasive tumors Grant support: The Swedish Cancer Society, the Swedish research council, the Nilsson Cancer foundation, the BioCARE Strategic Cancer Research program, the Lund Medical Faculty, and FoU Landstinget Kronoberg and SÄdra Region <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 791-797.	0.8	106
47	A high frequency of MDSCs in sepsis patients, with the granulocytic subtype dominating in gram-positive cases. <i>Journal of Leukocyte Biology</i> , 2014, 96, 685-693.	1.5	128
48	Wnt5a Inhibits Human Monocyte-Derived Myeloid Dendritic Cell Generation. <i>Scandinavian Journal of Immunology</i> , 2013, 78, 194-204.	1.3	21
49	Wnt5a Induces a Tolerogenic Phenotype of Macrophages in Sepsis and Breast Cancer Patients. <i>Journal of Immunology</i> , 2012, 188, 5448-5458.	0.4	100
50	The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. <i>BMC Cancer</i> , 2012, 12, 306.	1.1	531
51	T cells developing in fetal thymus of T-cell receptor α -chain transgenic mice colonize β T-cell-specific epithelial niches but lack long-term reconstituting potential. <i>Immunology</i> , 2006, 119, 134-142.	2.0	6
52	Wnt-5a mRNA translation is suppressed by the Elav-like protein HuR in human breast epithelial cells. <i>Nucleic Acids Research</i> , 2006, 34, 3988-3999.	6.5	86
53	Wnt-5a/Ca ²⁺ -Induced NFAT Activity Is Counteracted by Wnt-5a/Yes-Cdc42-Casein Kinase 1α Signaling in Human Mammary Epithelial Cells. <i>Molecular and Cellular Biology</i> , 2006, 26, 6024-6036.	1.1	144
54	Expression and signaling activity of Wnt-5a/discoidin domain receptor-1 and Syk plays distinct but decisive roles in breast cancer patient survival. <i>Clinical Cancer Research</i> , 2005, 11, 520-8.	3.2	89