

# Andreas Villunger

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1561478/publications.pdf>

Version: 2024-02-01

141  
papers

14,979  
citations

38742

50  
h-index

19749

117  
g-index

151  
all docs

151  
docs citations

151  
times ranked

21577  
citing authors

#	ARTICLE	IF	CITATIONS
1	BCL-2 family protein tBID can act as a BAX-like effector of apoptosis. <i>EMBO Journal</i> , 2022, 41, e108690.	7.8	74
2	PIDD1 in cell cycle control, sterile inflammation and cell death. <i>Biochemical Society Transactions</i> , 2022, 50, 813-824.	3.4	11
3	The miR-26 family regulates early B cell development and transformation. <i>Life Science Alliance</i> , 2022, 5, e202101303.	2.8	5
4	The SKP2-p27 axis defines susceptibility to cell death upon CHK1 inhibition. <i>Molecular Oncology</i> , 2022, 16, 2771-2787.	4.6	4
5	Cell-Specific Immune Regulation by Glucocorticoids in Murine Models of Infection and Inflammation. <i>Cells</i> , 2022, 11, 2126.	4.1	3
6	P53 clears aneuploid cells by entosis. <i>Cell Death and Differentiation</i> , 2021, 28, 818-820.	11.2	10
7	Differential roles of miR-15a/16-1 and miR-497/195 clusters in immune cell development and homeostasis. <i>FEBS Journal</i> , 2021, 288, 1533-1545.	4.7	6
8	Biallelic mutations in the death domain of PIDD1 impair caspase-2 activation and are associated with intellectual disability. <i>Translational Psychiatry</i> , 2021, 11, 1.	4.8	334
9	Polyploidy control in hepatic health and disease. <i>Journal of Hepatology</i> , 2021, 75, 1177-1191.	3.7	19
10	Centriolar distal appendages activate the centrosome-PIDDosome-p53 signalling axis via ANKRD26. <i>EMBO Journal</i> , 2021, 40, e104844.	7.8	40
11	At a Crossroads to Cancer: How p53-Induced Cell Fate Decisions Secure Genome Integrity. <i>International Journal of Molecular Sciences</i> , 2021, 22, 10883.	4.1	30
12	Lack of Bmf Facilitates the Selection of Highly Responsive B-Cell Receptor Clones in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2021, 138, 1543-1543.	1.4	0
13	The BH3-only protein NOXA serves as an independent predictor of breast cancer patient survival and defines susceptibility to microtubule targeting agents. <i>Cell Death and Disease</i> , 2021, 12, 1151.	6.3	11
14	Dynein light chain binding determines complex formation and posttranslational stability of the Bcl-2 family members Bmf and Bim. <i>Cell Death and Differentiation</i> , 2020, 27, 434-450.	11.2	19
15	Cell-Cycle Cross Talk with Caspases and Their Substrates. <i>Cold Spring Harbor Perspectives in Biology</i> , 2020, 12, a036475.	5.5	17
16	Uncovering the PIDDosome and caspase-2 as regulators of organogenesis and cellular differentiation. <i>Cell Death and Differentiation</i> , 2020, 27, 2037-2047.	11.2	24
17	Drp1 modulates mitochondrial stress responses to mitotic arrest. <i>Cell Death and Differentiation</i> , 2020, 27, 2620-2634.	11.2	18
18	E2F-Family Members Engage the PIDDosome to Limit Hepatocyte Ploidy in Liver Development and Regeneration. <i>Developmental Cell</i> , 2020, 52, 335-349.e7.	7.0	40

#	ARTICLE	IF	CITATIONS
19	MARCH5-dependent degradation of MCL1/NOXA complexes defines susceptibility to antimitotic drug treatment. <i>Cell Death and Differentiation</i> , 2020, 27, 2297-2312.	11.2	31
20	PIDDosome-induced p53-dependent ploidy restriction facilitates hepatocarcinogenesis. <i>EMBO Reports</i> , 2020, 21, e50893.	4.5	29
21	Checkpoint kinase 1 is essential for fetal and adult hematopoiesis. <i>EMBO Reports</i> , 2019, 20, e47026.	4.5	15
22	TET enzymes control antibody production and shape the mutational landscape in germinal centre B cells. <i>FEBS Journal</i> , 2019, 286, 3566-3581.	4.7	37
23	CHK1 dosage in germinal center B cells controls humoral immunity. <i>Cell Death and Differentiation</i> , 2019, 26, 2551-2567.	11.2	14
24	Glucocorticoid Receptor-Deficient Foxp3+ Regulatory T Cells Fail to Control Experimental Inflammatory Bowel Disease. <i>Frontiers in Immunology</i> , 2019, 10, 472.	4.8	28
25	RIPK1 and Caspase-8 Ensure Chromosome Stability Independently of Their Role in Cell Death and Inflammation. <i>Molecular Cell</i> , 2019, 73, 413-428.e7.	9.7	50
26	BIRC3 Expression Predicts CLL Progression and Defines Treatment Sensitivity via Enhanced NF- $\kappa$ B Nuclear Translocation. <i>Clinical Cancer Research</i> , 2019, 25, 1901-1912.	7.0	23
27	Perturbing mitosis for anti-cancer therapy: is cell death the only answer?. <i>EMBO Reports</i> , 2018, 19, .	4.5	67
28	Differential effects of Vav-promoter-driven overexpression of BCLX and BFL1 on lymphocyte survival and B cell lymphomagenesis. <i>FEBS Journal</i> , 2018, 285, 1403-1418.	4.7	5
29	Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. <i>Cell Death and Differentiation</i> , 2018, 25, 486-541.	11.2	4,036
30	BOK promotes chemical-induced hepatocarcinogenesis in mice. <i>Cell Death and Differentiation</i> , 2018, 25, 708-720.	11.2	26
31	CDK6 Antagonizes p53-Induced Responses during Tumorigenesis. <i>Cancer Discovery</i> , 2018, 8, 884-897.	9.4	53
32	Deletion of the p53 Target Gene PUIMA Prevents Bone Marrow Failure in a Dyskeratosis Congenita Mouse Model. <i>Blood</i> , 2018, 132, 648-648.	1.4	0
33	The PIDDosome activates p53 in response to supernumerary centrosomes. <i>Genes and Development</i> , 2017, 31, 34-45.	5.9	153
34	The BCL-2 pro-survival protein A1 is dispensable for T cell homeostasis on viral infection. <i>Cell Death and Differentiation</i> , 2017, 24, 523-533.	11.2	29
35	Characterisation of mice lacking all functional isoforms of the pro-survival BCL-2 family member A1 reveals minor defects in the haematopoietic compartment. <i>Cell Death and Differentiation</i> , 2017, 24, 534-545.	11.2	60
36	There is something about BOK we just don't get yet. <i>FEBS Journal</i> , 2017, 284, 708-710.	4.7	6

#	ARTICLE	IF	CITATIONS
37	DNA-binding of the Tet-transactivator curtails antigen-induced lymphocyte activation in mice. <i>Nature Communications</i> , 2017, 8, 1028.	12.8	8
38	Signalling strength determines proapoptotic functions of STING. <i>Nature Communications</i> , 2017, 8, 427.	12.8	321
39	The miRâ€15 family reinforces the transition from proliferation to differentiation in preâ€B cells. <i>EMBO Reports</i> , 2017, 18, 1604-1617.	4.5	34
40	Checkpoint kinase 1 is essential for normal B cell development and lymphomagenesis. <i>Nature Communications</i> , 2017, 8, 1697.	12.8	28
41	The corepressor NCOR1 regulates the survival of single-positive thymocytes. <i>Scientific Reports</i> , 2017, 7, 15928.	3.3	14
42	The resurrection of the PIDDosome â€ emerging roles in the DNA-damage response and centrosome surveillance. <i>Journal of Cell Science</i> , 2017, 130, 3779-3787.	2.0	39
43	The cyanobacterial metabolite nocuolin a is a natural oxadiazine that triggers apoptosis in human cancer cells. <i>PLoS ONE</i> , 2017, 12, e0172850.	2.5	43
44	T-cell autonomous death induced by regeneration of inert glucocorticoid metabolites. <i>Cell Death and Disease</i> , 2017, 8, e2948-e2948.	6.3	17
45	The RNA-binding protein tristetraprolin schedules apoptosis of pathogen-engaged neutrophils during bacterial infection. <i>Journal of Clinical Investigation</i> , 2017, 127, 2051-2065.	8.2	28
46	Janus Kinase 1 Is Essential for Inflammatory Cytokine Signaling and Mammary Gland Remodeling. <i>Molecular and Cellular Biology</i> , 2016, 36, 1673-1690.	2.3	24
47	MOMP in the absence of BH3-only proteins. <i>Genes and Development</i> , 2016, 30, 878-880.	5.9	8
48	Canonical NF-Î²B signaling is uniquely required for the long-term persistence of functional mature B cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 5065-5070.	7.1	20
49	LDHA-Associated Lactic Acid Production Blunts Tumor Immunosurveillance by T and NK Cells. <i>Cell Metabolism</i> , 2016, 24, 657-671.	16.2	1,126
50	Interrogating the relevance of mitochondrial apoptosis for vertebrate development and postnatal tissue homeostasis. <i>Genes and Development</i> , 2016, 30, 2133-2151.	5.9	56
51	Cooperation of ETV6/RUNX1 and BCL2 enhances immunoglobulin production and accelerates glomerulonephritis in transgenic mice. <i>Oncotarget</i> , 2016, 7, 12191-12205.	1.8	6
52	Beclin 1 is dispensable for chromosome congression and proper outer kinetochore assembly. <i>EMBO Reports</i> , 2015, 16, 1233-1236.	4.5	5
53	Embryonic stem cell differentiation requires full length Chd1. <i>Scientific Reports</i> , 2015, 5, 8007.	3.3	23
54	The p53 binding protein PDCD5 is not rate-limiting in DNA damage induced cell death. <i>Scientific Reports</i> , 2015, 5, 11268.	3.3	6

#	ARTICLE	IF	CITATIONS
55	The Nuclear Orphan Receptor NR2F6 Is a Central Checkpoint for Cancer Immune Surveillance. <i>Cell Reports</i> , 2015, 12, 2072-2085.	6.4	47
56	Knockdown of the Antiapoptotic Bcl-2 Family Member A1/Bfl-1 Protects Mice from Anaphylaxis. <i>Journal of Immunology</i> , 2015, 194, 1316-1322.	0.8	16
57	The NOXA-“MCL1”-BIM axis defines lifespan on extended mitotic arrest. <i>Nature Communications</i> , 2015, 6, 6891.	12.8	86
58	Lessons from gain- and loss-of-function models of pro-survival Bcl2 family proteins: implications for targeted therapy. <i>FEBS Journal</i> , 2015, 282, 834-849.	4.7	53
59	Replenishment of the B cell compartment after doxorubicin-induced hematopoietic toxicity is facilitated by STAT1. <i>Journal of Leukocyte Biology</i> , 2014, 95, 853-866.	3.3	6
60	Chemokine-mediated redirection of T cells constitutes a critical mechanism of glucocorticoid therapy in autoimmune CNS responses. <i>Acta Neuropathologica</i> , 2014, 127, 713-729.	7.7	46
61	SCID-dependent release of mitochondrial SMAC dampens XIAP-mediated immunity against <i>Shigella</i> . <i>EMBO Journal</i> , 2014, 33, 2171-2187.	7.8	52
62	Stop competing, start talking!. <i>EMBO Journal</i> , 2014, 33, 1849-1851.	7.8	5
63	Deregulated cell death and lymphocyte homeostasis cause premature lethality in mice lacking the BH3-only proteins Bim and Bmf. <i>Blood</i> , 2014, 123, 2652-2662.	1.4	40
64	Transient Bcl-XL Overexpression in Donor Stem Cells Increases Efficacy of Hematopoietic Stem Cell Transplantation without Increasing the Risk of Leukemogenesis. <i>Blood</i> , 2014, 124, 4350-4350.	1.4	1
65	AICAR induces Bax/Bak-dependent apoptosis through upregulation of the BH3-only proteins Bim and Noxa in mouse embryonic fibroblasts. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2013, 18, 1008-1016.	4.9	21
66	BH3-only protein Noxa contributes to apoptotic control of stress-erythropoiesis. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2013, 18, 1306-1318.	4.9	10
67	Haematopoietic stem cell survival and transplantation efficacy is limited by the BH3-only proteins Bim and Bmf. <i>EMBO Molecular Medicine</i> , 2013, 5, 122-136.	6.9	25
68	Lapatinib and doxorubicin enhance the Src-dependent antitumor immune response. <i>European Journal of Immunology</i> , 2013, 43, 2718-2729.	2.9	108
69	Possible pitfalls investigating cell death responses in genetically engineered mouse models and derived cell lines. <i>Methods</i> , 2013, 61, 130-137.	3.8	8
70	Neuronal caspase 2 activity and function requires RAIDD, but not PIDD. <i>Biochemical Journal</i> , 2012, 444, 591-599.	3.7	37
71	PINCH-1 promotes Bcl-2-dependent survival signalling and inhibits JNK-mediated apoptosis in the primitive endoderm.. <i>Journal of Cell Science</i> , 2012, 125, 5233-40.	2.0	25
72	PIDDosome-independent tumor suppression by Caspase-2. <i>Cell Death and Differentiation</i> , 2012, 19, 1722-1732.	11.2	60

#	ARTICLE	IF	CITATIONS
73	Caspase-2 at a glance. <i>Journal of Cell Science</i> , 2012, 125, 5911-5915.	2.0	74
74	Targeting antiapoptotic A1/Bfl-1 by in vivo RNAi reveals multiple roles in leukocyte development in mice. <i>Blood</i> , 2012, 119, 6032-6042.	1.4	52
75	Necrosis-like death can engage multiple pro-apoptotic Bcl-2 protein family members. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2012, 17, 1197-1209.	4.9	48
76	Upregulation of miRâ€24 is associated with a decreased DNA damage response upon etoposide treatment in highly differentiated CD8<sup>+</sup> T cells sensitizing them to apoptotic cell death. <i>Aging Cell</i> , 2012, 11, 579-587.	6.7	78
77	Defective cell death signalling along the Bcl-2 regulated apoptosis pathway compromises Treg cell development and limits their functionality in mice. <i>Journal of Autoimmunity</i> , 2012, 38, 59-69.	6.5	36
78	A1/Bfl-1 in leukocyte development and cell death. <i>Experimental Cell Research</i> , 2012, 318, 1291-1303.	2.6	44
79	GSK3 TIPping Off p53 to Unleash PUMA. <i>Molecular Cell</i> , 2011, 42, 555-556.	9.7	4
80	The cooperating mutation or â€œsecond hitâ€ determines the immunologic visibility toward MYC-induced murine lymphomas. <i>Blood</i> , 2011, 118, 4635-4645.	1.4	30
81	Genome-wide association analysis in primary sclerosing cholangitis identifies two non-HLA susceptibility loci. <i>Nature Genetics</i> , 2011, 43, 17-19.	21.4	221
82	Shaping the Tâ€cell repertoire: a matter of life and death. <i>Immunology and Cell Biology</i> , 2011, 89, 33-39.	2.3	33
83	Apoptosis: A barrier against cancer no more?. <i>Hepatology</i> , 2011, 54, 1121-1124.	7.3	7
84	Deciphering the Molecular Events Necessary for Synergistic Tumor Cell Apoptosis Mediated by the Histone Deacetylase Inhibitor Vorinostat and the BH3 Mimetic ABT-737. <i>Cancer Research</i> , 2011, 71, 3603-3615.	0.9	51
85	BCL-2 Modifying Factor (BMF) Is a Central Regulator of Anoikis in Human Intestinal Epithelial Cells. <i>Journal of Biological Chemistry</i> , 2011, 286, 26533-26540.	3.4	42
86	Induction of Noxa-Mediated Apoptosis by Modified Vaccinia Virus Ankara Depends on Viral Recognition by Cytosolic Helicases, Leading to IRF-3/IFN-Î²-Dependent Induction of Pro-Apoptotic Noxa. <i>PLoS Pathogens</i> , 2011, 7, e1002083.	4.7	48
87	Generation and Evaluation of an IPTG-Regulated Version of Vav-Gene Promoter for Mouse Transgenesis. <i>PLoS ONE</i> , 2011, 6, e18051.	2.5	11
88	Suppression of B-cell lymphomagenesis by the BH3-only proteins Bmf and Bad. <i>Blood</i> , 2010, 115, 995-1005.	1.4	53
89	AICAR induces apoptosis independently of AMPK and p53 through up-regulation of the BH3-only proteins BIM and NOXA in chronic lymphocytic leukemia cells. <i>Blood</i> , 2010, 116, 3023-3032.	1.4	95
90	Infiltrating CD11b<sup>+</sup>CD11c<sup>+</sup> cells have the potential to mediate inducible nitric oxide synthaseâ€dependent cell death in mammary carcinomas of HERâ€2/neu transgenic mice. <i>International Journal of Cancer</i> , 2010, 126, 896-908.	5.1	34

#	ARTICLE	IF	CITATIONS
91	Only the Strong Survive. <i>Immunity</i> , 2010, 32, 729-731.	14.3	3
92	Apoptosis of leukocytes triggered by acute DNA damage promotes lymphoma formation. <i>Genes and Development</i> , 2010, 24, 1602-1607.	5.9	95
93	The Anti-apoptotic Protein BCL2L1/Bcl-xL Is Neutralized by Pro-apoptotic PMAIP1/Noxa in Neuroblastoma, Thereby Determining Bortezomib Sensitivity Independent of Prosurvival MCL1 Expression. <i>Journal of Biological Chemistry</i> , 2010, 285, 6904-6912.	3.4	66
94	PUMA-mediated tumor suppression: A tale of two stories. <i>Cell Cycle</i> , 2010, 9, 4269-4275.	2.6	16
95	Apoptosis and necroptosis are induced in rainbow trout cell lines exposed to cadmium. <i>Aquatic Toxicology</i> , 2010, 99, 73-85.	4.0	63
96	Role for BH3-Only Protein NOXA In Growth-Factor Deprivation and Early Erythropoiesis. <i>Blood</i> , 2010, 116, 4235-4235.	1.4	0
97	Lack of the BH3-Only Proteins Bim, Bmf and Puma In Haematopoietic Stem and Progenitor Cells Facilitates Early Reconstitution and Long Term Haematopoiesis.. <i>Blood</i> , 2010, 116, 1542-1542.	1.4	0
98	Deletion of Puma and p21Waf1 In Mice Deactivates p53-Induced Cell Death and Cell Cycle Arrest, but Protects Mice From Irradiation-Induced Lymphomagenesis by a Mechanism Involving Hemopoietic Stem Cell Quiescence. <i>Blood</i> , 2010, 116, 90-90.	1.4	5
99	Loss of the pro-apoptotic BH3-only Bcl-2 family member Bim sustains B lymphopoiesis in the absence of IL-7. <i>International Immunology</i> , 2009, 21, 715-725.	4.0	20
100	Caspase-2 activation in the absence of PIDDosome formation. <i>Journal of Cell Biology</i> , 2009, 185, 291-303.	5.2	144
101	Bcl2 family proteins in carcinogenesis and the treatment of cancer. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2009, 14, 584-596.	4.9	288
102	BH3-only protein Bim more critical than Puma in tyrosine kinase inhibitorâ€“induced apoptosis of human leukemic cells and transduced hematopoietic progenitors carrying oncogenic FLT3. <i>Blood</i> , 2009, 113, 2302-2311.	1.4	31
103	The Nuclear Orphan Receptor NR2F6 Suppresses Lymphocyte Activation and T Helper 17-Dependent Autoimmunity. <i>Immunity</i> , 2008, 29, 205-216.	14.3	93
104	Loss of the BH3-only protein Bmf impairs B cell homeostasis and accelerates $\beta$ irradiationâ€“induced thymic lymphoma development. <i>Journal of Experimental Medicine</i> , 2008, 205, 641-655.	8.5	116
105	Ultraviolet radiation triggers apoptosis of fibroblasts and skin keratinocytes mainly via the BH3-only protein Noxa. <i>Journal of Cell Biology</i> , 2007, 176, 415-424.	5.2	96
106	Deletion of the BH3-only protein <i>puma</i> protects motoneurons from ER stress-induced apoptosis and delays motoneuron loss in ALS mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 20606-20611.	7.1	122
107	Impact of cellular lifespan on the T cell receptor repertoire. <i>European Journal of Immunology</i> , 2007, 37, 1978-1985.	2.9	8
108	FOXO3a-dependent regulation of Puma in response to cytokine/growth factor withdrawal. <i>Journal of Experimental Medicine</i> , 2006, 203, 1657-1663.	8.5	367

#	ARTICLE	IF	CITATIONS
109	Apoptosis in Activated T Cells – What Are the Triggers, and What the Signal Transducers?. <i>Cell Cycle</i> , 2006, 5, 2421-2424.	2.6	24
110	BH3-Only Proapoptotic Bcl-2 Family Members Noxa and Puma Mediate Neural Precursor Cell Death. <i>Journal of Neuroscience</i> , 2006, 26, 7257-7264.	3.6	61
111	p14 – MP1-MEK1 signaling regulates endosomal traffic and cellular proliferation during tissue homeostasis. <i>Journal of Cell Biology</i> , 2006, 175, 861-868.	5.2	195
112	The NF- $\kappa$ B regulator Bcl-3 and the BH3-only proteins Bim and Puma control the death of activated T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 10979-10984.	7.1	80
113	Bim and Bad mediate imatinib-induced killing of Bcr/Abl+ leukemic cells, and resistance due to their loss is overcome by a BH3 mimetic. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 14907-14912.	7.1	310
114	Puma cooperates with Bim, the rate-limiting BH3-only protein in cell death during lymphocyte development, in apoptosis induction. <i>Journal of Experimental Medicine</i> , 2006, 203, 2939-2951.	8.5	209
115	Key roles of BIM-driven apoptosis in epithelial tumors and rational chemotherapy. <i>Cancer Cell</i> , 2005, 7, 227-238.	16.8	276
116	Phagocytosis-Induced Apoptosis in Macrophages Is Mediated by Up-Regulation and Activation of the Bcl-2 Homology Domain 3-Only Protein Bim. <i>Journal of Immunology</i> , 2005, 174, 671-679.	0.8	52
117	Mutually Exclusive Subsets of BH3-Only Proteins Are Activated by the p53 and c-Jun N-Terminal Kinase/c-Jun Signaling Pathways during Cortical Neuron Apoptosis Induced by Arsenite. <i>Molecular and Cellular Biology</i> , 2005, 25, 8732-8747.	2.3	74
118	Death squads enlisted by the tumour suppressor p53. <i>Biochemical and Biophysical Research Communications</i> , 2005, 331, 786-798.	2.1	112
119	BH3-only proteins Puma and Bim are rate-limiting for $\hat{\gamma}$ -radiation – and glucocorticoid-induced apoptosis of lymphoid cells in vivo. <i>Blood</i> , 2005, 106, 4131-4138.	1.4	259
120	Negative selection of semimature CD4+8-HSA+ thymocytes requires the BH3-only protein Bim but is independent of death receptor signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 7052-7057.	7.1	71
121	The Bcl-2 protein family and its role in the development of neoplastic disease. <i>Experimental Gerontology</i> , 2004, 39, 1125-1135.	2.8	49
122	p53- and Drug-Induced Apoptotic Responses Mediated by BH3-Only Proteins Puma and Noxa. <i>Science</i> , 2003, 302, 1036-1038.	12.6	1,187
123	Essential role for the BH3-only protein Bim but redundant roles for Bax, Bcl-2, and Bcl-w in the control of granulocyte survival. <i>Blood</i> , 2003, 101, 2393-2400.	1.4	133
124	Bmf: A Proapoptotic BH3-Only Protein Regulated by Interaction with the Myosin V Actin Motor Complex, Activated by Anoikis. <i>Science</i> , 2001, 293, 1829-1832.	12.6	555
125	T cell expressed PKC $\hat{\zeta}$ demonstrates cell-type selective function. <i>European Journal of Immunology</i> , 2000, 30, 3645-3654.	2.9	54
126	Unique Structural and Functional Properties of the ATP-binding Domain of Atypical Protein Kinase C- $\hat{\zeta}$ . <i>Journal of Biological Chemistry</i> , 2000, 275, 33289-33296.	3.4	44



#	ARTICLE	IF	CITATIONS
127	Fas Ligand-Induced c-Jun Kinase Activation in Lymphoid Cells Requires Extensive Receptor Aggregation But Is Independent of DAXX, and Fas-Mediated Cell Death Does Not Involve DAXX, RIP, or RAIDD. <i>Journal of Immunology</i> , 2000, 165, 1337-1343.	0.8	61
128	FAS Ligand, Bcl-2, Granulocyte Colony-Stimulating Factor, and p38 Mitogen-Activated Protein Kinase. <i>Journal of Experimental Medicine</i> , 2000, 192, 647-658.	8.5	103
129	Evidence That Atypical Protein Kinase C $\beta$ and Atypical Protein Kinase C $\eta$ Participate in Ras-mediated Reorganization of the F-actin Cytoskeleton. <i>Journal of Cell Biology</i> , 1999, 144, 413-425.	5.2	134
130	The great escape: Is immune evasion required for tumor progression?. <i>Nature Medicine</i> , 1999, 5, 874-875.	30.7	33
131	Protein kinase C $\delta$ , a selective upstream regulator of JNK/SAPK and IL-2 promoter activation in Jurkat T cells. <i>European Journal of Immunology</i> , 1999, 29, 132-142.	2.9	110
132	Synergistic action of protein kinase C $\delta$ and calcineurin is sufficient for Fas ligand expression and induction of a crmA-sensitive apoptosis pathway in Jurkat T cells. <i>European Journal of Immunology</i> , 1999, 29, 3549-3561.	2.9	49
133	Transcriptional activation of c-fos by oncogenic Ha-Ras in mouse mammary epithelial cells requires the combined activities of PKC $\delta$ , $\mu$ and $\eta$ . <i>EMBO Journal</i> , 1998, 17, 4046-4055.	7.8	66
134	Functional granulocyte/macrophage colony stimulating factor receptor is constitutively expressed on neoplastic plasma cells and mediates tumour cell longevity. <i>British Journal of Haematology</i> , 1998, 102, 1069-1080.	2.5	13
135	On the Role and Significance of Fas (Apo-1/CD95) Ligand (FasL) Expression in Immune Privileged Tissues and Cancer Cells Using Multiple Myeloma as a Model*. <i>Leukemia and Lymphoma</i> , 1998, 31, 477-490.	1.3	28
136	The interleukin 1 $\beta$ -converting enzyme inhibitor CrmA prevents Apo1/Fas- but not glucocorticoid-induced poly(ADP-ribose) polymerase cleavage and apoptosis in lymphoblastic leukemia cells. <i>FEBS Letters</i> , 1997, 402, 36-40.	2.8	35
137	Expression of Apo $\beta$ 1/Fas (CD95), Bcl $\beta$ 2, Bax and Bcl $\beta$ 3 in myeloma cell lines: relationship between responsiveness to anti $\beta$ 1/Fas mab and p53 functional status. <i>British Journal of Haematology</i> , 1997, 97, 418-428.	2.5	39
138	Modulation of Apo-1/Fas (CD95)-induced programmed cell death in myeloma cells by interferon- $\gamma$ . <i>European Journal of Immunology</i> , 1996, 26, 3119-3126.	2.9	70
139	Constituents of autocrine IL-6 loops in myeloma cell lines and their targeting for suppression of neoplastic growth by antibody strategies. , 1996, 65, 498-505.		23
140	2 $\beta$ -Fluorodeoxycytidine (Gemcitabine) Induces Apoptosis in Myeloma Cell Lines Resistant to Steroids and 2 $\beta$ -Chlorodeoxyadenosine (2 $\beta$ -CdA). <i>Stem Cells</i> , 1996, 14, 351-362.	3.2	22
141	Lactogenic Hormone and Cell Type-Specific Control of the Whey Acidic Protein Gene Promoter in Transfected Mouse Cells. <i>Molecular Endocrinology</i> , 1991, 5, 1624-1632.	3.7	68