

Johann Lehrner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1567082/publications.pdf>

Version: 2024-02-01

73
papers

2,468
citations

270111

25
h-index

242451

47
g-index

78
all docs

78
docs citations

78
times ranked

3394
citing authors

#	ARTICLE	IF	CITATIONS
1	Motor Reaction Times as an Embedded Measure of Performance Validity: a Study with a Sample of Austrian Early Retirement Claimants. <i>Psychological Injury and Law</i> , 2022, 15, 200-212.	1.0	9
2	Symptom and Performance Validity Assessment in European Countries: an Update. <i>Psychological Injury and Law</i> , 2022, 15, 116-127.	1.0	8
3	Neuropsychological prediction of dementia using the neuropsychological test battery Vienna – A retrospective study. <i>Brain Disorders</i> , 2022, 5, 100028.	1.1	0
4	Reduction of physical activity during the COVID-19 pandemic is related to increased neuropsychiatric symptoms in memory clinic patients. <i>Clinical Medicine</i> , 2022, 22, 177-180.	0.8	4
5	No effect of thyroid hormones on 5-year mortality in patients with subjective cognitive decline, mild cognitive disorder, and Alzheimer's disease. <i>Journal of Neuroendocrinology</i> , 2022, 34, e13107.	1.2	5
6	Connectome Analysis in an Individual with SETD1B-Related Neurodevelopmental Disorder and Epilepsy. <i>Journal of Developmental and Behavioral Pediatrics</i> , 2022, 43, e419-e422.	0.6	3
7	Individual cognitive changes in subjective cognitive decline, mild cognitive impairment and Alzheimer's disease using the reliable change index methodology. <i>Wiener Klinische Wochenschrift</i> , 2021, 133, 1064-1069.	1.0	5
8	Patient satisfaction after breast cancer surgery. <i>Wiener Klinische Wochenschrift</i> , 2021, 133, 6-13.	1.0	11
9	Symptom and performance validation in patients with subjective cognitive decline and mild cognitive impairment. <i>Applied Neuropsychology Adult</i> , 2021, 28, 269-281.	0.7	14
10	Assessing visuo-constructive functions in patients with subjective cognitive decline, mild cognitive impairment and Alzheimer's disease with the Vienna Visuo-Constructional Test 3.0 (VVIT 3.0). <i>Neuropsychiatrie</i> , 2021, 35, 147-155.	1.3	3
11	Basal myokine levels are associated with quality of life and depressed mood in older adults. <i>Psychophysiology</i> , 2021, 58, e13799.	1.2	12
12	Depressive symptoms and olfactory function in patients with subjective cognitive decline, mild cognitive impairment and Alzheimer's disease. <i>Brain Disorders</i> , 2021, 2, 100014.	1.1	1
13	Awareness of Olfactory Dysfunction in Subjective Cognitive Decline, Mild Cognitive Decline, and Alzheimer's Disease. <i>Chemosensory Perception</i> , 2020, 13, 59-70.	0.7	4
14	Transcranial Pulse Stimulation with Ultrasound in Alzheimer's Disease – A New Navigated Focal Brain Therapy. <i>Advanced Science</i> , 2020, 7, 1902583.	5.6	117
15	Comparing a visual and verbal semantic memory test on the effects of gender, age and education as assessed in a cognitively healthy sample. <i>Neuropsychiatrie</i> , 2020, 34, 140-147.	1.3	4
16	Focal Brain Therapy: Transcranial Pulse Stimulation with Ultrasound in Alzheimer's Disease – A New Navigated Focal Brain Therapy (Adv. Sci. 3/2020). <i>Advanced Science</i> , 2020, 7, 2070017.	5.6	4
17	Depression, quality of life, activities of daily living, and subjective memory after deep brain stimulation in Parkinson disease – A reliable change index analysis. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 1698-1705.	1.3	11
18	Identification of odors, faces, cities and naming of objects in patients with subjective cognitive decline, mild cognitive impairment and Alzheimer's disease: a longitudinal study. <i>International Psychogeriatrics</i> , 2019, 31, 537-549.	0.6	22

#	ARTICLE	IF	CITATIONS
19	High intensity endurance training is associated with better quality of life, but not with improved cognitive functions in elderly marathon runners. <i>Scientific Reports</i> , 2019, 9, 4629.	1.6	12
20	Differences regarding the five-factor personality model in patients with subjective cognitive decline and mild cognitive impairment. <i>Neuropsychiatrie</i> , 2019, 33, 35-45.	1.3	9
21	Awareness of olfactory dysfunction in Parkinson's disease. <i>Neuropsychology</i> , 2019, 33, 633-641.	1.0	32
22	Individual cognitive change after DBS-surgery in Parkinson's disease patients using Reliable Change Index Methodology. <i>Neuropsychiatrie</i> , 2018, 32, 149-158.	1.3	9
23	Screening for dementia with the Vienna Visuo-Constructional Test 3.0 screening (VWT 3.0 screening). <i>Neuropsychiatrie</i> , 2018, 32, 196-203.	1.3	5
24	Early dysfunctions of fronto-parietal praxis networks in Parkinson's disease. <i>Brain Imaging and Behavior</i> , 2017, 11, 512-525.	1.1	9
25	Assessment of individual cognitive changes after deep brain stimulation surgery in Parkinson's disease using the Neuropsychological Test Battery Vienna short version. <i>Wiener Klinische Wochenschrift</i> , 2017, 129, 564-571.	1.0	9
26	Semantic memory and depressive symptoms in patients with subjective cognitive decline, mild cognitive impairment, and Alzheimer's disease. <i>International Psychogeriatrics</i> , 2017, 29, 1123-1135.	0.6	17
27	Activities of Daily Living and Depressive Symptoms in Patients with Subjective Cognitive Decline, Mild Cognitive Impairment, and Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2016, 49, 1043-1050.	1.2	48
28	Facial emotion recognition and its relationship to cognition and depressive symptoms in patients with Parkinson's disease. <i>International Psychogeriatrics</i> , 2016, 28, 1165-1179.	0.6	24
29	The impact of depressive symptoms on health-related quality of life in patients with subjective cognitive decline, mild cognitive impairment, and Alzheimer's disease. <i>International Psychogeriatrics</i> , 2016, 28, 2045-2054.	0.6	21
30	Facial emotion recognition in patients with subjective cognitive decline and mild cognitive impairment. <i>International Psychogeriatrics</i> , 2016, 28, 477-485.	0.6	20
31	Self-reported and informant-reported memory functioning and awareness in patients with mild cognitive impairment and Alzheimer's disease. <i>Neuropsychiatrie</i> , 2016, 30, 103-112.	1.3	13
32	rs6295 [C]-Allele Protects Against Depressive Mood in Elderly Endurance Athletes. <i>Journal of Sport and Exercise Psychology</i> , 2015, 37, 637-645.	0.7	8
33	Health-Related Quality of Life in Patients with Subjective Cognitive Decline and Mild Cognitive Impairment and its Relation to Activities of Daily Living. <i>Journal of Alzheimer's Disease</i> , 2015, 47, 479-486.	1.2	67
34	Physical Exercise Counteracts Genetic Susceptibility to Depression. <i>Neuropsychobiology</i> , 2015, 71, 168-175.	0.9	54
35	Finger dexterity deficits in Parkinson's disease and somatosensory cortical dysfunction. <i>Parkinsonism and Related Disorders</i> , 2015, 21, 259-265.	1.1	32
36	Higher level of neuroticism in patients with problems with the sense of smell. <i>Wiener Klinische Wochenschrift</i> , 2015, 127, 303-307.	1.0	7

#	ARTICLE	IF	CITATIONS
37	Visuo-constructional functions in patients with mild cognitive impairment, Alzheimer's disease, and Parkinson's disease. <i>Neuropsychiatrie</i> , 2015, 29, 112-119.	1.3	8
38	Gender-Specific Differences in Cognitive Profiles of Patients with Alzheimer's Disease: Results of the Prospective Dementia Registry Austria (PRODEM-Austria). <i>Journal of Alzheimer's Disease</i> , 2015, 46, 631-637.	1.2	20
39	Awareness of memory deficits in subjective cognitive decline, mild cognitive impairment, Alzheimer's disease and Parkinson's disease. <i>International Psychogeriatrics</i> , 2015, 27, 357-366.	0.6	74
40	Depressive Symptoms are the Main Predictor for Subjective Sleep Quality in Patients with Mild Cognitive Impairment—A Controlled Study. <i>PLoS ONE</i> , 2015, 10, e0128139.	1.1	12
41	A neuropsychological rehabilitation program for patients with Multiple Sclerosis based on the model of the ICF. <i>NeuroRehabilitation</i> , 2014, 35, 519-527.	0.5	19
42	Subjective memory complaints, depressive symptoms and cognition in Parkinson's disease patients. <i>European Journal of Neurology</i> , 2014, 21, 1276.	1.7	50
43	Subjective memory complaints, depressive symptoms and cognition in patients attending a memory outpatient clinic. <i>International Psychogeriatrics</i> , 2014, 26, 463-473.	0.6	53
44	Prevalence of mild cognitive impairment subtypes in patients attending a memory outpatient clinic—comparison of two modes of mild cognitive impairment classification. Results of the Vienna Conversion to Dementia Study. <i>Alzheimer's and Dementia</i> , 2013, 9, 366-376.	0.4	40
45	The "Sense of Coherence" and the coping capacity of patients with Parkinson disease. <i>International Psychogeriatrics</i> , 2012, 24, 1972-1979.	0.6	52
46	Development of a Brief Self-Report Inventory to Measure Olfactory Dysfunction and Quality of Life in Patients with Problems with the Sense of Smell. <i>Chemosensory Perception</i> , 2012, 5, 292-299.	0.7	30
47	Demenzsyndrome. , 2011, , 375-394.		1
48	Klinische Psychologie in der Neurologie. , 2011, , 223-242.		0
49	Gedächtnisstörungen. , 2011, , 541-559.		1
50	Cognitive function in elderly marathon runners: Cross-sectional data from the marathon trial (apsoem). <i>Wiener Klinische Wochenschrift</i> , 2010, 122, 704-716.	1.0	28
51	Neurocognitive training in patients with high-grade glioma: a pilot study. <i>Journal of Neuro-Oncology</i> , 2010, 97, 109-115.	1.4	78
52	FMRI correlates of apraxia in Parkinson's disease patients OFF medication. <i>Experimental Neurology</i> , 2010, 225, 416-422.	2.0	24
53	Odor Identification and Self-reported Olfactory Functioning in Patients with Subtypes of Mild Cognitive Impairment. <i>Clinical Neuropsychologist</i> , 2009, 23, 818-830.	1.5	37
54	Improvement of neurocognitive function after protected carotid artery stenting. <i>Catheterization and Cardiovascular Interventions</i> , 2008, 71, 114-119.	0.7	22

#	ARTICLE	IF	CITATIONS
55	Does modification of olfacto-gustatory stimulation diminish sensory-specific satiety in humans?. <i>Physiology and Behavior</i> , 2006, 87, 469-477.	1.0	26
56	Demenzsyndrome. , 2006, , 327-346.		1
57	Gedächtnisstörungen. , 2006, , 455-473.		0
58	Annual conversion to Alzheimer disease among patients with memory complaints attending an outpatient memory clinic: the influence of amnesic mild cognitive impairment and the predictive value of neuropsychological testing. <i>Wiener Klinische Wochenschrift</i> , 2005, 117, 629-635.	1.0	60
59	Depth of word processing in Alzheimer patients and normal controls: a magnetoencephalographic (MEG) study. <i>Journal of Neural Transmission</i> , 2005, 112, 713-730.	1.4	26
60	Neuropsychological Outcome 6 Months after Unilateral Carotid Stenting. <i>Journal of Clinical and Experimental Neuropsychology</i> , 2005, 27, 859-866.	0.8	39
61	Ambient odors of orange and lavender reduce anxiety and improve mood in a dental office. <i>Physiology and Behavior</i> , 2005, 86, 92-95.	1.0	346
62	Olfaction and face encoding in humans: a magnetoencephalographic study. <i>Cognitive Brain Research</i> , 2003, 15, 105-115.	3.3	14
63	Olfaction and Depth of Word Processing: A Magnetoencephalographic Study. <i>NeuroImage</i> , 2003, 18, 104-116.	2.1	18
64	Evidence of conscious and subconscious olfactory information processing during word encoding: a magnetoencephalographic (MEG) study. <i>Cognitive Brain Research</i> , 2002, 14, 309-316.	3.3	21
65	Ambient odor of orange in a dental office reduces anxiety and improves mood in female patients. <i>Physiology and Behavior</i> , 2000, 71, 83-86.	1.0	251
66	Odor Identification, Consistency of Label Use, Olfactory Threshold and their Relationships to Odor Memory over the Human Lifespan. <i>Chemical Senses</i> , 1999, 24, 337-346.	1.1	150
67	Health-related quality of life (HRQOL), activity of daily living (ADL) and depressive mood disorder in temporal lobe epilepsy patients. <i>Seizure: the Journal of the British Epilepsy Association</i> , 1999, 8, 88-92.	0.9	186
68	Language-related hemispheric asymmetry in healthy subjects and patients with temporal lobe epilepsy as studied by event-related brain potentials and intracarotid amobarbital test. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1998, 108, 274-282.	2.0	23
69	Event-related potentials in patients with temporal lobe epilepsy reveal topography specific lateralization in relation to the side of the epileptic focus. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1998, 108, 567-576.	2.0	7
70	Olfactory Functions in Parkinson's Disease and Alzheimer's Disease. <i>Chemical Senses</i> , 1997, 22, 105-110.	1.1	39
71	Olfactory Prodromal Symptoms and Unilateral Olfactory Dysfunction Are Associated in Patients with Right Mesial Temporal Lobe Epilepsy. <i>Epilepsia</i> , 1997, 38, 1042-1044.	2.6	22
72	Impaired olfactory function in Parkinson's disease. <i>Lancet, The</i> , 1995, 345, 1054-1055.	6.3	36

#	ARTICLE	IF	CITATIONS
73	Long-term Olfactory Functions in Patients with Subjective Cognitive Decline and Mild Cognitive Impairment. <i>Chemosensory Perception</i> , 0, , 1.	0.7	0