

# Ewald Moser

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1570933/publications.pdf>

Version: 2024-02-01

269  
papers

14,718  
citations

16411

64  
h-index

24915

109  
g-index

312  
all docs

312  
docs citations

312  
times ranked

16401  
citing authors

#	ARTICLE	IF	CITATIONS
1	Transitory ischemic attack associated with a rare fenestration of the cervical segment of the internal carotid artery: a case report. <i>Journal of Medical Case Reports</i> , 2022, 16, 13.	0.4	4
2	Technical note: A PET/MR coil with an integrated, orbiting 511 keV transmission source for PET/MR imaging validated in an animal study. <i>Medical Physics</i> , 2022, 49, 2366-2372.	1.6	3
3	Medical Physics and Imaging – A Timely Perspective. <i>Frontiers in Physics</i> , 2021, 9, .	1.0	5
4	3D localized lactate detection in muscle tissue using double-quantum filtered <sup>1</sup> H MRS with adiabatic refocusing pulses at 7 T. <i>Magnetic Resonance in Medicine</i> , 2021, , .	1.9	2
5	Interleaved <sup>31</sup> P MRS/ <sup>1</sup> H ASL for analysis of metabolic and functional heterogeneity along human lower leg muscles at 7T. <i>Magnetic Resonance in Medicine</i> , 2020, 83, 1909-1919.	1.9	20
6	Disparity of time-contrast curves generated by various types of power injectors used in magnetic resonance imaging. <i>Scientific Reports</i> , 2020, 10, 19568.	1.6	0
7	Advanced Analysis of the Water/Fat Distribution in Skeletal Muscle Tissue Using Magnetic Resonance Imaging in Patients With Neuromuscular Disease. <i>Frontiers in Physics</i> , 2020, 8, .	1.0	2
8	Design, Implementation, and Evaluation of a Head and Neck MRI RF Array Integrated with a 511 keV Transmission Source for Attenuation Correction in PET/MR. <i>Sensors</i> , 2019, 19, 3297.	2.1	5
9	Prefrontal networks dynamically related to recovery from major depressive disorder: a longitudinal pharmacological fMRI study. <i>Translational Psychiatry</i> , 2019, 9, 64.	2.4	43
10	Dynamic multivoxel-localized <sup>31</sup> P MRS during plantar flexion exercise with variable knee angle. <i>NMR in Biomedicine</i> , 2018, 31, e3905.	1.6	13
11	Effects of 1,8-cineole and (E)-linalool on Functional Brain Activation in a Working Memory Task. <i>Flavour and Fragrance Journal</i> , 2018, 33, 235-244.	1.2	11
12	Immediate and delayed neuroendocrine responses to social exclusion in males and females. <i>Psychoneuroendocrinology</i> , 2018, 93, 56-64.	1.3	23
13	In vivo MRI of the human finger at 7 T. <i>Magnetic Resonance in Medicine</i> , 2018, 79, 588-592.	1.9	23
14	A head coil system with an integrated orbiting transmission point source mechanism for attenuation correction in PET/MRI. <i>Physics in Medicine and Biology</i> , 2018, 63, 225014.	1.6	12
15	Flexible 23-channel coil array for high-resolution magnetic resonance imaging at 3 Tesla. <i>PLoS ONE</i> , 2018, 13, e0206963.	1.1	24
16	A flexible 12-channel transceiver array of transmission line resonators for 7 T MRI. <i>Journal of Magnetic Resonance</i> , 2018, 296, 47-59.	1.2	13
17	Self-Managed Belief as Part of the “Scientific Method” Part II – Examples From Published Scientific Work. <i>Frontiers in Physics</i> , 2018, 6, .	1.0	0
18	Hybrid Imaging: Instrumentation and Data Processing. <i>Frontiers in Physics</i> , 2018, 6, .	1.0	30

#	ARTICLE	IF	CITATIONS
19	Pros and cons of ultra-high-field MRI/MRS for human application. Progress in Nuclear Magnetic Resonance Spectroscopy, 2018, 109, 1-50.	3.9	331
20	Magnetic resonance imaging T1- and T2-mapping to assess renal structure and function: a systematic review and statement paper. Nephrology Dialysis Transplantation, 2018, 33, ii41-ii50.	0.4	75
21	Handy magnetic resonance coils. Nature Biomedical Engineering, 2018, 2, 557-558.	11.6	8
22	A Quantitative Comparison of Clinically Employed Parameters in the Assessment of Acute Cerebral Ischemia Using Dynamic Susceptibility Contrast Magnetic Resonance Imaging. Frontiers in Physiology, 2018, 9, 1945.	1.3	2
23	Normalised time-to-peak-distribution curves correlate with cerebral white matter hyperintensities â€œ Could this improve early diagnosis?. Journal of Cerebral Blood Flow and Metabolism, 2017, 37, 444-455.	2.4	13
24	High-sensitivity TMS/fMRI of the Human Motor Cortex Using a Dedicated Multichannel MR Coil. NeuroImage, 2017, 150, 262-269.	2.1	43
25	Impact of self-esteem and sex on stress reactions. Scientific Reports, 2017, 7, 17210.	1.6	50
26	Interleaved multivoxel <sup>31</sup> P MR spectroscopy. Magnetic Resonance in Medicine, 2017, 77, 921-927.	1.9	16
27	Ultra-High Field NMR and MRIâ€”The Role of Magnet Technology to Increase Sensitivity and Specificity. Frontiers in Physics, 2017, 5, .	1.0	62
28	Editorial: In Vivo Magnetic Resonance at Ultra High Field. Frontiers in Physics, 2017, 5, .	1.0	1
29	Dynamic PCr and pH imaging of human calf muscles during exercise and recovery using <sup>31</sup> P gradientâ€œEcho MRI at 7 Tesla. Magnetic Resonance in Medicine, 2016, 75, 2324-2331.	1.9	31
30	Skeletal muscle ATP synthesis and cellular H <sup>+</sup> handling measured by localized <sup>31</sup> P-MRS during exercise and recovery. Scientific Reports, 2016, 6, 32037.	1.6	33
31	Multi-turn multi-gap transmission line resonators â€œ Concept, design and first implementation at 4.7 T and 7 T. Journal of Magnetic Resonance, 2016, 273, 65-72.	1.2	18
32	Sex differences in the functional connectivity of the amygdalae in association with cortisol. NeuroImage, 2016, 134, 410-423.	2.1	62
33	fMRI of Emotion. Neuromethods, 2016, , 451-494.	0.2	1
34	Simultaneous and interleaved acquisition of <sup>1</sup> H NMR signals from different nuclei with a clinical MRI scanner. Magnetic Resonance in Medicine, 2016, 76, 1636-1641.	1.9	29
35	Simultaneous and interleaved acquisition of NMR signals from different nuclei with a clinical MRI scanner. Magnetic Resonance in Medicine, 2016, 76, spcone-spcone.	1.9	1
36	Oppositional COMT Val158Met effects on resting state functional connectivity in adolescents and adults. Brain Structure and Function, 2016, 221, 103-114.	1.2	31

#	ARTICLE	IF	CITATIONS
37	fMRI measurements of amygdala activation are confounded by stimulus correlated signal fluctuation in nearby veins draining distant brain regions. <i>Scientific Reports</i> , 2015, 5, 10499.	1.6	104
38	Novel inductive decoupling technique for flexible transceiver arrays of monolithic transmission line resonators. <i>Magnetic Resonance in Medicine</i> , 2015, 73, 1669-1681.	1.9	26
39	A form-fitted three channel <sup>31</sup> P, two channel <sup>1</sup> H transceiver coil array for calf muscle studies at 7 T. <i>Magnetic Resonance in Medicine</i> , 2015, 73, 2376-2389.	1.9	40
40	Power balance and loss mechanism analysis in RF transmit coil arrays. <i>Magnetic Resonance in Medicine</i> , 2015, 74, 1165-1176.	1.9	33
41	A functional polymorphism in the prodynorphin gene affects cognitive flexibility and brain activation during reversal learning. <i>Frontiers in Behavioral Neuroscience</i> , 2015, 9, 172.	1.0	13
42	Identification of Voxels Confounded by Venous Signals Using Resting-State fMRI Functional Connectivity Graph Community Identification. <i>Frontiers in Neuroscience</i> , 2015, 9, 472.	1.4	13
43	Localized semi-LASER dynamic <sup>31</sup> P magnetic resonance spectroscopy of the soleus during and following exercise at 7T. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2015, 28, 493-501.	1.1	23
44	Reduced default mode network suppression during a working memory task in remitted major depression. <i>Journal of Psychiatric Research</i> , 2015, 64, 9-18.	1.5	99
45	fMRI correlates of different components of Braille reading by the blind. <i>Neurology Psychiatry and Brain Research</i> , 2015, 21, 137-145.	2.0	12
46	In vivo MR imaging of the human skin at subnanoliter resolution using a superconducting surface coil at 1.5 tesla. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 496-504.	1.9	21
47	Dynamic ASL and T2*-weighted MRI in exercising calf muscle at 7 T: A feasibility study. <i>Magnetic Resonance in Medicine</i> , 2015, 73, 1190-1195.	1.9	39
48	New ultra-thin multichannel receive coil for concurrent TMS/fMRI experiments. <i>Brain Stimulation</i> , 2015, 8, 426-427.	0.7	0
49	A novel coil array for combined TMS/fMRI experiments at 3 T. <i>Magnetic Resonance in Medicine</i> , 2015, 74, 1492-1501.	1.9	46
50	Disrupted Effective Connectivity Between the Amygdala and Orbitofrontal Cortex in Social Anxiety Disorder During Emotion Discrimination Revealed by Dynamic Causal Modeling for fMRI. <i>Cerebral Cortex</i> , 2015, 25, 895-903.	1.6	139
51	Big Data Approaches for the Analysis of Large-Scale fMRI Data Using Apache Spark and GPU Processing: A Demonstration on Resting-State fMRI Data from the Human Connectome Project. <i>Frontiers in Neuroscience</i> , 2015, 9, 492.	1.4	48
52	Improved Quantification of Cerebral Hemodynamics Using Individualized Time Thresholds for Assessment of Peak Enhancement Parameters Derived from Dynamic Susceptibility Contrast Enhanced Magnetic Resonance Imaging. <i>PLoS ONE</i> , 2014, 9, e114999.	1.1	2
53	Scanning fast and slow: current limitations of 3 Tesla functional MRI and future potential. <i>Frontiers in Physics</i> , 2014, 2, 00001.	1.0	20
54	On the generalizability of resting-state fMRI machine learning classifiers. <i>Frontiers in Human Neuroscience</i> , 2014, 8, 502.	1.0	9

#	ARTICLE	IF	CITATIONS
55	Technical Note: Evaluation of the Uncertainties in (Choline+Creatine)/Citrate Ratios Measured by Proton MR Spectroscopic Imaging in Patients Suspicious for Prostate Cancer. <i>RoFo Fortschritte Auf Dem Gebiet Der Rontgenstrahlen Und Der Bildgebenden Verfahren</i> , 2014, 186, 698-702.	0.7	2
56	Exercising calf muscle changes correlate with pH, PCr recovery and maximum oxidative phosphorylation. <i>NMR in Biomedicine</i> , 2014, 27, 553-560.	1.6	31
57	Lower Fasting Muscle Mitochondrial Activity Relates to Hepatic Steatosis in Humans. <i>Diabetes Care</i> , 2014, 37, 468-474.	4.3	26
58	Additive Gene-Environment Effects on Hippocampal Structure in Healthy Humans. <i>Journal of Neuroscience</i> , 2014, 34, 9917-9926.	1.7	59
59	A Genetic Polymorphism of the Endogenous Opioid Dynorphin Modulates Monetary Reward Anticipation in the Corticostriatal Loop. <i>PLoS ONE</i> , 2014, 9, e89954.	1.1	13
60	Platelet Serotonin Transporter Function Predicts Default-Mode Network Activity. <i>PLoS ONE</i> , 2014, 9, e92543.	1.1	19
61	The Spectral Diversity of Resting-State Fluctuations in the Human Brain. <i>PLoS ONE</i> , 2014, 9, e93375.	1.1	76
62	High-resolution functional MRI of the human amygdala at 7T. <i>European Journal of Radiology</i> , 2013, 82, 728-733.	1.2	71
63	RESCALE: Voxel-specific task-fMRI scaling using resting state fluctuation amplitude. <i>NeuroImage</i> , 2013, 70, 80-88.	2.1	34
64	MR/PET or PET/MRI: does it matter?. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2013, 26, 1-4.	1.1	11
65	Automatic model-based analysis of skeletal muscle BOLD-fMRI in reactive hyperemia. <i>Journal of Magnetic Resonance Imaging</i> , 2013, 38, 963-969.	1.9	12
66	Discontinuous Patterns of Brain Activation in the Psychotherapy Process of Obsessive-Compulsive Disorder: Converging Results from Repeated fMRI and Daily Self-Reports. <i>PLoS ONE</i> , 2013, 8, e71863.	1.1	53
67	Grand challenges in biomedical physics. <i>Frontiers in Physics</i> , 2013, 1, .	1.0	10
68	Beyond Noise: Using Temporal ICA to Extract Meaningful Information from High-Frequency fMRI Signal Fluctuations during Rest. <i>Frontiers in Human Neuroscience</i> , 2013, 7, 168.	1.0	149
69	Effects of individual glucose levels on the neuronal correlates of emotions. <i>Frontiers in Human Neuroscience</i> , 2013, 7, 212.	1.0	11
70	P.1.e.017 Consistency of resting-state networks in a multi-centre dataset. <i>European Neuropsychopharmacology</i> , 2012, 22, S201-S202.	0.3	0
71	A highly parallelized framework for computationally intensive MR data analysis. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2012, 25, 313-320.	1.1	14
72	Culture but not gender modulates amygdala activation during explicit emotion recognition. <i>BMC Neuroscience</i> , 2012, 13, 54.	0.8	35

#	ARTICLE	IF	CITATIONS
73	Heme arginate improves reperfusion patterns after ischemia: a randomized, placebo-controlled trial in healthy male subjects. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2012, 14, 35.	1.6	22
74	The Human Factor: Behavioral and Neural Correlates of Humanized Perception in Moral Decision Making. <i>PLoS ONE</i> , 2012, 7, e47698.	1.1	39
75	Fully exploratory network independent component analysis of the 1000 functional connectomes database. <i>Frontiers in Human Neuroscience</i> , 2012, 6, 301.	1.0	55
76	Comparison of measuring energy metabolism by different <sup>31</sup> P magnetic resonance spectroscopy techniques in resting, ischemic, and exercising muscle. <i>Magnetic Resonance in Medicine</i> , 2012, 67, 898-905.	1.9	35
77	Comparing localized and nonlocalized dynamic <sup>31</sup> P magnetic resonance spectroscopy in exercising muscle at 7T. <i>Magnetic Resonance in Medicine</i> , 2012, 68, 1713-1723.	1.9	55
78	A Single Nucleotide Polymorphism Associates With the Response of Muscle ATP Synthesis to Long-Term Exercise Training in Relatives of Type 2 Diabetic Humans. <i>Diabetes Care</i> , 2012, 35, 350-357.	4.3	25
79	No amygdala attenuation in schizophrenic patients treated with atypical antipsychotics. <i>Psychiatry Research - Neuroimaging</i> , 2012, 202, 168-171.	0.9	10
80	7â€ MRâ€”from research to clinical applications?. <i>NMR in Biomedicine</i> , 2012, 25, 695-716.	1.6	168
81	Increased Neural Habituation in the Amygdala and Orbitofrontal Cortex in Social Anxiety Disorder Revealed by fMRI. <i>PLoS ONE</i> , 2012, 7, e50050.	1.1	82
82	Orbitofrontal hyperactivity in social anxiety disorder patients: An fmri study. <i>European Psychiatry</i> , 2011, 26, 179-179.	0.1	1
83	Reduced connectivity in the uncinate fiber tract between the frontal cortex and limbic subcortical areas in social phobia. <i>European Psychiatry</i> , 2011, 26, 182-182.	0.1	5
84	Biological alterations during remission of major depressive disorder. <i>European Psychiatry</i> , 2011, 26, 633-633.	0.1	0
85	Peripheral serotonin uptake is related to neural activation in the cingulate cortex. <i>European Psychiatry</i> , 2011, 26, 684-684.	0.1	0
86	Increased functional coupling between basal ganglia and cingulate and prefrontal cortex during resting state conditions in remitted major depressive disorder. <i>European Psychiatry</i> , 2011, 26, 915-915.	0.1	0
87	Increased coupling of resting state activity between amygdala and cortical emotion processing regions in remitted major depressive disorder. <i>European Psychiatry</i> , 2011, 26, 931-931.	0.1	0
88	Remitted major depression is related to increased functional coupling between ventral striatum and cortical regions in resting state fMRI. <i>European Psychiatry</i> , 2011, 26, 948-948.	0.1	1
89	Reduced resting-state functional connectivity between amygdala and orbitofrontal cortex in social anxiety disorder. <i>NeuroImage</i> , 2011, 56, 881-889.	2.1	353
90	Slice-timing effects and their correction in functional MRI. <i>NeuroImage</i> , 2011, 58, 588-594.	2.1	309

#	ARTICLE	IF	CITATIONS
91	Model-free fMRI group analysis using FENICA. <i>NeuroImage</i> , 2011, 55, 185-193.	2.1	35
92	Effect of ischemic preconditioning in skeletal muscle measured by functional magnetic resonance imaging and spectroscopy: a randomized crossover trial. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2011, 13, 32.	1.6	68
93	P.1.033 Serotonin uptake in platelets predicts neural activation in the cingulate cortex. <i>European Neuropsychopharmacology</i> , 2011, 21, S27.	0.3	0
94	P.2.c.020 Synthesising evidence in clinical psychopharmacology and beyond: a re-examination of published meta-analyses. <i>European Neuropsychopharmacology</i> , 2011, 21, S394-S395.	0.3	0
95	Semi-LASER localized dynamic <sup>31</sup> P magnetic resonance spectroscopy in exercising muscle at ultra-high magnetic field. <i>Magnetic Resonance in Medicine</i> , 2011, 65, 1207-1215.	1.9	39
96	Magnetic resonance microimaging of human skin vasculature in vivo at 3 Tesla. <i>Magnetic Resonance in Medicine</i> , 2011, 65, 1718-1723.	1.9	19
97	Body and Liver Fat Mass Rather Than Muscle Mitochondrial Function Determine Glucose Metabolism in Women With a History of Gestational Diabetes Mellitus. <i>Diabetes Care</i> , 2011, 34, 430-436.	4.3	42
98	Liver ATP Synthesis Is Lower and Relates to Insulin Sensitivity in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2011, 34, 448-453.	4.3	177
99	Group ICA of resting-state data: a comparison. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2010, 23, 317-325.	1.1	41
100	In vivo MR imaging of brain networks: illusion or revolution?. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2010, 23, 275-277.	1.1	3
101	Multiple serial picture presentation with millisecond resolution using a three-way LC-shutter-tachistoscope. <i>Journal of Neuroscience Methods</i> , 2010, 187, 235-242.	1.3	6
102	Fully exploratory network ICA (FENICA) on resting-state fMRI data. <i>Journal of Neuroscience Methods</i> , 2010, 192, 207-213.	1.3	65
103	Windows on the Human Body – in Vivo High-Field Magnetic Resonance Research and Applications in Medicine and Psychology. <i>Sensors</i> , 2010, 10, 5724-5757.	2.1	12
104	Modulation of hypothalamus and amygdalar activation levels with stimulus valence. <i>NeuroImage</i> , 2010, 51, 324-328.	2.1	26
105	Inducing perceptual unawareness: Tachistoscopic stimulus presentation versus visual masking. <i>International Journal of Psychophysiology</i> , 2010, 77, 324-325.	0.5	1
106	Non-invasive assessment of hepatic fat accumulation in chronic hepatitis C by 1H magnetic resonance spectroscopy. <i>European Journal of Radiology</i> , 2010, 74, e60-e66.	1.2	50
107	P.1.028 Altered functional connectivity of the amygdala in social anxiety disorder. <i>European Neuropsychopharmacology</i> , 2010, 20, S24-S25.	0.3	0
108	P.1.e.011 Fronto-limbic interaction of working memory and emotion in the medial prefrontal cortex. <i>European Neuropsychopharmacology</i> , 2010, 20, S294.	0.3	0

#	ARTICLE	IF	CITATIONS
109	Area-specific modulation of neural activation comparing escitalopram and citalopram revealed by pharmacof-MRI: A randomized cross-over study. <i>NeuroImage</i> , 2010, 49, 1161-1170.	2.1	111
110	Multi-subject analyses with dynamic causal modeling. <i>NeuroImage</i> , 2010, 49, 3065-3074.	2.1	61
111	Ultra-high-field magnetic resonance: Why and when?. <i>World Journal of Radiology</i> , 2010, 2, 37.	0.5	35
112	Amygdala activation during recognition of emotions in a foreign ethnic group is associated with duration of stay. <i>Social Neuroscience</i> , 2009, 4, 294-307.	0.7	50
113	Short-Term Exercise Training Does Not Stimulate Skeletal Muscle ATP Synthesis in Relatives of Humans With Type 2 Diabetes. <i>Diabetes</i> , 2009, 58, 1333-1341.	0.3	62
114	Diffusion-weighted MR for Differentiation of Breast Lesions at 3.0 T: How Does Selection of Diffusion Protocols Affect Diagnosis?. <i>Radiology</i> , 2009, 253, 341-351.	3.6	262
115	Amygdala activity to fear and anger in healthy young males is associated with testosterone. <i>Psychoneuroendocrinology</i> , 2009, 34, 687-693.	1.3	166
116	A resting state network in the motor control circuit of the basal ganglia. <i>BMC Neuroscience</i> , 2009, 10, 137.	0.8	134
117	General and specific responsiveness of the amygdala during explicit emotion recognition in females and males. <i>BMC Neuroscience</i> , 2009, 10, 91.	0.8	76
118	Abnormal hepatic energy homeostasis in type 2 diabetes. <i>Hepatology</i> , 2009, 50, 1079-1086.	3.6	166
119	Assessment of $T_1\rho$ relaxation times in the human calf muscle: A comparison between 3 T and 7 T in vivo. <i>Magnetic Resonance in Medicine</i> , 2009, 62, 574-582.	1.9	118
120	Magnetic resonance imaging methodology. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2009, 36, 30-41.	3.3	40
121	Correlations and anticorrelations in resting-state functional connectivity MRI: A quantitative comparison of preprocessing strategies. <i>NeuroImage</i> , 2009, 47, 1408-1416.	2.1	745
122	fMRI of Emotion. <i>NeuroMethods</i> , 2009, , 411-456.	0.2	8
123	Estimating Activation of Denervated, Degenerated Muscle after Functional Electrical Stimulation with Magnetic Resonance Imaging. <i>IFMBE Proceedings</i> , 2009, , 191-194.	0.2	0
124	Brain Activity Movie Functional MRI with Ultra-High Temporal Resolution at 7 Tesla. <i>IFMBE Proceedings</i> , 2009, , 192-194.	0.2	1
125	Effects of functional electrical stimulation in denervated thigh muscles of paraplegic patients mapped with T2 imaging. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2008, 21, 219-226.	1.1	13
126	The impact of EPI voxel size on SNR and BOLD sensitivity in the anterior medio-temporal lobe: a comparative group study of deactivation of the Default Mode. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2008, 21, 279-290.	1.1	45



#	ARTICLE	IF	CITATIONS
127	Re-establishment of normal blood flow is mandatory to restore intramuscular high energy phosphate levels after transient ischemia. <i>BMC Pharmacology</i> , 2008, 8, .	0.4	0
128	Three-dimensional high-resolution magnetic resonance spectroscopic imaging for absolute quantification of <sup>31</sup> P metabolites in human liver. <i>Magnetic Resonance in Medicine</i> , 2008, 60, 796-802.	1.9	47
129	Quantitative ATP synthesis in human liver measured by localized <sup>31</sup> P spectroscopy using the magnetization transfer experiment. <i>NMR in Biomedicine</i> , 2008, 21, 437-443.	1.6	61
130	Time-resolved analysis of fMRI signal changes using Brain Activation Movies. <i>Journal of Neuroscience Methods</i> , 2008, 169, 222-230.	1.3	16
131	Functional Electrical Stimulation of Long-term Denervated, Degenerated Human Skeletal Muscle: Estimating Activation Using T2-Parameter Magnetic Resonance Imaging Methods. <i>Artificial Organs</i> , 2008, 32, 604-608.	1.0	16
132	Facial emotion recognition and amygdala activation are associated with menstrual cycle phase. <i>Psychoneuroendocrinology</i> , 2008, 33, 1031-1040.	1.3	156
133	Altered reward processing in the nucleus accumbens and mesial prefrontal cortex of patients with posttraumatic stress disorder. <i>Neuropsychologia</i> , 2008, 46, 2836-2844.	0.7	169
134	Emotion recognition accuracy in healthy young females is associated with cycle phase. <i>Hormones and Behavior</i> , 2008, 53, 90-95.	1.0	160
135	Metabolic changes in the normal ageing brain: Consistent findings from short and long echo time proton spectroscopy. <i>European Journal of Radiology</i> , 2008, 68, 320-327.	1.2	76
136	The suppressive influence of SMA on M1 in motor imagery revealed by fMRI and dynamic causal modeling. <i>NeuroImage</i> , 2008, 40, 828-837.	2.1	219
137	Multimodal imaging of human early visual cortex by combining functional and molecular measurements with fMRI and PET. <i>NeuroImage</i> , 2008, 41, 204-211.	2.1	32
138	The spatial resolution in dosimetry with normoxic polymer gels investigated with the dose modulation transfer approach. <i>Medical Physics</i> , 2008, 35, 1756-1769.	1.6	18
139	Neurological and brain MRS findings in patients with Gaucher disease type 1. <i>Molecular Genetics and Metabolism</i> , 2007, 91, 390-395.	0.5	16
140	Treatment of cognition and affect in schizophrenia. <i>European Psychiatry</i> , 2007, 22, S13.	0.1	0
141	Diffusion tensor imaging and optimized fiber tracking in glioma patients: Histopathologic evaluation of tumor-invaded white matter structures. <i>NeuroImage</i> , 2007, 34, 949-956.	2.1	117
142	The functional role of dorso-lateral premotor cortex during mental rotation. <i>NeuroImage</i> , 2007, 36, 1374-1386.	2.1	69
143	Imaging the changing role of feedback during learning in decision-making. <i>NeuroImage</i> , 2007, 37, 1474-1486.	2.1	24
144	Absolute quantification of phosphorus metabolite concentrations in human muscle <i>in vivo</i> by <sup>31</sup> P MRS: a quantitative review. <i>NMR in Biomedicine</i> , 2007, 20, 555-565.	1.6	256

#	ARTICLE	IF	CITATIONS
145	Direct noninvasive quantification of lactate and high energy phosphates simultaneously in exercising human skeletal muscle by localized magnetic resonance spectroscopy. <i>Magnetic Resonance in Medicine</i> , 2007, 57, 654-660.	1.9	39
146	Amygdala activation at 3T in response to human and avatar facial expressions of emotions. <i>Journal of Neuroscience Methods</i> , 2007, 161, 126-133.	1.3	110
147	Decrease of NAA with aging outside the seizure focus in mesial temporal lobe epilepsy – A proton-MRS study at 3T. <i>Brain Research</i> , 2007, 1179, 131-139.	1.1	15
148	Amygdala activation and facial expressions: Explicit emotion discrimination versus implicit emotion processing. <i>Neuropsychologia</i> , 2007, 45, 2369-2377.	0.7	171
149	Changes in fiber integrity, diffusivity, and metabolism of the pyramidal tract adjacent to gliomas: a quantitative diffusion tensor fiber tracking and MR spectroscopic imaging study. <i>American Journal of Neuroradiology</i> , 2007, 28, 462-9.	1.2	66
150	The selection of intended actions and the observation of others' actions: A time-resolved fMRI study. <i>NeuroImage</i> , 2006, 29, 1294-1302.	2.1	123
151	EEG reveals the effect of fMRI scanner noise on noise-sensitive subjects. <i>NeuroImage</i> , 2006, 31, 332-341.	2.1	28
152	Basic investigations on the performance of a normoxic polymer gel with tetrakis-hydroxy-methyl-phosphonium chloride as an oxygen scavenger: Reproducibility, accuracy, stability, and dose rate dependence. <i>Medical Physics</i> , 2006, 33, 2506-2518.	1.6	37
153	<sup>1</sup> H magnetic resonance spectroscopy at 3 T in cryptogenic and mesial temporal lobe epilepsy. <i>NMR in Biomedicine</i> , 2006, 19, 544-553.	1.6	39
154	Preoperative Grading of Gliomas by Using Metabolite Quantification with High-Spatial-Resolution Proton MR Spectroscopic Imaging. <i>Radiology</i> , 2006, 238, 958-969.	3.6	168
155	Gliomas: Histopathologic Evaluation of Changes in Directionality and Magnitude of Water Diffusion at Diffusion-Tensor MR Imaging. <i>Radiology</i> , 2006, 240, 803-810.	3.6	181
156	Proton Magnetic Resonance Spectroscopic Imaging Integrated into Image-guided Surgery: Correlation to Standard Magnetic Resonance Imaging and Tumor Cell Density. <i>Operative Neurosurgery</i> , 2005, 56, ONS-291-ONS-298.	0.4	65
157	Dynamic interleaved <sup>1</sup> H/ <sup>31</sup> P STEAM MRS at 3 Tesla using a pneumatic force-controlled plantar flexion exercise rig. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2005, 18, 257-262.	1.1	39
158	fMRI of the Emotions: Towards an Improved Understanding of Amygdala Function. <i>Current Medical Imaging</i> , 2005, 1, 115-129.	0.4	14
159	Premovement activity of the pre-supplementary motor area and the readiness for action: Studies of time-resolved event-related functional MRI. <i>Human Movement Science</i> , 2005, 24, 644-656.	0.6	141
160	Proton magnetic resonance spectroscopic imaging in brain tumor diagnosis. <i>Neurosurgery Clinics of North America</i> , 2005, 16, 101-114.	0.8	32
161	Diagnostic value of MRI in comparison to scintigraphy, PET, MS-CT and PET/CT for the detection of metastases of bone. <i>European Journal of Radiology</i> , 2005, 55, 41-55.	1.2	174
162	Integration of biochemical images of a tumor into frameless stereotaxy achieved using a magnetic resonance imaging/magnetic resonance spectroscopy hybrid data set. <i>Journal of Neurosurgery</i> , 2004, 101, 287-294.	0.9	63

#	ARTICLE	IF	CITATIONS
163	A quantitative comparison of functional MRI cluster analysis. <i>Artificial Intelligence in Medicine</i> , 2004, 31, 57-71.	3.8	84
164	High resolution MR based polymer dosimetry versus film densitometry: a systematic study based on the modulation transfer function approach. <i>Physics in Medicine and Biology</i> , 2004, 49, 4087-4108.	1.6	24
165	<sup>1</sup> H NMR relaxation times of skeletal muscle metabolites at 3 T. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2004, 16, 155-159.	1.1	75
166	Robust field map generation using a triple-echo acquisition. <i>Journal of Magnetic Resonance Imaging</i> , 2004, 20, 730-734.	1.9	59
167	Optimized 3 T EPI of the amygdalae. <i>NeuroImage</i> , 2004, 22, 203-210.	2.1	125
168	Wavelet-based multifractal analysis of fMRI time series. <i>NeuroImage</i> , 2004, 22, 1195-1202.	2.1	89
169	Improved delineation of brain tumors: an automated method for segmentation based on pathologic changes of <sup>1</sup> H-MRSI metabolites in gliomas. <i>NeuroImage</i> , 2004, 23, 454-461.	2.1	118
170	The dose modulation transfer function concept (DMTF) in polymer dosimetry for quality control in fine photon fields. <i>Journal of Physics: Conference Series</i> , 2004, 3, 200-204.	0.3	0
171	High resolution dosimetric imaging of polymer gels prepared under normoxic conditions: Stereotactic radiation with a 16 mm collimator. <i>Journal of Physics: Conference Series</i> , 2004, 3, 272-275.	0.3	5
172	Fuzzy cluster analysis of high-field functional MRI data. <i>Artificial Intelligence in Medicine</i> , 2003, 29, 203-223.	3.8	40
173	Automated unwrapping of MR phase images applied to BOLD MR-venography at 3 Tesla. <i>Journal of Magnetic Resonance Imaging</i> , 2003, 18, 175-180.	1.9	98
174	High-resolution 3D proton spectroscopic imaging of the human brain at 3 T: SNR issues and application for anatomy-matched voxel sizes. <i>Magnetic Resonance in Medicine</i> , 2003, 49, 299-306.	1.9	61
175	Relaxation times of <sup>31</sup> P-metabolites in human calf muscle at 3 T. <i>Magnetic Resonance in Medicine</i> , 2003, 49, 620-625.	1.9	47
176	Scaling laws and persistence in human brain activity. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2003, 326, 511-521.	1.2	53
177	The preparation and readiness for voluntary movement: a high-field event-related fMRI study of the Bereitschafts-BOLD response. <i>NeuroImage</i> , 2003, 20, 404-412.	2.1	211
178	Human motor cortex activity during mental rotation. <i>NeuroImage</i> , 2003, 20, 225-232.	2.1	51
179	Degree of Hypomyelination and Magnetic Resonance Spectroscopy Findings in Patients with Pelizaeus Merzbacher Phenotype. <i>Neuropediatrics</i> , 2003, 34, 127-136.	0.3	44
180	Title is missing!. <i>Investigative Radiology</i> , 2003, 38, 467-472.	3.5	1

#	ARTICLE	IF	CITATIONS
181	Title is missing!. Investigative Radiology, 2003, 38, 403-408.	3.5	10
182	Title is missing!. Investigative Radiology, 2003, 38, 460-466.	3.5	1
183	Title is missing!. Investigative Radiology, 2003, 38, 452-459.	3.5	2
184	Title is missing!. Investigative Radiology, 2003, 38, 375-376.	3.5	3
185	Quantification of Metabolic Differences in the Frontal Brain of Depressive Patients and Controls Obtained by 1H-MRS at 3 Tesla. Investigative Radiology, 2003, 38, 403-408.	3.5	136
186	High-Resolution Diffusivity Imaging at 3.0 T for the Detection of Degenerative Changes. Investigative Radiology, 2003, 38, 460-466.	3.5	21
187	Bone Homogeneity Factor. Investigative Radiology, 2003, 38, 467-472.	3.5	10
188	Diffusivity- and T2 Imaging at 3 Tesla for the Detection of Degenerative Changes in Human-Excised Tissue with High Resolution. Investigative Radiology, 2003, 38, 452-459.	3.5	12
189	3.0 Tesla MR Systems. Investigative Radiology, 2003, 38, 375-376.	3.5	8
190	Effects of Oral Creatine Supplementation in a Patient with MELAS Phenotype and Associated Nephropathy. Neuropediatrics, 2002, 33, 157-161.	0.3	52
191	The Preparation and Execution of Self-Initiated and Externally-Triggered Movement: A Study of Event-Related fMRI. NeuroImage, 2002, 15, 373-385.	2.1	516
192	Consistency of inter-trial activation using single-trial fMRI: assessment of regional differences. Cognitive Brain Research, 2002, 13, 129-138.	3.3	13
193	Comparison of multi-echo spiral and echo planar imaging in functional MRI. Magnetic Resonance Imaging, 2002, 20, 359-364.	1.0	10
194	On the origin of respiratory artifacts in BOLD-EPI of the human brain. Magnetic Resonance Imaging, 2002, 20, 575-582.	1.0	149
195	Finger Somatotopy in Human Motor Cortex. NeuroImage, 2001, 13, 1016-1026.	2.1	132
196	Evidence for Premotor Cortex Activity during Dynamic Visuospatial Imagery from Single-Trial Functional Magnetic Resonance Imaging and Event-Related Slow Cortical Potentials. NeuroImage, 2001, 14, 268-283.	2.1	173
197	Comparative detectability of bone metastases and impact on therapy of magnetic resonance imaging and bone scintigraphy in patients with breast cancer. European Journal of Radiology, 2001, 40, 16-23.	1.2	54
198	High resolution polymer gel dosimetry by parameter selective MR-microimaging on a whole body scanner at 3 T. Medical Physics, 2001, 28, 833-843.	1.6	41

#	ARTICLE	IF	CITATIONS
199	Spatial Distribution of Prostate Cancers Undetected on Initial Needle Biopsies. <i>European Urology</i> , 2001, 39, 662-668.	0.9	32
200	In vivo magnetic resonance micro-imaging of the human toe at 3 tesla. <i>Magnetic Resonance Imaging</i> , 2001, 19, 1235-1238.	1.0	11
201	Autocorrelation analysis of bone structure. <i>Journal of Magnetic Resonance Imaging</i> , 2001, 14, 87-93.	1.9	36
202	High-resolution blood flow velocity measurements in the human finger. <i>Magnetic Resonance in Medicine</i> , 2001, 45, 716-719.	1.9	121
203	Proton T1 and T2 relaxation times of human brain metabolites at 3 Tesla. <i>NMR in Biomedicine</i> , 2001, 14, 325-331.	1.6	255
204	Characterization of BOLD activation in multi-echo fMRI data using fuzzy cluster analysis and a comparison with quantitative modeling. <i>NMR in Biomedicine</i> , 2001, 14, 484-489.	1.6	12
205	Low-Power Water Suppression by Hyperbolic Secant Pulses with Controlled Offsets and Delays (WASHCODE). <i>Journal of Magnetic Resonance</i> , 2001, 152, 168-178.	1.2	19
206	Co-registration of EEG and MRI data using matching of spline interpolated and MRI-segmented reconstructions of the scalp surface. <i>Brain Topography</i> , 2001, 14, 93-100.	0.8	31
207	Multivoxel 3D proton spectroscopy in the brain at 1.5 versus 3.0 T: signal-to-noise ratio and resolution comparison. <i>American Journal of Neuroradiology</i> , 2001, 22, 1727-31.	1.2	102
208	High-Resolution MR Venography at 3.0 Tesla. <i>Journal of Computer Assisted Tomography</i> , 2000, 24, 949-957.	0.5	190
209	Very short echo time proton MR spectroscopy of human brain with a standard transmit/receive surface coil. <i>Magnetic Resonance in Medicine</i> , 2000, 44, 964-967.	1.9	18
210	Spatial resolution in echo planar imaging: shifting the acquisition window in k-space. <i>Magnetic Resonance Imaging</i> , 2000, 18, 825-834.	1.0	16
211	Wavelet domain de-noising of time-courses in MR image sequences. <i>Magnetic Resonance Imaging</i> , 2000, 18, 1129-1134.	1.0	36
212	A wavelet-based method for improving signal-to-noise ratio and contrast in MR images. <i>Magnetic Resonance Imaging</i> , 2000, 18, 169-180.	1.0	85
213	Improvement of presurgical patient evaluation by generation of functional magnetic resonance risk maps. <i>Neuroscience Letters</i> , 2000, 290, 13-16.	1.0	48
214	P02.188 1H-Magnetic resonance spectroscopy at 3.0 tesla reveals reduced n-acetyl aspartate, choline and myo-inositol levels in depression. <i>European Psychiatry</i> , 2000, 15, 372s-372s.	0.1	0
215	Rapid impairment of skeletal muscle glucose transport/phosphorylation by free fatty acids in humans. <i>Diabetes</i> , 1999, 48, 358-364.	0.3	175
216	The use of single event $\text{AE}'\text{MRI}$ and fuzzy clustering analysis to examine haemodynamic response timecourses in supplementary motor and primary motor cortical areas.. <i>Biomedizinische Technik</i> , 1999, 44, 116-119.	0.9	5

#	ARTICLE	IF	CITATIONS
217	Quantification of Supplementary Motor Area activation with movements of different practice levels using Functional Magnetic Resonance Imaging at 3T. Biomedizinische Technik, 1999, 44, 120-122.	0.9	0
218	High-resolution, multiple gradient-echo functional MRI at 1.5 T. Magnetic Resonance Imaging, 1999, 17, 321-329.	1.0	54
219	A hierarchical clustering method for analyzing functional MR images. Magnetic Resonance Imaging, 1999, 17, 817-826.	1.0	91
220	Functional MRI of the human motor cortex using single-shot, multiple gradient-echo spiral imaging. Magnetic Resonance Imaging, 1999, 17, 1239-1243.	1.0	35
221	Fractal Analysis: An Objective Method for Identifying Atypical Nuclei in Dysplastic Lesions of the Cervix Uteri. Gynecologic Oncology, 1999, 75, 78-83.	0.6	65
222	Explorative signal processing in functional MR imaging. International Journal of Imaging Systems and Technology, 1999, 10, 166-176.	2.7	26
223	Bone marrow scintigraphy using Technetium-99m antigranulocyte antibody in malignant lymphomas. Annals of Oncology, 1999, 10, 79-86.	0.6	6
224	Myo-inositol in depressive and healthy subjects determined by frontal 1H-magnetic resonance spectroscopy at 1.5 tesla. Journal of Psychiatric Research, 1998, 32, 411-420.	1.5	91
225	Absolute metabolite quantification by in vivo NMR spectroscopy: II. a multicentre trial of protocols for in vivo localised proton studies of human brain. Magnetic Resonance Imaging, 1998, 16, 1093-1106.	1.0	98
226	Quantification in Functional Magnetic Resonance Imaging: Fuzzy Clustering vs. Correlation Analysis. Magnetic Resonance Imaging, 1998, 16, 115-125.	1.0	117
227	RETROSPEKTIVE BESTIMMUNG PHYSIOLOGISCHER BEWEGUNGEN IN DER FUNKTIONELLEN KERNSPINTOMOGRAPHIE. Biomedizinische Technik, 1998, 43, 27-28.	0.9	0
228	Comparative diagnostic accuracy of magnetic resonance imaging and immunoscintigraphy for detection of bone marrow involvement in patients with malignant lymphoma.. Journal of Clinical Oncology, 1997, 15, 1754-1760.	0.8	46
229	Magnetoencephalography May Help to Improve Functional MRI Brain Mapping. European Journal of Neuroscience, 1997, 9, 1072-1077.	1.2	39
230	Nonlinear motion artifact reduction in event-triggered gradient-echo FMRI. Magnetic Resonance Imaging, 1997, 15, 163-167.	1.0	3
231	Quantification of signal changes in gradient recalled echo FMRI. Magnetic Resonance Imaging, 1997, 15, 753-762.	1.0	11
232	Modulation of signal changes in gradient-recalled echo functional MRI with increasing echo time correlate with model calculations. Magnetic Resonance Imaging, 1997, 15, 745-752.	1.0	19
233	Three-dimensional reconstruction of the liver venous system using the preservation solution as contrast agent. Journal of Magnetic Resonance Imaging, 1997, 7, 600-602.	1.9	1
234	Fuzzy clustering of gradient-echo functional MRI in the human visual cortex. Part I: Reproducibility. Journal of Magnetic Resonance Imaging, 1997, 7, 1094-1101.	1.9	128

#	ARTICLE	IF	CITATIONS
235	Fuzzy clustering of gradient-echo functional MRI in the human visual cortex. Part II: Quantification. Journal of Magnetic Resonance Imaging, 1997, 7, 1102-1108.	1.9	68
236	On the correlation between tissue hydration state and proton NMR relaxation rates in experimental liver transplantation. , 1997, 10, 143-150.		1
237	Proton NMR relaxation times of human blood samples at 1.5 T and implications for functional MRI. Cellular and Molecular Biology, 1997, 43, 783-91.	0.3	73
238	Improved estimation of tissue hydration and bound water fraction in rat liver tissue. Magnetic Resonance Materials in Physics, Biology, and Medicine, 1996, 4, 55-59.	1.1	15
239	Quantification of statistical type i and ii errors in correlation analysis of simulated functional magnetic resonance imaging data. Magnetic Resonance Materials in Physics, Biology, and Medicine, 1996, 4, 251-256.	1.1	7
240	Reproducibility and postprocessing of gradient-echo functional MRI to improve localization of brain activity in the human visual cortex. Magnetic Resonance Imaging, 1996, 14, 567-579.	1.0	65
241	Proton magnetic resonance spectroscopy in patients with glial tumors: a multicenter study. Journal of Neurosurgery, 1996, 84, 449-458.	0.9	332
242	Quantification of intensity variations in functional MR images using rotated principal components. Physics in Medicine and Biology, 1996, 41, 1425-1438.	1.6	77
243	Evaluating conventional fMRI with respect to noninvasive localization of neuronal activity. International Journal of Imaging Systems and Technology, 1995, 6, 225-229.	2.7	3
244	Comparing Localization of Conventional Functional Magnetic Resonance Imaging and Magnetoencephalography. European Journal of Neuroscience, 1995, 7, 1121-1124.	1.2	67
245	Quality assessment in in vivo NMR spectroscopy: IV. A multicentre trial of test objects and protocols for performance assessment in clinical NMR spectroscopy. Magnetic Resonance Imaging, 1995, 13, 139-157.	1.0	41
246	Temperature- and pH-dependence of proton relaxation rates in rat liver tissue. Magnetic Resonance Imaging, 1995, 13, 429-440.	1.0	26
247	Funktionelles Magnet Resonanz Imaging (fMRI) – Ergebnisse und Kombinationsmöglichkeiten mit der Magneto- und Elektroenzephalographie. Verhandlungen Der Deutschen Gesellschaft für Neurologie, 1995, , 189-192.	0.0	1
248	Clinical proton spectroscopy – towards routine diagnostic use. NMR in Biomedicine, 1994, 7, 341-342.	1.6	4
249	Proton spin-lattice relaxation time as liver transplantation graft viability parameter. Magnetic Resonance Imaging, 1993, 11, 229-239.	1.0	10
250	Proton spin-spin relaxation times as liver transplantation graft viability parameter. Magnetic Resonance Imaging, 1993, 11, 749-759.	1.0	7
251	VII. Analysis of the first international data bank on in vitro NMR relaxation times: Animal liver. Magnetic Resonance Imaging, 1993, 11, 865-872.	1.0	8
252	A possible role of in-flow effects in functional MR-imaging. Magnetic Resonance Materials in Physics, Biology, and Medicine, 1993, 1, 109-113.	1.1	25

#	ARTICLE	IF	CITATIONS
253	Influence of warm ischemia on cold-stored, University of Wisconsin solution-protected rat liver: investigation by 1H-nuclear magnetic resonance relaxometry. Transplantation Proceedings, 1993, 25, 1902-3.	0.3	3
254	COLD-PRESERVED RAT LIVER VIABILITY TESTING BY PROTON NUCLEAR MAGNETIC RESONANCE RELAXOMETRY. Transplantation, 1992, 53, 536-539.	0.5	13
255	Comparison of HTK- and UW-solution for liver preservation tested in an orthotopic liver transplantation model in the pig. Transplant International, 1992, 5, S403-S407.	0.8	16
256	In vitro NMR investigation of controlled single and repeated isoflurane anesthesia. Magnetic Resonance Imaging, 1992, 10, 393-400.	1.0	3
257	Liver tissue characterization by in vitro NMR: Tissue handling and biological variation. Magnetic Resonance in Medicine, 1992, 24, 213-220.	1.9	12
258	Comparison of HTK- and UW-solution for liver preservation tested in an orthotopic liver transplantation model in the pig. , 1992, 5 Suppl 1, 403-407.		6
259	Changes in hepatic tissue water content in EC-, UW-, and HTK-preserved livers tested in a pig liver transplant model. Transplantation Proceedings, 1991, 23, 2414-5.	0.3	3
260	Viability testing by 48-hour 1H-NMR-relaxometry in rat livers preserved in University of Wisconsin solution. Transplantation Proceedings, 1991, 23, 1973.	0.3	0
261	Viability testing of transplantation donor liver by 1H NMR relaxometry. Magnetic Resonance in Medicine, 1990, 16, 173-178.	1.9	12
262	Analysis of relaxation time data from a low-resolution 1H-NMR-pulse-spectrometer. Physiological Chemistry and Physics and Medical NMR, 1989, 21, 123-32.	0.2	2
263	Systematic investigation of degradation effects on spin-spin relaxation times in mouse-liver by low resolution NMR. Physiological Chemistry and Physics and Medical NMR, 1989, 21, 133-44.	0.2	1
264	Systematic investigation of degradation effects on spin-lattice relaxation times in mouse-liver by low resolution NMR. Physiological Chemistry and Physics and Medical NMR, 1989, 21, 73-80.	0.2	1
265	Anisotropy effects in tantalum, niobium, and vanadium down to the millikelvin temperature range. Journal of Low Temperature Physics, 1987, 66, 191-208.	0.6	24
266	Superconductive properties of vanadium and their impurity dependence. Journal of Low Temperature Physics, 1982, 49, 585-607.	0.6	20
267	Superconductivity and Hc2-anisotropy in vanadium. Physica B: Physics of Condensed Matter & C: Atomic, Molecular and Plasma Physics, Optics, 1981, 107, 295-296.	0.9	3
268	Detecting regions of interest in fMRI: an application on exploratory-based data analysis. , 0, , .		1
269	Self-Managed Belief as Part of the "Scientific Method" Part I "A Guide on Mental Modus Operandi as Exemplified by Research in Nuclear Magnetic Resonance. Frontiers in Physics, 0, 6, .	1.0	1